# Exercise 4.1

## Question 1:

Prove the following by using the principle of mathematical induction for all  $n \in \mathbb{N}$ :

$$1+3+3^2+...+3^{n-1}=\frac{\left(3^n-1\right)}{2}$$

### **Answer**

Let the given statement be P(n), i.e.,

P(n): 1 + 3 + 3<sub>2</sub> + ...+ 3<sub>n-1</sub> = 
$$\frac{(3^{n} - 1)}{2}$$

For n = 1, we have

P(1): 1 = 
$$\frac{(3^1-1)}{2} = \frac{3-1}{2} = \frac{2}{2} = 1$$
, which is true.

Let P(k) be true for some positive integer k, i.e.,

$$1+3+3^2+...+3^{k-1} = \frac{\left(3^k-1\right)}{2}$$
 ...(i)

We shall now prove that P(k + 1) is true.

$$1 + 3 + 32 + ... + 3k-1 + 3(k+1)-1$$
  
= (1 + 3 + 3<sub>2</sub> + ... + 3<sub>k-1</sub>) + 3<sub>k</sub>

$$= \frac{(3^{k} - 1)}{2} + 3^{k}$$
 [Using (i)]  

$$= \frac{(3^{k} - 1) + 2 \cdot 3^{k}}{2}$$
  

$$= \frac{(1 + 2)3^{k} - 1}{2}$$
  

$$= \frac{3 \cdot 3^{k} - 1}{2}$$
  

$$= \frac{3^{k+1} - 1}{2}$$

Thus, P(k + 1) is true whenever P(k) is true.

Hence, by the principle of mathematical induction, statement P(n) is true for all natural numbers i.e., n.

## Question 2:

Prove the following by using the principle of mathematical induction for all  $n \in N$ :

$$1^3 + 2^3 + 3^3 + ... + n^3 = \left(\frac{n(n+1)}{2}\right)^2$$

#### Answer

Let the given statement be P(n), i.e.,

P(n): 
$$1^3 + 2^3 + 3^3 + \dots + n^3 = \left(\frac{n(n+1)}{2}\right)^2$$

\_ ` ´ .

For n = 1, we have

P(1): 13 = 1 = 
$$\left( \frac{1 \left( 1+1 \right)}{2} \right)^2 = \left( \frac{1.2}{2} \right)^2 = 1^2 = 1$$
 , which is true.

Let P(k) be true for some positive integer k, i.e.,

$$1^{3} + 2^{3} + 3^{3} + \dots + k^{3} = \left(\frac{k(k+1)}{2}\right)^{2} \qquad \dots (i)$$

We shall now prove that P(k + 1) is true.

Consider

$$1_3 + 2_3 + 3_3 + ... + k_3 + (k + 1)_3$$

Email: care@mentorminutes.com

$$= \left(\frac{k(k+1)}{2}\right)^{2} + (k+1)^{3}$$

$$= \frac{k^{2}(k+1)^{2}}{4} + (k+1)^{3}$$

$$= \frac{k^{2}(k+1)^{2} + 4(k+1)^{3}}{4}$$

$$= \frac{(k+1)^{2}\left\{k^{2} + 4(k+1)\right\}}{4}$$

$$= \frac{(k+1)^{2}\left\{k^{2} + 4k + 4\right\}}{4}$$

$$= \frac{(k+1)^{2}(k+2)^{2}}{4}$$

$$= \frac{(k+1)^{2}(k+1+1)^{2}}{4}$$

$$= \left(\frac{(k+1)(k+1+1)}{2}\right)^{2}$$

=  $(1_3 + 2_3 + 3_3 + .... + k_3) + (k + 1)_3$ 

Thus, P(k + 1) is true whenever P(k) is true.

Hence, by the principle of mathematical induction, statement P(n) is true for all natural numbers i.e., n.

## Question 3:

Prove the following by using the principle of mathematical induction for all  $n \in \mathbb{N}$ :

$$1 + \frac{1}{(1+2)} + \frac{1}{(1+2+3)} + \dots + \frac{1}{(1+2+3+\dots n)} = \frac{2n}{(n+1)}$$

## Answer

Let the given statement be P(n), i.e.,

P(n): 
$$1 + \frac{1}{1+2} + \frac{1}{1+2+3} + \dots + \frac{1}{1+2+3+\dots n} = \frac{2n}{n+1}$$

For n = 1, we have

P(1): 
$$1 = \frac{2.1}{1+1} = \frac{2}{2} = 1$$
 which is true.

Let P(k) be true for some positive integer k, i.e.,



$$1 + \frac{1}{1+2} + \dots + \frac{1}{1+2+3} + \dots + \frac{1}{1+2+3+\dots+k} = \frac{2k}{k+1}$$
 ... (i)

We shall now prove that P(k + 1) is true.

Consider

$$1 + \frac{1}{1+2} + \frac{1}{1+2+3} + \dots + \frac{1}{1+2+3+\dots+k} + \frac{1}{1+2+3+\dots+k+(k+1)}$$

$$= \left(1 + \frac{1}{1+2} + \frac{1}{1+2+3} + \dots + \frac{1}{1+2+3+\dots+k}\right) + \frac{1}{1+2+3+\dots+k+(k+1)}$$

$$= \frac{2k}{k+1} + \frac{1}{1+2+3+\dots+k+(k+1)}$$
[Using (i)]
$$= \frac{2k}{k+1} + \frac{1}{\left(\frac{(k+1)(k+1+1)}{2}\right)}$$

$$= \frac{2k}{(k+1)} + \frac{2}{(k+1)(k+2)}$$

$$= \frac{2}{(k+1)} \left(k + \frac{1}{k+2}\right)$$

### Question 4:

Prove the following by using the principle of mathematical induction for all  $n \in \mathbb{N}$ : 1.2.3

$$\frac{n(n+1)(n+2)(n+3)}{4}$$

#### Answer

Let the given statement be P(n), i.e.,

P(n): 1.2.3 + 2.3.4 + ... + 
$$n(n + 1)$$
 (n + 2) = 
$$\frac{n(n+1)(n+2)(n+3)}{4}$$
For  $n = 1$ , we have

P(1): 1.2.3 = 6 = 
$$\frac{1(1+1)(1+2)(1+3)}{4} = \frac{1.2.3.4}{4} = 6$$
, which is true.

Let P(k) be true for some positive integer k, i.e.,

$$= \frac{k(k+1)(k+2)(k+3)}{4} \dots (i)$$

We shall now prove that P(k + 1) is true.

Consider

$$1.2.3 + 2.3.4 + ... + k(k + 1) (k + 2) + (k + 1) (k + 2) (k + 3)$$

$$= \{1.2.3 + 2.3.4 + ... + k(k + 1) (k + 2)\} + (k + 1) (k + 2) (k + 3)$$

$$= \frac{k(k+1)(k+2)(k+3)}{4} + (k+1)(k+2)(k+3)$$
 [Using (i)]  

$$= (k+1)(k+2)(k+3)\left(\frac{k}{4}+1\right)$$
  

$$= \frac{(k+1)(k+2)(k+3)(k+4)}{4}$$
  

$$= \frac{(k+1)(k+1+1)(k+1+2)(k+1+3)}{4}$$

Thus, P(k + 1) is true whenever P(k) is true.

Hence, by the principle of mathematical induction, statement P(n) is true for all natural numbers i.e., n.

# Question 5:

Prove the following by using the principle of mathematical induction for all  $n \in N$ :

$$1.3 + 2.3^{2} + 3.3^{3} + ... + n.3^{n} = \frac{(2n-1)3^{n+1} + 3}{4}$$

## **Answer**

Let the given statement be P(n), i.e.,

P(n): 
$$1.3 + 2.3^{2} + 3.3^{3} + ... + n3^{n} = \frac{(2n-1)3^{n+1} + 3}{4}$$

For n = 1, we have

P(1): 1.3 = 3 = 
$$\frac{(2.1-1)3^{1+1}+3}{4} = \frac{3^2+3}{4} = \frac{12}{4} = 3$$
, which is true.

Let P(k) be true for some positive integer k, i.e.,

$$1.3 + 2.3^{2} + 3.3^{3} + ... + k3^{k} = \frac{(2k-1)3^{k+1} + 3}{4} \qquad ... (i)$$

We shall now prove that P(k + 1) is true.



Consider

$$1.3 + 2.32 + 3.33 + ... + k3k + (k + 1) 3k + 1$$

$$= (1.3 + 2.32 + 3.33 + ... + k.3k) + (k + 1) 3k + 1$$

$$= \frac{(2k-1)3^{k+1} + 3}{4} + (k+1)3^{k+1}$$

$$= \frac{(2k-1)3^{k+1} + 3 + 4(k+1)3^{k+1}}{4}$$

$$= \frac{3^{k+1} \{2k-1+4(k+1)\} + 3}{4}$$

$$= \frac{3^{k+1} \{6k+3\} + 3}{4}$$

$$= \frac{3^{k+1} \cdot 3\{2k+1\} + 3}{4}$$

$$= \frac{3^{(k+1)+1} \{2k+1\} + 3}{4}$$

$$= \frac{\{2(k+1)-1\}3^{(k+1)+1} + 3}{4}$$

Thus, P(k + 1) is true whenever P(k) is true.

Hence, by the principle of mathematical induction, statement P(n) is true for all natural numbers i.e., n.

# Question 6:

Prove the following by using the principle of mathematical induction for all  $n \in \mathbb{N}$ :

$$1.2 + 2.3 + 3.4 + ... + n.(n+1) = \left\lceil \frac{n(n+1)(n+2)}{3} \right\rceil$$

#### **Answer**

Let the given statement be P(n), i.e.,

$$1.2 + 2.3 + 3.4 + \dots + n.(n+1) = \left[\frac{n(n+1)(n+2)}{3}\right]$$

P(n):

For n = 1, we have

P(1): 
$$1.2 = 2 = \frac{1(1+1)(1+2)}{3} = \frac{1.2.3}{3} = 2$$
, which is true.

Let P(k) be true for some positive integer k, i.e.,

$$1.2 + 2.3 + 3.4 + \dots + k.(k+1) = \left\lceil \frac{k(k+1)(k+2)}{3} \right\rceil \dots (i)$$

We shall now prove that P(k + 1) is true.

Consider

$$1.2 + 2.3 + 3.4 + ... + k.(k + 1) + (k + 1).(k + 2)$$

$$= [1.2 + 2.3 + 3.4 + ... + k.(k + 1)] + (k + 1).(k + 2)$$

$$= \frac{k(k+1)(k+2)}{3} + (k+1)(k+2)$$
 [Using (i)]
$$= (k+1)(k+2)(\frac{k}{3}+1)$$

$$= \frac{(k+1)(k+2)(k+3)}{3}$$

$$= \frac{(k+1)(k+1+1)(k+1+2)}{3}$$

Thus, P(k + 1) is true whenever P(k) is true.

Hence, by the principle of mathematical induction, statement P(n) is true for all natural numbers i.e., n.

### Question 7:

Prove the following by using the principle of mathematical induction for all  $n \in N$ :

$$1.3+3.5+5.7+...+(2n-1)(2n+1)=\frac{n(4n^2+6n-1)}{3}$$

#### Answer

Let the given statement be P(n), i.e.,

P(n): 
$$1.3+3.5+5.7+...+(2n-1)(2n+1)=\frac{n(4n^2+6n-1)}{3}$$

For n = 1, we have

$$P(1):1.3=3=\frac{1(4.1^2+6.1-1)}{3}=\frac{4+6-1}{3}=\frac{9}{3}=3$$
 , which is true.

Let P(k) be true for some positive integer k, i.e.,

$$1.3 + 3.5 + 5.7 + \dots + (2k-1)(2k+1) = \frac{k(4k^2 + 6k - 1)}{3} \dots (i)$$

We shall now prove that P(k + 1) is true.

Consider

$$(1.3 + 3.5 + 5.7 + ... + (2k - 1) (2k + 1) + \{2(k + 1) - 1\}\{2(k + 1) + 1\}$$

$$= \frac{k(4k^2 + 6k - 1)}{3} + (2k + 2 - 1)(2k + 2 + 1) \qquad [Using (i)]$$

$$= \frac{k(4k^2 + 6k - 1)}{3} + (2k + 1)(2k + 3)$$

$$= \frac{k(4k^2 + 6k - 1)}{3} + (4k^2 + 8k + 3)$$

$$= \frac{k(4k^2 + 6k - 1) + 3(4k^2 + 8k + 3)}{3}$$

$$= \frac{4k^3 + 6k^2 - k + 12k^2 + 24k + 9}{3}$$

$$= \frac{4k^3 + 18k^2 + 23k + 9}{3}$$

$$= \frac{4k^3 + 14k^2 + 9k + 4k^2 + 14k + 9}{3}$$

$$= \frac{k(4k^2 + 14k + 9) + 1(4k^2 + 14k + 9)}{3}$$

$$= \frac{(k + 1)(4k^2 + 14k + 9)}{3}$$

$$= \frac{(k + 1)\{4(k^2 + 2k + 1) + 6(k + 1) - 1\}}{3}$$

$$= \frac{(k + 1)\{4(k^2 + 2k + 1) + 6(k + 1) - 1\}}{3}$$

$$= \frac{(k + 1)\{4(k + 1)^2 + 6(k + 1) - 1\}}{3}$$

Thus, P(k + 1) is true whenever P(k) is true.

Hence, by the principle of mathematical induction, statement P(n) is true for all natural numbers i.e., n.

### Question 8:

Prove the following by using the principle of mathematical induction for all  $n \in \mathbb{N}$ : 1.2 +

$$2.2_2 + 3.2_2 + ... + n.2_n = (n-1) 2_{n+1} + 2$$

#### **Answer**

Let the given statement be P(n), i.e.,

$$P(n)$$
: 1.2 + 2.2<sub>2</sub> + 3.2<sub>2</sub> + ... +  $n$ .2<sub>n</sub> =  $(n - 1)$  2<sub>n+1</sub> + 2

For n = 1, we have

P(1): 
$$1.2 = 2 = (1 - 1) 2_{1+1} + 2 = 0 + 2 = 2$$
, which is true.

Let P(k) be true for some positive integer k, i.e.,

$$1.2 + 2.22 + 3.22 + ... + k \cdot 2k = (k - 1) \cdot 2k + 1 + 2 ... (i)$$

We shall now prove that P(k + 1) is true.

Consider

$$\begin{aligned}
&\left\{1.2 + 2.2^{2} + 3.2^{3} + \dots + k.2^{k}\right\} + (k+1) \cdot 2^{k+1} \\
&= (k-1)2^{k+1} + 2 + (k+1)2^{k+1} \\
&= 2^{k+1} \left\{ (k-1) + (k+1) \right\} + 2 \\
&= 2^{k+1} \cdot 2k + 2 \\
&= k \cdot 2^{(k+1)+1} + 2 \\
&= \left\{ (k+1) - 1 \right\} 2^{(k+1)+1} + 2
\end{aligned}$$

Thus, P(k + 1) is true whenever P(k) is true.

Hence, by the principle of mathematical induction, statement P(n) is true for all natural numbers i.e., n.

### Question 9:

Prove the following by using the principle of mathematical induction for all  $n \in N$ :

$$\frac{1}{2} + \frac{1}{4} + \frac{1}{8} + \dots + \frac{1}{2^n} = 1 - \frac{1}{2^n}$$

### **Answer**

Let the given statement be P(n), i.e.,

P(n): 
$$\frac{1}{2} + \frac{1}{4} + \frac{1}{8} + \dots + \frac{1}{2^n} = 1 - \frac{1}{2^n}$$

For n = 1, we have

P(1): 
$$\frac{1}{2} = 1 - \frac{1}{2^1} = \frac{1}{2}$$
, which is true.

Let P(k) be true for some positive integer k, i.e.,

$$\frac{1}{2} + \frac{1}{4} + \frac{1}{8} + \dots + \frac{1}{2^k} = 1 - \frac{1}{2^k}$$
 ... (i)

We shall now prove that P(k + 1) is true.

Consider

$$\begin{split} &\left(\frac{1}{2} + \frac{1}{4} + \frac{1}{8} + \dots + \frac{1}{2^k}\right) + \frac{1}{2^{k+1}} \\ &= \left(1 - \frac{1}{2^k}\right) + \frac{1}{2^{k+1}} \\ &= 1 - \frac{1}{2^k} + \frac{1}{2 \cdot 2^k} \\ &= 1 - \frac{1}{2^k} \left(1 - \frac{1}{2}\right) \\ &= 1 - \frac{1}{2^k} \left(\frac{1}{2}\right) \\ &= 1 - \frac{1}{2^{k+1}} \end{split}$$
 [Using (i)]

Thus, P(k + 1) is true whenever P(k) is true.

Hence, by the principle of mathematical induction, statement P(n) is true for all natural numbers i.e., n.

### Question 10:

Prove the following by using the principle of mathematical induction for all  $n \in N$ :

$$\frac{1}{2.5} + \frac{1}{5.8} + \frac{1}{8.11} + \dots + \frac{1}{(3n-1)(3n+2)} = \frac{n}{(6n+4)}$$

# Answer

Let the given statement be P(n), i.e.,

$$\frac{1}{2.5} + \frac{1}{5.8} + \frac{1}{8.11} + \dots + \frac{1}{(3n-1)(3n+2)} = \frac{n}{(6n+4)}$$

For n = 1, we have

Email: care@mentorminutes.com



$$P(1) = \frac{1}{2.5} = \frac{1}{10} = \frac{1}{6.1 + 4} = \frac{1}{10}$$
, which is true.

Let P(k) be true for some positive integer k, i.e.,

$$\frac{1}{2.5} + \frac{1}{5.8} + \frac{1}{8.11} + \dots + \frac{1}{(3k-1)(3k+2)} = \frac{k}{6k+4}$$
 ... (i)

We shall now prove that P(k + 1) is true.

Consider

$$\frac{1}{2.5} + \frac{1}{5.8} + \frac{1}{8.11} + \dots + \frac{1}{(3k-1)(3k+2)} + \frac{1}{\{3(k+1)-1\}\{3(k+1)+2\}}$$

$$= \frac{k}{6k+4} + \frac{1}{(3k+2)(3k+5)}$$

$$= \frac{k}{2(3k+2)} + \frac{1}{(3k+2)(3k+5)}$$

$$= \frac{1}{(3k+2)} \left(\frac{k}{2} + \frac{1}{3k+5}\right)$$

$$= \frac{1}{(3k+2)} \left(\frac{k(3k+5)+2}{2(3k+5)}\right)$$

$$= \frac{1}{(3k+2)} \left(\frac{3k^2+5k+2}{2(3k+5)}\right)$$

$$= \frac{1}{(3k+2)} \left(\frac{(3k+2)(k+1)}{2(3k+5)}\right)$$

$$= \frac{(k+1)}{6k+10}$$

$$= \frac{(k+1)}{6(k+1)+4}$$

Thus, P(k + 1) is true whenever P(k) is true.

Hence, by the principle of mathematical induction, statement P(n) is true for all natural numbers i.e., n.



# Question 11:

Prove the following by using the principle of mathematical induction for all  $n \in N$ :

$$\frac{1}{1.2.3} + \frac{1}{2.3.4} + \frac{1}{3.4.5} + \dots + \frac{1}{n(n+1)(n+2)} = \frac{n(n+3)}{4(n+1)(n+2)}$$

#### **Answer**

Let the given statement be P(n), i.e.,

$$\frac{1}{1.2.3} + \frac{1}{2.3.4} + \frac{1}{3.4.5} + \dots + \frac{1}{n(n+1)(n+2)} = \frac{n(n+3)}{4(n+1)(n+2)}$$

For n = 1, we have

$$P(1): \frac{1}{1 \cdot 2 \cdot 3} = \frac{1 \cdot (1+3)}{4(1+1)(1+2)} = \frac{1 \cdot 4}{4 \cdot 2 \cdot 3} = \frac{1}{1 \cdot 2 \cdot 3}$$
, which is true.

Let P(k) be true for some positive integer k, i.e.,

$$\frac{1}{1 \cdot 2 \cdot 3} + \frac{1}{2 \cdot 3 \cdot 4} + \frac{1}{3 \cdot 4 \cdot 5} + \dots + \frac{1}{k(k+1)(k+2)} = \frac{k(k+3)}{4(k+1)(k+2)}$$
 ... (i)

We shall now prove that P(k + 1) is true.

$$\left[\frac{1}{1 \cdot 2 \cdot 3} + \frac{1}{2 \cdot 3 \cdot 4} + \frac{1}{3 \cdot 4 \cdot 5} + \dots + \frac{1}{k(k+1)(k+2)}\right] + \frac{1}{(k+1)(k+2)(k+3)}$$

$$= \frac{k(k+3)}{4(k+1)(k+2)} + \frac{1}{(k+1)(k+2)(k+3)} \qquad [Using (i)]$$

$$= \frac{1}{(k+1)(k+2)} \left\{ \frac{k(k+3)}{4} + \frac{1}{k+3} \right\}$$

$$= \frac{1}{(k+1)(k+2)} \left\{ \frac{k(k+3)^2 + 4}{4(k+3)} \right\}$$

$$= \frac{1}{(k+1)(k+2)} \left\{ \frac{k(k^2 + 6k + 9) + 4}{4(k+3)} \right\}$$

$$= \frac{1}{(k+1)(k+2)} \left\{ \frac{k^3 + 6k^2 + 9k + 4}{4(k+3)} \right\}$$

$$= \frac{1}{(k+1)(k+2)} \left\{ \frac{k^3 + 2k^2 + k + 4k^2 + 8k + 4}{4(k+3)} \right\}$$

$$= \frac{1}{(k+1)(k+2)} \left\{ \frac{k(k^2 + 2k + 1) + 4(k^2 + 2k + 1)}{4(k+3)} \right\}$$

$$= \frac{1}{(k+1)(k+2)} \left\{ \frac{k(k+1)^2 + 4(k+1)^2}{4(k+3)} \right\}$$

$$= \frac{(k+1)^2(k+4)}{4(k+1)(k+2)(k+3)}$$

$$= \frac{(k+1)^2(k+4)}{4(k+1)(k+2)(k+3)}$$

$$= \frac{(k+1)^2(k+4)}{4(k+1)(k+2)(k+3)}$$

Thus, P(k + 1) is true whenever P(k) is true.

Hence, by the principle of mathematical induction, statement P(n) is true for all natural numbers i.e., n.

### Question 12:

Prove the following by using the principle of mathematical induction for all  $n \in \mathbb{N}$ :

$$a + ar + ar^{2} + ... + ar^{n-1} = \frac{a(r^{n} - 1)}{r - 1}$$

#### **Answer**

Let the given statement be P(n), i.e.,

$$P(n): a + ar + ar^2 + ... + ar^{n-1} = \frac{a(r^n - 1)}{r - 1}$$

For n = 1, we have

$$\mathbf{P}\!\left(1\right)\!:\!a=\!\frac{a\!\left(r^1\!-\!1\right)}{\left(r\!-\!1\right)}\!=\!a$$
 , which is true.

Let P(k) be true for some positive integer k, i.e.,

$$a + ar + ar^{2} + \dots + ar^{k-1} = \frac{a(r^{k} - 1)}{r - 1}$$
 ... (i)

We shall now prove that P(k + 1) is true.

Consider

Thus, P(k + 1) is true whenever P(k) is true.

Hence, by the principle of mathematical induction, statement P(n) is true for all natural numbers i.e., n.

### Question 13:

Prove the following by using the principle of mathematical induction for all  $n \in \mathbb{N}$ :

$$\left(1+\frac{3}{1}\right)\left(1+\frac{5}{4}\right)\left(1+\frac{7}{9}\right)...\left(1+\frac{(2n+1)}{n^2}\right)=\left(n+1\right)^2$$

#### **Answer**

Let the given statement be P(n), i.e.,

$$P(n): \left(1 + \frac{3}{1}\right) \left(1 + \frac{5}{4}\right) \left(1 + \frac{7}{9}\right) ... \left(1 + \frac{(2n+1)}{n^2}\right) = (n+1)^2$$

For n = 1, we have

$$P(1): (1+\frac{3}{1})=4=(1+1)^2=2^2=4$$
, which is true.

Let P(k) be true for some positive integer k, i.e.,

$$\left(1 + \frac{3}{1}\right)\left(1 + \frac{5}{4}\right)\left(1 + \frac{7}{9}\right) \dots \left(1 + \frac{(2k+1)}{k^2}\right) = (k+1)^2 \dots (1)$$

We shall now prove that P(k + 1) is true.

Consider

$$\left[ \left( 1 + \frac{3}{1} \right) \left( 1 + \frac{5}{4} \right) \left( 1 + \frac{7}{9} \right) \dots \left( 1 + \frac{(2k+1)}{k^2} \right) \right] \left\{ 1 + \frac{\{2(k+1)+1\}}{(k+1)^2} \right\} 
= (k+1)^2 \left( 1 + \frac{2(k+1)+1}{(k+1)^2} \right) \qquad \left[ \text{Using}(1) \right] 
= (k+1)^2 \left[ \frac{(k+1)^2 + 2(k+1)+1}{(k+1)^2} \right] 
= (k+1)^2 + 2(k+1)+1 
= \{(k+1)+1\}^2$$

Thus, P(k + 1) is true whenever P(k) is true.

Hence, by the principle of mathematical induction, statement P(n) is true for all natural numbers i.e., n.

### Question 14:

Prove the following by using the principle of mathematical induction for all  $n \in \mathbb{N}$ :

$$\left(1+\frac{1}{1}\right)\left(1+\frac{1}{2}\right)\left(1+\frac{1}{3}\right)...\left(1+\frac{1}{n}\right)=(n+1)$$

#### Answer

Let the given statement be P(n), i.e.,

$$P(n): \left(1+\frac{1}{1}\right)\left(1+\frac{1}{2}\right)\left(1+\frac{1}{3}\right)...\left(1+\frac{1}{n}\right) = (n+1)$$

For n = 1, we have

$$P\left(1\right)\!:\!\left(1\!+\!\frac{1}{1}\right)\!=2=\left(1\!+\!1\right)$$
 , which is true.

Let P(k) be true for some positive integer k, i.e.,

$$P(k): \left(1 + \frac{1}{1}\right) \left(1 + \frac{1}{2}\right) \left(1 + \frac{1}{3}\right) \dots \left(1 + \frac{1}{k}\right) = (k+1)$$
 ... (1)

We shall now prove that P(k + 1) is true.

Consider

$$\left[ \left( 1 + \frac{1}{1} \right) \left( 1 + \frac{1}{2} \right) \left( 1 + \frac{1}{3} \right) \dots \left( 1 + \frac{1}{k} \right) \right] \left( 1 + \frac{1}{k+1} \right) \\
= (k+1) \left( 1 + \frac{1}{k+1} \right) \qquad \left[ \text{Using (1)} \right] \\
= (k+1) \left( \frac{(k+1)+1}{(k+1)} \right) \\
= (k+1)+1$$

Thus, P(k + 1) is true whenever P(k) is true.

Hence, by the principle of mathematical induction, statement P(n) is true for all natural numbers i.e., n.

### Question 15:

Prove the following by using the principle of mathematical induction for all  $n \in N$ :

$$1^{2} + 3^{2} + 5^{2} + ... + (2n-1)^{2} = \frac{n(2n-1)(2n+1)}{3}$$

#### **Answer**

Let the given statement be P(n), i.e.,

$$P(n) = 1^2 + 3^2 + 5^2 + \dots + (2n-1)^2 = \frac{n(2n-1)(2n+1)}{3}$$

For n = 1, we have

$$P(1) = 1^2 = 1 = \frac{1(2.1-1)(2.1+1)}{3} = \frac{1.1.3}{3} = 1$$
, which is true.

Let P(k) be true for some positive integer k, i.e.,

$$P(k) = 1^2 + 3^2 + 5^2 + \dots + (2k-1)^2 = \frac{k(2k-1)(2k+1)}{3}$$
 ... (1)

We shall now prove that P(k + 1) is true.

$$\begin{cases}
1^{2} + 3^{2} + 5^{2} + \dots + (2k-1)^{2} \\
= \frac{k(2k-1)(2k+1)}{3} + (2k+2-1)^{2} \\
= \frac{k(2k-1)(2k+1)}{3} + (2k+1)^{2} \\
= \frac{k(2k-1)(2k+1)}{3} + (2k+1)^{2} \\
= \frac{k(2k-1)(2k+1) + 3(2k+1)^{2}}{3} \\
= \frac{(2k+1)\{k(2k-1) + 3(2k+1)\}}{3} \\
= \frac{(2k+1)\{2k^{2} - k + 6k + 3\}}{3}$$

$$= \frac{(2k+1)\{2k^2+5k+3\}}{3}$$

$$= \frac{(2k+1)\{2k^2+2k+3k+3\}}{3}$$

$$= \frac{(2k+1)\{2k(k+1)+3(k+1)\}}{3}$$

$$= \frac{(2k+1)(k+1)(2k+3)}{3}$$

$$= \frac{(2k+1)\{2(k+1)-1\}\{2(k+1)+1\}}{3}$$

Thus, P(k + 1) is true whenever P(k) is true.

Hence, by the principle of mathematical induction, statement P(n) is true for all natural numbers i.e., n.

### Question 16:

Prove the following by using the principle of mathematical induction for all  $n \in \mathbb{N}$ :

$$\frac{1}{1.4} + \frac{1}{4.7} + \frac{1}{7.10} + \dots + \frac{1}{(3n-2)(3n+1)} = \frac{n}{(3n+1)}$$

### **Answer**

Let the given statement be P(n), i.e.,

$$P(n): \frac{1}{1.4} + \frac{1}{4.7} + \frac{1}{7.10} + \dots + \frac{1}{(3n-2)(3n+1)} = \frac{n}{(3n+1)}$$

For n = 1, we have

$$P(1) = \frac{1}{1.4} = \frac{1}{3.1+1} = \frac{1}{4} = \frac{1}{1.4}$$
, which is true.

Let P(k) be true for some positive integer k, i.e.,

$$P(k) = \frac{1}{1.4} + \frac{1}{4.7} + \frac{1}{7.10} + \dots + \frac{1}{(3k-2)(3k+1)} = \frac{k}{3k+1}$$
 ... (1)

We shall now prove that P(k + 1) is true.

$$\left\{ \frac{1}{1.4} + \frac{1}{4.7} + \frac{1}{7.10} + \dots + \frac{1}{(3k-2)(3k+1)} \right\} + \frac{1}{\{3(k+1)-2\}\{3(k+1)+1\}}$$

$$= \frac{k}{3k+1} + \frac{1}{(3k+1)(3k+4)} \qquad [Using (1)]$$

$$= \frac{1}{(3k+1)} \left\{ k + \frac{1}{(3k+4)} \right\}$$

$$= \frac{1}{(3k+1)} \left\{ \frac{k(3k+4)+1}{(3k+4)} \right\}$$

$$= \frac{1}{(3k+1)} \left\{ \frac{3k^2 + 4k + 1}{(3k+4)} \right\}$$

$$= \frac{1}{(3k+1)} \left\{ \frac{3k^2 + 3k + k + 1}{(3k+4)} \right\}$$

$$= \frac{(3k+1)(k+1)}{(3k+1)(3k+4)}$$

$$= \frac{(k+1)}{3(k+1)+1}$$

Thus, P(k + 1) is true whenever P(k) is true.

Hence, by the principle of mathematical induction, statement P(n) is true for all natural numbers i.e., n.

## Question 17:

Prove the following by using the principle of mathematical induction for all  $n \in \mathbb{N}$ :

$$\frac{1}{3.5} + \frac{1}{5.7} + \frac{1}{7.9} + \dots + \frac{1}{(2n+1)(2n+3)} = \frac{n}{3(2n+3)}$$

#### Answer

Let the given statement be P(n), i.e.,

$$P(n): \frac{1}{3.5} + \frac{1}{5.7} + \frac{1}{7.9} + \dots + \frac{1}{(2n+1)(2n+3)} = \frac{n}{3(2n+3)}$$

For n = 1, we have

$$P(1): \frac{1}{3.5} = \frac{1}{3(2.1+3)} = \frac{1}{3.5}$$
, which is true.

Let P(k) be true for some positive integer k, i.e.,

$$P(k): \frac{1}{3.5} + \frac{1}{5.7} + \frac{1}{7.9} + \dots + \frac{1}{(2k+1)(2k+3)} = \frac{k}{3(2k+3)}$$
 ... (1)

We shall now prove that P(k + 1) is true.

Consider

$$\left[\frac{1}{3.5} + \frac{1}{5.7} + \frac{1}{7.9} + \dots + \frac{1}{(2k+1)(2k+3)}\right] + \frac{1}{\{2(k+1)+1\}\{2(k+1)+3\}}$$

$$= \frac{k}{3(2k+3)} + \frac{1}{(2k+3)(2k+5)} \qquad [Using (1)]$$

$$= \frac{1}{(2k+3)} \left[\frac{k}{3} + \frac{1}{(2k+5)}\right]$$

$$= \frac{1}{(2k+3)} \left[\frac{k(2k+5)+3}{3(2k+5)}\right]$$

$$= \frac{1}{(2k+3)} \left[\frac{2k^2+5k+3}{3(2k+5)}\right]$$

$$= \frac{1}{(2k+3)} \left[\frac{2k^2+2k+3k+3}{3(2k+5)}\right]$$

$$= \frac{1}{(2k+3)} \left[\frac{2k(k+1)+3(k+1)}{3(2k+5)}\right]$$

$$= \frac{(k+1)(2k+3)}{3(2k+3)(2k+5)}$$

$$= \frac{(k+1)}{3\{2(k+1)+3\}}$$

Thus, P(k + 1) is true whenever P(k) is true.

Hence, by the principle of mathematical induction, statement P(n) is true for all natural numbers i.e., n.

### Question 18:

 $1 < \frac{1}{8}(2.1+1)^2 = \frac{9}{8}$ 

Prove the following by using the principle of mathematical induction for all  $n \in N$ :

$$1+2+3+...+n < \frac{1}{8}(2n+1)^2$$

#### **Answer**

Let the given statement be P(n), i.e.,

$$P(n): 1+2+3+...+n < \frac{1}{8}(2n+1)^2$$

It can be noted that P(n) is true for n = 1 since Let P(k) be true for some positive integer k, i.e.,

$$1+2+...+k < \frac{1}{8}(2k+1)^2$$
 ... (1)

We shall now prove that P(k + 1) is true whenever P(k) is true.

Consider

$$(1+2+...+k)+(k+1)<\frac{1}{8}(2k+1)^{2}+(k+1)$$

$$<\frac{1}{8}\{(2k+1)^{2}+8(k+1)\}$$

$$<\frac{1}{8}\{4k^{2}+4k+1+8k+8\}$$

$$<\frac{1}{8}\{4k^{2}+12k+9\}$$

$$<\frac{1}{8}(2k+3)^{2}$$

$$<\frac{1}{8}\{2(k+1)+1\}^{2}$$

Hence, 
$$(1+2+3+...+k)+(k+1)<\frac{1}{8}(2k+1)^2+(k+1)$$

Thus, P(k + 1) is true whenever P(k) is true.

Hence, by the principle of mathematical induction, statement P(n) is true for all natural numbers i.e., n.

### Question 19:

Prove the following by using the principle of mathematical induction for all  $n \in \mathbb{N}$ : n (n + 1) (n + 5) is a multiple of 3.

#### **Answer**

Let the given statement be P(n), i.e.,

P(n): n(n + 1)(n + 5), which is a multiple of 3.

It can be noted that P(n) is true for n = 1 since 1 (1 + 1) (1 + 5) = 12, which is a multiple of 3.

Let P(k) be true for some positive integer k, i.e.,

k(k+1)(k+5) is a multiple of 3.

: 
$$k(k + 1)(k + 5) = 3m$$
, where  $m \in \mathbb{N}$  ... (1)

We shall now prove that P(k + 1) is true whenever P(k) is true.

Consider

$$(k+1)\{(k+1)+1\}\{(k+1)+5\}$$

$$= (k+1)(k+2)\{(k+5)+1\}$$

$$= (k+1)(k+2)(k+5)+(k+1)(k+2)$$

$$= \{k(k+1)(k+5)+2(k+1)(k+5)\}+(k+1)(k+2)$$

$$= 3m+(k+1)\{2(k+5)+(k+2)\}$$

$$= 3m+(k+1)\{2k+10+k+2\}$$

$$= 3m+(k+1)(3k+12)$$

$$= 3m+3(k+1)(k+4)$$

$$= 3\{m+(k+1)(k+4)\} = 3\times q, \text{ where } q = \{m+(k+1)(k+4)\} \text{ is some natural number}$$
Therefore,  $(k+1)\{(k+1)+1\}\{(k+1)+5\}$  is a multiple of 3.

Thus, P(k + 1) is true whenever P(k) is true.

Hence, by the principle of mathematical induction, statement P(n) is true for all natural numbers i.e., n.

#### Question 20:

Prove the following by using the principle of mathematical induction for all  $n \in \mathbb{N}$ :  $10_{2n-1} + 1$  is divisible by 11.

# Answer

Let the given statement be P(n), i.e.,

P(n): 10<sub>2n-1</sub> + 1 is divisible by 11.

It can be observed that P(n) is true for n = 1 since  $P(1) = 10_{2.1 - 1} + 1 = 11$ , which is divisible by 11.

Let P(k) be true for some positive integer k, i.e.,

 $10_{2k-1} + 1$  is divisible by 11.

∴102k - 1 + 1 = 11
$$m$$
, where  $m \in \mathbb{N}$  ... (1)

We shall now prove that P(k + 1) is true whenever P(k) is true.

#### Consider

$$10^{2(k+1)-1} + 1$$

$$= 10^{2k+2-1} + 1$$

$$= 10^{2k+1} + 1$$

$$= 10^{2} (10^{2k-1} + 1 - 1) + 1$$

$$= 10^{2} (10^{2k-1} + 1) - 10^{2} + 1$$

$$= 10^{2} .11m - 100 + 1 \qquad [Using (1)]$$

$$= 100 \times 11m - 99$$

$$= 11(100m - 9)$$

$$= 11r, \text{ where } r = (100m - 9) \text{ is some natural number}$$
Therefore,  $10^{2(k+1)-1} + 1$  is divisible by 11.

Thus, P(k + 1) is true whenever P(k) is true.

Hence, by the principle of mathematical induction, statement P(n) is true for all natural numbers i.e., n.

#### Question 21:

Prove the following by using the principle of mathematical induction for all  $n \in \mathbb{N}$ :  $x_{2n} - y_{2n}$  is divisible by x + y.

# Answer

Let the given statement be P(n), i.e.,

P(n):  $x_{2n} - y_{2n}$  is divisible by x + y.

It can be observed that P(n) is true for n = 1.

This is so because  $x_2 \times 1 - y_2 \times 1 = x_2 - y_2 = (x + y)(x - y)$  is divisible by (x + y).

Let P(k) be true for some positive integer k, i.e.,

 $x_{2k} - y_{2k}$  is divisible by x + y.

$$\therefore x_{2k} - y_{2k} = m(x + y)$$
, where  $m \in \mathbb{N}$  ... (1)

We shall now prove that P(k + 1) is true whenever P(k) is true.

#### Consider

$$x^{2(k+1)} - y^{2(k+1)}$$

$$= x^{2k} \cdot x^2 - y^{2k} \cdot y^2$$

$$= x^2 \left( x^{2k} - y^{2k} + y^{2k} \right) - y^{2k} \cdot y^2$$

$$= x^2 \left\{ m(x+y) + y^{2k} \right\} - y^{2k} \cdot y^2 \qquad \left[ \text{Using (1)} \right]$$

$$= m(x+y)x^2 + y^{2k} \cdot x^2 - y^{2k} \cdot y^2$$

$$= m(x+y)x^2 + y^{2k} \left( x^2 - y^2 \right)$$

$$= m(x+y)x^2 + y^{2k} \left( x + y \right) (x-y)$$

$$= (x+y) \left\{ mx^2 + y^{2k} \left( x - y \right) \right\}, \text{ which is a factor of } (x+y).$$

Thus, P(k + 1) is true whenever P(k) is true.

Hence, by the principle of mathematical induction, statement P(n) is true for all natural numbers i.e., n.

### Question 22:

Prove the following by using the principle of mathematical induction for all  $n \in \mathbb{N}$ : 32n + 2 - 8n - 9 is divisible by 8.

#### **Answer**

Let the given statement be P(n), i.e.,

P(n): 32n + 2 - 8n - 9 is divisible by 8.

It can be observed that P(n) is true for n = 1 since  $32 \times 1 + 2 - 8 \times 1 - 9 = 64$ , which is divisible by 8.

Let P(k) be true for some positive integer k, i.e.,

32k + 2 - 8k - 9 is divisible by 8.

∴
$$32k + 2 - 8k - 9 = 8m$$
; where  $m \in \mathbb{N}$  ... (1)

We shall now prove that P(k + 1) is true whenever P(k) is true.

Class XI



$$3^{2(k+1)+2} - 8(k+1) - 9$$

$$= 3^{2k+2} \cdot 3^2 - 8k - 8 - 9$$

$$= 3^2 (3^{2k+2} - 8k - 9 + 8k + 9) - 8k - 17$$

$$= 3^2 (3^{2k+2} - 8k - 9) + 3^2 (8k + 9) - 8k - 17$$

$$= 9.8m + 9(8k + 9) - 8k - 17$$

$$= 9.8m + 72k + 81 - 8k - 17$$

$$= 9.8m + 64k + 64$$

$$= 8(9m + 8k + 8)$$

$$= 8r, \text{ where } r = (9m + 8k + 8) \text{ is a natural number}$$
Therefore,  $3^{2(k+1)+2} - 8(k+1) - 9$  is divisible by 8.

Thus, P(k + 1) is true whenever P(k) is true.

Hence, by the principle of mathematical induction, statement P(n) is true for all natural numbers i.e., n.

## Question 23:

Prove the following by using the principle of mathematical induction for all  $n \in N$ :  $41_n - 14_n$  is a multiple of 27.

### **Answer**

Let the given statement be P(n), i.e.,

 $P(n):41_n - 14_n$  is a multiple of 27.

It can be observed that P(n) is true for n = 1 since 27.

 $41^1 - 14^1 = 27$ , which is a multiple of

Let P(k) be true for some positive integer k, i.e.,

41k - 14k is a multiple of 27

∴41k - 14k = 27m, where  $m \in \mathbb{N}$  ... (1)

We shall now prove that P(k + 1) is true whenever P(k) is true.

$$41^{k+1} - 14^{k+1}$$

$$= 41^{k} \cdot 41 - 14^{k} \cdot 14$$

$$= 41(41^{k} - 14^{k} + 14^{k}) - 14^{k} \cdot 14$$

$$= 41(41^{k} - 14^{k}) + 41.14^{k} - 14^{k} \cdot 14$$

$$= 41.27m + 14^{k} (41 - 14)$$

$$= 41.27m + 27.14^{k}$$

$$= 27(41m - 14^{k})$$

=  $27 \times r$ , where  $r = (41m - 14^k)$  is a natural number

Therefore,  $41^{k+1} - 14^{k+1}$  is a multiple of 27.

Thus, P(k + 1) is true whenever P(k) is true.

Hence, by the principle of mathematical induction, statement P(n) is true for all natural numbers i.e., n.

## Question 24:

Prove the following by using the principle of mathematical induction for all  $n \in \mathbb{N}$ :  $(2n+7) < (n+3)_2$ 

### Answer

Let the given statement be P(n), i.e.,


$$P(n)$$
:  $(2n + 7) < (n + 3)_2$ 

It can be observed that P(n) is true for n = 1 since  $2.1 + 7 = 9 < (1 + 3)_2 = 16$ , which is true.

Let P(k) be true for some positive integer k, i.e.,

$$(2k + 7) < (k + 3)_2 \dots (1)$$

We shall now prove that P(k + 1) is true whenever P(k) is true.



Mathematics

$$\begin{aligned} & \left\{ 2\left(k+1\right)+7\right\} = \left(2k+7\right)+2 \\ & \therefore \left\{ 2\left(k+1\right)+7\right\} = \left(2k+7\right)+2 < \left(k+3\right)^2+2 \\ & \left[ u \sin g \ (1) \right] \\ & 2\left(k+1\right)+7 < k^2+6k+9+2 \\ & 2\left(k+1\right)+7 < k^2+6k+11 \\ & \text{Now, } k^2+6k+11 < k^2+8k+16 \\ & \therefore 2\left(k+1\right)+7 < \left(k+4\right)^2 \\ & 2\left(k+1\right)+7 < \left\{ \left(k+1\right)+3 \right\}^2 \end{aligned}$$

Thus, P(k + 1) is true whenever P(k) is true.

Hence, by the principle of mathematical induction, statement P(n) is true for all natural numbers i.e., n.