



(Division of Aakash Educational Services Ltd.)

Regd. Office: Aakash Tower, Plot No.-4, Sec-11, MLU, Dwarka, New Delhi-110075 Ph.: 011-47623456 Fax: 011-47623472

HINTS & SOLUTIONS for CBSE Preliminary 2011

by Aakash Institute

- 1. The dimensions of $(\mu_0 \epsilon_0)^{-\frac{1}{2}}$ are
 - $(1) \quad \left[L^{-\frac{1}{2}} T^{\frac{1}{2}} \right]$

(2) $\left[L^{\frac{1}{2}} T^{-\frac{1}{2}} \right]$

(3) $[L^{-1}T]$

(4) $[LT^{-1}]$

Sol. Answer (4)

Given expression is that of speed of light.

- 2. The moment of inertia of a thin uniform rod of mass M and length L about an axis passing through its midpoint and perpendicular to its length is I_0 . Its moment of inertia about an axis passing through one of its ends and perpendicular to its length is
 - (1) $I_0 + ML^2$

 $(2) I_0 + \frac{ML^2}{2}$

(3) $I_0 + \frac{ML^2}{4}$

(4) $I_0 + 2ML^2$

Sol. Answer (3)

$$I = I_{CM} + Mh^2$$

(Parallel axis theorem)

- 3. A boy standing at the top of a tower of 20 m height drops a stone. Assuming $g = 10 \text{ ms}^{-2}$, the velocity with which it hits the ground is
 - (1) 5.0 m/s

(2) 10.0 m/s

(3) 20.0 m/s

(4) 40.0 m/s

Sol. Answer (3)

$$v = \sqrt{2gh} = \sqrt{2 \times 10 \times 20} = 20 \text{ m/s}$$

- 4. A person of mass 60 kg is inside a lift of mass 940 kg and presses the button one control panel. The lift starts moving upwards with an acceleration 1.0 m/s^2 . If $g = 10 \text{ ms}^{-2}$, the tension in the supporting cable is
 - (1) 1200 N

(2) 8600 N

(3) 9680 N

(4) 11000 N

Sol. Answer (4)

$$T = (M + m) (g + a) = (940 + 60) (10 + 1) = 11000 N$$

- 5. A body projected vertically from the earth reaches a height equal to earth's radius before returning to the earth. The power exerted by the gravitational force is greatest
 - (1) At the instant just after the body is projected
 - (2) At the highest position of the body
 - (3) At the instant just before the body hits the earth
 - (4) It remains constant all through

Sol. Answer (3)

$$P = \vec{F} \cdot \vec{v} = Fv \cos \theta$$

Just before hitting θ is zero and both F, v are maximum.

6. The instantaneous angular position of a point on a rotating wheel is given by the equation $\theta(t) = 2t^3 - 6t^2$. The torque on the wheel becomes zero at

(1)
$$t = 2 s$$

(2)
$$t = 1 \text{ s}$$

(3)
$$t = 0.2 \text{ s}$$

(4)
$$t = 0.25 \text{ s}$$

Sol. Answer (2)

Torque zero means, $\,\alpha\,$ zero

$$\therefore \frac{d^2\theta}{dt^2} = 0$$

$$\Rightarrow$$
 12t - 12 = 0

$$t = 1$$
 second

7. A particle moves in a circle of radius 5 cm with constant speed and time period 0.2 π s. The acceleration of the particle is

(1)
$$5 \text{ m/s}^2$$

Sol. Answer (1)

$$a = \omega^2 R = \left(\frac{2\pi}{0.2 \,\pi}\right)^2 \left(5 \times 10^{-2}\right) = 5 \text{ m/s}^2$$

- 8. A body of mass *M* hits normally a rigid wall with velocity *V* and bounces back with the same velocity. The impulse experienced by the body is
 - (1) Zero

(2) MV

(3) 1.5 MV

(4) 2 MV

Sol. Answer (4)

Impulse = MV - (-MV) = 2 MV

Aakash Institute

9. A planet moving along an elliptical orbit is closest to the sun at a distance r_1 and farthest away at a distance of r_2 . If v_1 and v_2 are the linear velocities at these points respectively. Then the ratio $\frac{v_1}{v_2}$ is

(1)
$$\frac{r_1}{r_2}$$

$$(2) \quad \left(\frac{r_1}{r_2}\right)^2$$

(3)
$$\frac{r_2}{r_1}$$

(4)
$$\left(\frac{r_2}{r_1}\right)^2$$

Sol. Answer (3)

 $v_1 r_1 = v_2 r_2$ (: angular momentum is constant)

10. A radioactive nucleus of mass M emits a photon of frequency v and the nucleus recoils. The recoil energy will be

(2)
$$Mc^2 - hv$$

$$(3) \quad \frac{h^2 v^2}{2Mc^2}$$

Sol. Answer (3)

$$E = \frac{(\text{momentum})^2}{2M} = \frac{\left(\frac{hv}{c}\right)^2}{2M}$$

11. The potential energy of a system increases if work is done

(1) Upon the system by a conservative force

(2) Upon the system by a non-conservative force

(3) By the system against a conservative force

(4) By the system against a non-conservative force

Sol. Answer (3)

By definition

12. A body is moving with velocity 30 m/s towards east. After 10 seconds its velocity becomes 40 m/s towards north. The average acceleration of the body is

$$(1)$$
 5 m/s²

(3)
$$7 \text{ m/s}^2$$

(4)
$$\sqrt{7}$$
 m/s²

Sol. Answer (1)

$$a = \frac{\left|\vec{V}_{f} - \vec{V}_{i}\right|}{t} = \frac{\sqrt{30^{2} + 40^{2}}}{10} = 5 \text{ m/s}^{2}$$

13. A missile is fired for maximum range with an initial velocity of 20 m/s. If g = 10 m/s², the range of the missile is

(1) 20 m

(2) 40 m


(3) 50 m

(4) 60 m

Sol. Answer (2)

$$R_{\text{max}} = \frac{u^2}{g} = \frac{(20)^2}{10} = 40 \text{ m}$$

14. Force *F* on a particle moving in a straight line varies with distance *d* as shown in the figure. The work done on the particle during its displacement of 12 m is



(1) 13 J

(2) 18 J

(3) 21 J

(4) 26 J

Sol. Answer (1)

Work = Area under
$$(F - d)$$
 graph
= $8 + 5 = 13$ J

- 15. A charge Q is enclosed by a Gaussian spherical surface of radius R. If the radius is doubled, then the outward electric flux will
 - (1) Be doubled

(2) Increase four times


(3) Be reduced to half

(4) Remain the same

Sol. Answer (4)

$$\phi_{E} = \frac{Q_{\text{enclosed}}}{\varepsilon_{0}}, \ Q_{\text{enclosed}} \text{ remains unchanged}$$

16. Four electric charges +q, +q, -q and -q are placed at the corners of a square of side 2L (see figure). The electric potential at point A, midway between the two charges +q and +q, is



(1) Zero

(2) $\frac{1}{4\pi\epsilon_0} \frac{2q}{L} (1+\sqrt{5})$

 $(3) \quad \frac{1}{4\pi\varepsilon_0} \frac{2q}{L} \left(1 + \frac{1}{\sqrt{5}} \right)$

 $(4) \quad \frac{1}{4\pi\varepsilon_0} \frac{2q}{L} \left(1 - \frac{1}{\sqrt{5}} \right)$

Sol. Answer (4)

$$V = \frac{2Q}{4\pi\epsilon_0 L} - \frac{2Q}{4\pi\epsilon_0 L\sqrt{5}} = \frac{2Q}{4\pi\epsilon_0 L} \left[1 - \frac{1}{\sqrt{5}} \right] \text{volt}$$

17. A parallel plate condenser has a uniform electric field E (V/m) in the space between the plates. If the distance between the plates is d (m) and area of each plate is A (m²) the energy (joules) stored in the condenser is

(1)
$$\frac{1}{2}\varepsilon_0 E^2 A d$$

(2)
$$E^2Ad/\varepsilon_0$$


(3)
$$\frac{1}{2}\varepsilon_0 E^2$$

(4)
$$\varepsilon_0 EAd$$

Sol. Answer (1)

Direct formula : (energy density = $\frac{1}{2} \varepsilon_0 E^2$) × volume

18. If power dissipated in the 9 Ω resistor in the circuit shown is 36 Watt, the potential difference across the 2 Ω resistor is



(1) 2 Volt

(2) 4 Volt

(3) 8 Volt

(4) 10 Volt

Sol. Answer (4)

Current in 9Ω is 2A, so that in 6Ω is 3A. Total current is 2 + 3 = 5A. Potential drop = $5 \times 2 = 10V$

19. A current of 2 A flows through a 2 Ω resistor when connected across a battery. The same battery supplies a current of 0.5 A when connected across a 9 Ω resistor. The internal resistance of the battery is

(1) 1Ω

(2) 0.5Ω

(3) $1/3 \Omega$

(4) $1/4 \Omega$

Sol. Answer (3)

$$2 = \frac{\varepsilon}{2 + r}$$

$$0.5 = \frac{\varepsilon}{9 + r}$$

or,
$$\frac{2}{0.5} = \frac{9+r}{2+r}$$
 :: $r = \frac{1}{3}\Omega$

20. The rate of increase of thermo e.m.f. with temperature at the neutral temperature of a thermocouple


- (1) Is negative
- (2) Is positive
- (3) Is zero

(4) Depends upon the choice of the two materials of the thermocouple

Sol. Answer (3)

Slope is zero at neutral temperature.

21. A current carrying closed loop in the form of a right angle isosceles triangle ABC is placed in a uniform magnetic field acting along AB. If the magnetic force on the arm BC is \vec{F} , the force on the arm AC is



(1) $\sqrt{2}\vec{F}$

(2) $-\sqrt{2}\vec{F}$

(3) $-\vec{F}$

(4) **F**

Sol. Answer (3)

 $\vec{F} = I\vec{L} \times \vec{B}$

- 22. The power obtained in a reactor using U²³⁵ disintegration is 1000 kW. The mass decay of U²³⁵ per hour is
 - (1) 1 microgram

(2) 10 microgram

(3) 20 microgram

(4) 40 microgram

Sol. Answer (4)

 $E = mc^2$

 $1000 \times 10^3 \times 3600 = m(3 \times 10^8)^2$

- 23. There are four light-weight-rod samples, A, B, C, D separately suspended by threads. A bar magnet is slowly brought near each sample and the following observations are noted
 - (i) A is feebly repelled
 - (ii) B is feebly attracted
 - (iii) C is strongly attracted
 - (iv) D remains unaffected

Which one of the following is true?

- (1) A is of a non-magnetic material
- (2) B is of a paramagnetic material
- (3) C is of a diamagnetic material
- (4) D is of a ferromagnetic material

Sol. Answer (2)

Diamagnetic will be feebly repelled. Paramagnetic will be feebly attracted. Ferromagnetic will be strongly attracted

- 24. The electric and the magnetic field, associated with an e.m. wave, propagating along the +z-axis, can be represented by
 - $(1) \quad \left[\vec{E} = E_0 \hat{j}, \vec{B} = B_0 \hat{k} \right]$

(2) $\left[\vec{E} = E_0 \hat{i}, \vec{B} = B_0 \hat{j}\right]$

(3) $\left[\vec{E} = E_0 \hat{k}, \vec{B} = B_0 \hat{i}\right]$

(4) $\left[\vec{E} = E_0 \hat{j}, \vec{B} = B_0 \hat{i}\right]$

Sol. Answer (2)

 $\vec{E} \times \vec{B}$ points in the direction of wave propagation.

Aakash Institute

25. A uniform electric field and a uniform magnetic field are acting along the same direction in certain region. If an electron is projected in the region such that its velocity is pointed along direction of fields, then the electron

(1) Will turn towards left of direction of motion

(2) Will turn towards right of direction of motion


(3) Speed will decrease


(4) Speed will increase


Sol. Answer (3)


B field will not apply any force. E field will apply a force opposite to velocity of the electron hence, speed will decrease.


26. The current *i* in a coil varies with time as shown in the figure. The variation of induced emf with time would be











Sol. Answer (2)

$$\varepsilon \propto -\frac{di}{dt}$$

27. In an ac circuit an alternating voltage $e = 200\sqrt{2} \sin 100t$ volts is connected to capacitor of capacity 1 μF. The r.m.s. value of the current in the circuit is

(4)

(1) 20 mA

(2) 10 mA

(3) 100 mA

(4) 200 mA

Sol. Answer (1)

$$I_{RMS} = \frac{\varepsilon_0 / \sqrt{2}}{1/\omega C} = 200 \times 100 \times 10^{-6} \text{ A}$$

- 28. An ac voltage is applied to a resistance R and inductor L in series. If R and the inductive reactance are both equal to 3 Ω , the phase difference between the applied voltage and the current in the circuit is
 - (1) Zero

(2) $\pi/6$

(3) $\pi/4$

(4) $\pi/2$

Sol. Answer (3)

$$\tan\theta = \frac{X_L}{R} = 1$$
 : $\phi = 45^{\circ}$ or $\pi/4$

29. When 1 kg of ice at 0°C melts to water at 0° C, the resulting change in its entropy, taking latent heat of ice to be 80 cal/°C is

(1) 293 cal/K

(2) 273 cal/K

(3) $8 \times 10^4 \text{ cal/K}$

(4) 80 cal/K

Sol. Answer (1)

$$\Delta S = \frac{\Delta Q}{T} = \frac{80 \times 1000}{273} \approx 293 \text{ cal/K}$$

- 30. During an isothermal expansion, a confined ideal gas does –150 J of work against its surrounding. This implies that
 - (1) 150 J of heat has been added to the gas
 - (2) 150 J of heat has been removed from the gas
 - (3) 300 J of heat has been added to the gas
 - (4) No heat is transferred because the process is isothermal


Sol. Answer (2)


 $\Delta Q = \Delta U + \Delta W$


 $\Delta Q = O - 150 J$

So, heat has been given by the system

31. A particle of mass *m* is released from rest and follows a parabolic path as shown. Assuming that the displacement of the mass from the origin is small, which graph correctly depicts the position of the particle as a function of time?







(4)

Sol. Answer (2)

Motion given here is SHM starting from rest.

Aakash Institute

Two waves are represented by the equations $y_1 = a\sin(\omega t + kx + 0.57)m$ and $y_2 = a\cos(\omega t + kx)m$, where x is in meter and t in s. The phase difference between them is

(1) 0.57 radian

1.0 radian

(3) 1.25 radian

1.57 radian

Sol. Answer (2)

$$y_2 = a \sin (wt + kx + \frac{\pi}{2})$$

$$y_1 = a \sin (wt + kx + 0.57)$$

Phase difference = $\frac{\pi}{2}$ - 0.57 = 1 radian

33. Out of the following functions representing motion of a particle which represents SHM?

(A) $y = \sin \omega t - \cos \omega t$

(B) $y = \sin^3 \omega t$

(C) $y = 5\cos\left(\frac{3\pi}{4} - 3\omega t\right)$

(D) $y = 1 + \omega t + \omega^2 t^2$

(1) Only (A) and (B)

(2) Only (A)

(3) Only (D) does not represent SHM

(4) Only (A) and (C)

Sol. Answer (4)

For SHM,
$$\frac{d^2y}{dt^2} \propto -y$$

34. Sound waves travel at 350 m/s through a warm air and at 3500 m/s through brass. The wavelength of a 700 Hz acoustic wave as it enters brass from warm air

(1) Decreases by a factor 20

(2) Decreases by a factor 10

(3) Increases by a factor 20

(4) Increases by a factor 10

Sol. Answer (4)

... frequency is same in both the medium

∴ $\lambda \propto$ speed

35. The decreasing order of wavelength of infrared, microwave, ultraviolet and gamma rays is

(1) Infrared, microwave, ultraviolet, gamma rays

(2) Microwave, infrared, ultraviolet, gamma rays

(3) Gamma rays, ultraviolet, infrared, microwaves (4) Microwaves, gamma rays, infrared, ultraviolet

Sol. Answer (2)

Fact based question

The wavelength of the first line of Lyman series for hydrogen atom is equal to that of the second line of Balmer series for a hydrogen like ion. The atomic number Z of hydrogen like ion is

(1) 2

(2)3

(3) 4

(4)

Sol. Answer (1)

$$\left(1-\frac{1}{4}\right)=z^2\left[\frac{1}{4}-\frac{1}{16}\right]$$

 \therefore z = 2

- 37. Which of the following is not due to total internal reflection?
 - (1) Brilliance of diamond
 - (2) Working of optical fibre
 - (3) Difference between apparent and real depth of a pond
 - (4) Mirage on hot summer days

Sol. Answer (3)

Real & apparent depth are explained on the basis of refraction only. TIR not involved here.

- 38. A biconvex lens has a radius of curvature of magnitude 20 cm. Which one of the following options describe best the image formed of an object of height 2 cm placed 30 cm from the lens?
 - (1) Real, inverted, height = 1 cm
- (2) Virtual, upright, height = 1 cm
- (3) Virtual, upright, height = 0.5 cm
- (4) Real, inverted, height = 4 cm

Sol. Answer (4)

In general we have assumed $\mu = 1.5$ so, f = 20cm and calculate v = 60 cm

so, magnification is $\frac{v}{u} = -2$

$$\frac{\left|h_{i}\right|}{\left|h_{o}\right|}=2$$

$$|\mathbf{h}_{\mathsf{i}}| = 2 \times |\mathbf{h}_{\mathsf{0}}|$$

- 39. In photoelectric emission process from a metal of work function 1.8 eV, the kinetic energy of most energetic electrons is 0.5 eV. The corresponding stopping potential is
 - (1) 2.3 V

(2) 1.8 V

(3) 1.3 V

(4) 0.5 V

Sol. Answer (4)

$$eV = KE_{max}$$

- 40. Electrons used in an electron microscope are accelerated by a voltage of 25 kV. If the voltage is increased to 100 km then the de-Broglie wavelength associated with the electrons would
 - (1) Increase by 4 times

(2) Increase by 2 times

(3) Decrease by 2 times

(4) Decrease by 4 times

Sol. Answer (3)

$$\lambda \propto \frac{1}{\sqrt{\text{volt}}}$$

- 41. Light of two different frequencies whose photons have energies 1 eV and 2.5 eV respectively illuminate a metallic surface whose work function is 0.5 eV successively. Ratio of maximum speeds of emitted electrons will be
 - (1) 1:5

(2) 1:4

(3) 1:2

(4) 1:1

Sol. Answer (3)

$$\frac{v_1}{v_2} = \sqrt{\frac{1 - 0.5}{2.5 - 0.5}} = \frac{1}{2}$$

Aakash Institute

- 42. In the Davisson and Germer experiment, the velocity of electrons emitted from the electron gun can be increased by
 - (1) Decreasing the potential difference between the anode and filament
 - (2) Increasing the potential difference between the anode and filament
 - (3) Increasing the filament current
 - (4) Decreasing the filament current

Sol. Answer (2)

Fact

- 43. The half life of a radioactive isotope X is 50 years. It decays to another element Y which is stable. The two elements X and Y were found to be in the ratio of 1 : 15 in a sample of a given rock. The age of the rock was estimated to be
 - (1) 100 years

(2) 150 years

(3) 200 years

(4) 250 years

Sol. Answer (3)

After t second fractional amount of X left is $\frac{1}{16}$ or $\left(\frac{1}{2}\right)^{1}$

$$\therefore t = 4 \times T_{1/2}$$

$$t = 4 \times 50$$

- 44. Photoelectric emission occurs only when the incident light has more than a certain minimum
 - (1) Frequency

(2) Power

(3) Wavelength

(4) Intensity

Sol. Answer (1)

Concept of threshold frequency

- 45. Fusion reaction takes place at high temperature because
 - (1) Molecules break up at high temperature
 - (2) Nuclei break up at high temperature
 - (3) Atoms get ionised at high temperature
 - (4) Kinetic energy is high enough to overcome the coulomb repulsion between nuclei

Sol. Answer (4)

Fact

- 46. A nucleus $_{n}^{m}$ X emits one α particle and two β -particles. The resulting nucleus is
 - (1) $_{n-2}^{m-4}$ Y

(2) ^{m-6}Z

(3) m-6 Z

(4) m-4

Sol. Answer (4)

 α emission decreases mass number by 4 and atomic number by 2. One β^- emission increases atomic number by one but leaves mass number unchanged.

- 47. A transistor is operated in common emitter configuration at V_C = 2 V such that a change in the base current from 100 μ A to 300 μ A produces a change in the collector current from 10 mA to 20 mA. The current gain is
 - (1) 25

(2) 50

(3) 75

(4) 100

Sol. Answer (2)

Current gain
$$\beta = \frac{(20-10)\times 10^{-3}}{(300-100)\times 10^{-6}} = 50$$

- 48. If a small amount of antimony is added to germanium crystal
 - (1) Its resistance is increased
 - (2) It becomes a p-type semiconductor
 - (3) The antimony becomes an acceptor atom
 - (4) There will be more free electrons than hole in the semiconductor

Sol. Answer (4)

Addition of antimony will make it an N-type semiconductor

- 49. In forward biasing of the p-n junction
 - (1) The positive terminal of the battery in connected to p-side and the depletion region becomes thin
 - (2) The positive terminal of the battery is connected to p-side and the depletion region becomes thick
 - (3) The positive terminal of the battery is connected to n-side and the depletion region becomes thin
 - (4) The positive terminal of the battery is connected to n-side and the depletion region becomes thick


Sol. Answer (1)


Fact

50. Symbolic representation of four logic gates are shown as









Pick out which ones are for AND, NAND and NOT gates, respectively:

(1) (ii), (iv) and (iii)

(2) (ii), (iii) and (iv)

(3) (iii), (ii) and (i)

(4) (iii), (ii) and (iv)

Sol. Answer (1)

Fact

Aakash Institute

- 51. The total number of atomic orbitals in fourth energy level of an atom is
 - (1) 4

(2) 8

(3) 16

(4) 32

Sol. Answer (3)

Number of atomic orbitals in an orbit = $n^2 = 4^2 = 16$

- 52. The electrode potentials for $Cu_{(aq)}^{2+} + e^- \rightarrow Cu_{(aq)}^+$ and $Cu_{(aq)}^+ + e^- \rightarrow Cu_{(s)}$ are +0.15 V and +0.50 V respectively. The value of E_{Cu}^{0} will be
 - (1) 0.150 V

(2) 0.500 V

(3) 0.325 V

(4) 0.650 V

Sol. Answer (3)

$$\Delta G_3 = \Delta G_1 + \Delta G_2$$

$$\Rightarrow$$
 -2 FE° = -1F × 0.15 + (-1F × 0.50)

$$\Rightarrow$$
 -2 FE° = -0.15F - 0.50 F

$$\Rightarrow$$
 -2 FE° = -F (0.15 + 0.50)

$$\therefore$$
 E° = $\frac{0.65}{2}$ = 0.325 volt

- 53. Mole fraction of the solute in a 1.00 molal aqueous solution is
 - (1) 1.7700

(2) 0.1770

(3) 0.0177

(4) 0.0344

Sol. Answer (3)

Mole fraction of solute = $\frac{1}{56.55}$ = 0.0177

- 54. By what factors does the average velocity of a gaseous molecule increase when the temperature (in Kelvin) is doubled?
 - (1) 1.4

(2) 2.0

(3) 2.8

(4) 4.0

Sol. Answer (1)

Average velocity = $\sqrt{\frac{8RT}{\pi M}}$

- 55. A buffer solution is prepared in which the concentration of NH_3 is 0.30 M and the concentration of NH_4^+ is 0.20 M. If the equilibrium constant, K_b for NH_3 equals 1.8 × 10⁻⁵, what is the pH of this solution?
 - (1) 8.73

(2) 9.08

(3) 9.43

(4) 11.72

Sol. Answer (3)

$$p^{OH} = pK_b + log \frac{[Salt]}{[Base]} = 4.74 + log \frac{0.20}{0.30} = 4.74 + (0.301 - 0.477)$$

$$= 4.74 - 0.176 = 4.56$$

$$\therefore$$
 PH = 14 - 4.56 = 9.44

- 56. Two gases A and B having the same volume diffuse through a porous partition in 20 and 10 seconds respectively. The molecular mass of A is 49 u. Molecular mass of B will be
 - (1) 25.00 u

(2) 50.00 u

(3) 12.25 u

(4) 6.50 u

Sol. Answer (3)

$$\frac{r_A}{r_B} = \sqrt{\frac{M_B}{M_A}}$$

$$\Rightarrow \frac{v_A}{t_A} \times \frac{t_B}{v_B} = \sqrt{\frac{M_B}{M_A}}$$

$$\Rightarrow \frac{10}{20} = \sqrt{\frac{M_B}{49}}$$

$$\Rightarrow \frac{1}{4} = \frac{M_B}{49}$$

$$M_{\rm B} = \frac{49}{4} = 12.25$$

- 57. Which of the following is correct option for free expansion of an ideal gas under adiabatic condition?
 - (1) $q = 0, \Delta T < 0, w \neq 0$

(2) $q = 0, \Delta T \neq 0, w = 0$

(3) $q \neq 0$, $\Delta T = 0$, w = 0

(4) $q = 0, \Delta T = 0, w = 0$

Sol. Answer (4)

For an ideal gas, for free expansion

$$q = 0$$
; $\Delta T = 0 & w = 0$

58. For the reaction $N_2(g) + O_2(g) \Longrightarrow 2NO(g)$, the equilibrium constant is K_1 . The equilibrium constant is $K_2(g) + O_2(g) \Longrightarrow 2NO_2(g)$. What is K for the reaction $NO_2(g) \Longrightarrow \frac{1}{2}N_2(g) + O_2(g)$?

(1)
$$\frac{1}{(K_1K_2)}$$

(2)
$$\frac{1}{(2K_1K_2)}$$

(3)
$$\frac{1}{(4K_1K_2)}$$

$$(4) \quad \left\lceil \frac{1}{\mathsf{K}_1 \mathsf{K}_2} \right\rceil^{1/2}$$

Sol. Answer (4)

$$N_2(g) + O_2(g) \rightleftharpoons 2NO(g); K_1$$

$$2NO(g) + O_2(g) \rightleftharpoons 2NO_2(g); K_2$$

$$N_2(g) + 2O_2(g) \rightleftharpoons 2 NO_2(g); K = K_1 \times K_2$$

$$\therefore \text{ For NO}_2(q) \iff \frac{1}{2} N_2(g) + O_2(g); K' = \left[\frac{1}{K_1 \cdot K_2}\right]^{\frac{1}{2}}$$

Aakash Institute

59. If x is amount of adsorbate and m is amount of adsorbent, which of the following relations is not related to adsorption process?

(1)
$$\frac{x}{m} = p \times T$$

(2)
$$\frac{x}{m} = f(p)$$
 at constant T

(3)
$$\frac{x}{m} = f(T)$$
 at constant p

(4)
$$p = f(T)$$
 at constant $\left(\frac{x}{m}\right)$

Sol. Answer (1)

 $x/m = P \times T$ is the incorrect relation

60. If the enthalpy change for the transition of liquid water to steam is 30 kJ mol⁻¹ at 27°C, the entropy change for the process would be

(4)
$$0.1 \text{ J mol}^{-1} \text{ K}^{-1}$$

Sol. Answer (1)

$$\Delta S_{\text{vap}} = \frac{\Delta H_{\text{vap}}}{T} = \frac{30 \text{KJmol}^{-1}}{300 \text{K}} = 100 \text{ J mol}^{-1} \text{ K}^{-1}$$

61. The van't Hoff factor i for a compound which undergoes dissociation in one solvent and association in other solvent is respectively

(1) Greater than one and greater than one

(2) Less than one and greater than one

(3) Less than one and less than one

(4) Greater than one and less than one

Sol. Answer (4)

Fact

62. Standard electrode potential for Sn^{4+}/Sn^{2+} couple is +0.15 V and that for the Cr^{3+}/Cr couple is -0.74 V. These two couples in their standard state are connected to make a cell. The cell potential will be

Sol. Answer (3)

$$E_{cell}^{o} = E_{cathode(RP)}^{o} - E_{anode(RP)}^{o}$$

= 0.15 - (-0.74)
= +0.89 V

63. A gaseous mixture was prepared by taking equal mole of CO and N₂. If the total pressure of the mixture was found 1 atmosphere, the partial pressure of the nitrogen (N₂) in the mixture is

(1) 1 atm

(2) 0.5 atm

(3) 0.8 atm

(4) 0.9 atm

Sol. Answer (2)

Fact

64. If the E_{cell}^{o} for a given reaction has a negative value, then which of the following gives the correct relationships for the values of ΔG° and K_{eq} ?

(1)
$$\Delta G^{\circ} > 0$$
; $K_{eq} < 1$

(2)
$$\Delta G^{\circ} > 0$$
; $K_{eq} > 1$

(3)
$$\Delta G^{\circ} < 0; K_{eq} > 1$$

(4)
$$\Delta G^{\circ} < 0$$
; $K_{eq} < 1$

Sol. Answer (1)

$$\Delta G^{\circ} = -nFE^{\circ}$$

&
$$\Delta G^{\circ} = -RT \log_e K_{eq}$$

65. The freezing point depression constant for water is -1.86°Cm⁻¹. If 5.00 g Na₂SO₄ is dissolved in 45.0 g H₂O, the freezing point is changed by -3.82°C. Calculate the van't Hoff factor for Na₂SO₄

Sol. Answer (3)

Using,
$$\Delta T_f = i \times K_f \times m$$

$$i = \frac{\Delta T_f \times W_A}{K_f \times n_B \times 1000} = \frac{3.82 \times 45}{1.86 \times \left(\frac{5}{142}\right) \times 1000} = 2.63$$

66. The energies E_1 and E_2 of two radiations are 25 eV and 50 eV respectively. The relation between their wavelengths i.e. λ_1 and λ_2 will be

$$(1) \quad \lambda_1 = \frac{1}{2}\lambda_2$$

(2)
$$\lambda_1 = \lambda_2$$

(3)
$$\lambda_1 = 2\lambda_2$$

$$(4) \quad \lambda_1 = 4\lambda_2$$

Sol. Answer (3)

$$\lambda_1 = 2\lambda_2$$

67. Standard electrode potential of three metals X, Y and Z are -1.2 V, +0.5 V and -3.0 V respectively. The reducing power of these metals will be

(1)
$$X > Y > Z$$

(2)
$$Y > Z > X$$

(3)
$$Y > X > Z$$

$$(4) \quad Z > X > Y$$

Sol. Answer (4)

Z > X > Y; higher the reduction potential lesser the reducing power

- 68. Which one of the following statements for the order of a reaction is incorrect?
 - (1) Order of reaction is always whole number
 - (2) Order can be determined only experimentally
 - (3) Order is not influenced by stoichiometric coefficient of the reactants
 - (4) Order of reaction is sum of power to the concentration terms of reactants to express the rate of reaction
- Sol. Answer (1)

Fact

Aakash Institute

69. Enthalpy change for the reaction, $4H_{(q)} \rightarrow 2H_{2(q)}$ is -869.6 kJ. The dissociation energy of H - H bond is

(1) +217.4 kJ

(2) -434.8 kJ

(3) -869.6 kJ

(4) +434.8 kJ

Sol. Answer (4)

The dissociation energy of H–H bond is = $\frac{869.6}{2}$ = 434.8KJ

70. If n = 6, the correct sequence of filling of electrons will be

(1) $ns \rightarrow np(n-1)d \rightarrow (n-2)f$

- (2) $ns \rightarrow n(n-2)f \rightarrow (n-1)d \rightarrow np$
- (3) $ns \rightarrow (n-1)d \rightarrow (n-2)f \rightarrow np$
- (4) $ns \rightarrow (n-2)f \rightarrow np \rightarrow (n-1)d$

Sol. Answer (2)

Fact

71. Which of the following compounds has the lowest melting point?

(1) CaF₂

(2) CaCl₂

(3) CaBr₂

(4) Cal₂

Sol. Answer (4)

Cal₂ has lowest melting point

72. Which of the following pairs of metals is purified by van Arkel method?

(1) Ni and Fe

(2) Ga and In

(3) Zr and Ti

(4) Ag and Au

Sol. Answer (3)

Zr and Ti are purified by van Arkel method

73. The correct order of increasing bond length of C - H, C - O, C - C and C = C is

- (1) C-H < C-O < C-C < C=C
- (2) C H < C = C < C O < C C
- (3) C C < C = C < C O < C H
- (4) C O < C H < C C < C = C

Sol. Answer (2)

Fact

74. Acidified K₂Cr₂O₇ solution turns green when Na₂SO₃ is added to it. This is due to the formation of

(1) CrSO₄

(2) $\operatorname{Cr}_{2}(\operatorname{SO}_{4})_{3}$

(3) CrO_4^{2-}

(4) $Cr_2(SO_3)_3$

Sol. Answer (2)

$$K_2Cr_2O_7 + 3 Na_2SO_3 + 4H_2SO_4 \rightarrow 3 Na_2SO_4 + K_2SO_4 + Cr_2 (SO_4)_3$$

75. For the four successive transition elements (Cr, Mn, Fe and Co), the stability of + 2 oxidation state will be there in which of the following order?

(1) Cr > Mn > Co > Fe

(2) Mn > Fe > Cr > Co

(3) Fe > Mn > Co > Cr

(4) Co > Mn > Fe > Cr

(At. nos. Cr = 24, Mn = 25, Fe = 26, Co = 27)

Sol. Answer (2)

On the basis of electrode potentials, the correct order is Mn > Fe > Cr > Co

Aakash Institute

CBSE Prelim. 2011 (Hints & Solutions) - Code A

76. Which of the two ions from the list given below that have the geometry that is explained by the same hybridization of orbitals, NO_2^- , NO_3^- , NH_2^- , NH_4^+ , SCN^- ?

(1) NO_2^- and NH_2^-

(2) NO_2^- and NO_3^-

(3) NH₄⁺ and NO₃⁻

(4) SCN^- and NH_2^-

Sol. Answer (2)

 $\mathrm{NO_2^{(-)}}$ and $\mathrm{NO_3^{(-)}}$ both have same hybridisation

77. Which of the following elements is present as the impurity to the maximum extent in the pig iron?

(1) Phosphorus

(2) Manganese

(3) Carbon

(4) Silicon

Sol. Answer (3)

Fact

78. Which of the following is least likely to behave is Lewis base?

(1) OH

(2) H₂O

(3) NH₃

(4) BF₃

Sol. Answer (4)

BF₃ is an electron deficient species

79. Which one of the following is present as an active ingredient in bleaching powder for bleaching action?

(1) CaCl₂

(2) CaOCl₂

(3) Ca $(OCI)_2$

(4) CaO₂Cl

Sol. Answer (3)

Fact

80. The complex, [Pt (Py) (INH₃) Br Cl] will have how many geometrical isomers?


(1) 2

(2) 3

(3) 4

(4) 0

Sol. Answer (2)



81. Name the type of the structure of silicate in which one oxygen atom of $[SiO_4]^{4-}$ is shared?


(1) Three dimensional

(2) Linear chain silicate

(3) Sheet silicate

(4) Pyrosilicate

Sol. Answer (4)



Aakash Institute

82. The complexes [Co (NH₃)₆] [Cr (CN)₆] and [Cr(NH₃)₆] [Co(CN)₆] are the examples of which type of isomerism?

(1) Geometrical isomerism

(2) Linkage isomerism

(3) Ionization isomerism

(4) Coordination isomerism

Sol. Answer (4)

Co-ordination isomerism

83. The d-electron configurations of Cr²⁺, Mn²⁺, Fe²⁺ and Co²⁺ are d⁴, d⁵, d⁶ and d⁷ respectively. Which one of the following will exhibit minimum paramagnetic behaviour?

(1) $[Cr(H_2O)_6]^{2+}$

(2) $[Mn(H_2O)_6]^{2+}$

(3) $[Fe(H_2O)_6]^{2+}$

(4) $[Co(H_2O)_6]^{2+}$

(At. nos. Cr = 24, Mn = 25, Fe = 26, Co = 27)

Sol. Answer (4)

$$\text{Co}^{2+}
ightarrow 3\text{d}^7 \ 4\text{s}^\circ$$

1111111

; having minimum no. of unpaired electrons

84. Of the following complex ions, which is diamagnetic in nature?

(1) $[CoF_6]^{3-}$


(2) $[NiCl_{4}]^{2-}$

(3) $[Ni(CN)_4]^{2-}$

(4) [CuCl₄]²⁻

Sol. Answer (3)

Ni has dsp² hybridisation where CN⁽⁻⁾ is a strong ligand



85. Which of the following has the minimum bond length?

(1) O_2

(2) O^{2}

(3) O_2^-

(4) O_2^{2-}

Sol. Answer (2)

$$O_2^{(+)}$$
 has B.O 2.5 and B.O $\propto \frac{1}{\text{Bond length}}$

86. The value of ΔH for the reaction

 $X_{2(g)}$ + $4Y_{2(g)}$ \Longrightarrow $2XY_{4(g)}$ is less than zero. Formation of $XY_{4(g)}$ will be favoured at

- (1) High pressure and low temperature
- (2) High temperature and high pressure
- (3) Low pressure and low temperature
- (4) High temperature and low pressure

Sol. Answer (1)

$$\Delta n_g = -ve$$
 and $\Delta H = -ve$

- 87. Of the following which one is classified as polyester polymer?
 - (1) Nylon-66

(2) Terylene

(3) Backelite

(4) Melamine

Sol. Answer (2)

Terylene is a polyester polymer because it is formed by the monomer units terephthalic acid and ethylene glycol

88. What is the product obtained in the following reaction

Sol. Answer (2)

89. In a set of reactions m-bromobenzoic acid gave a product D. Identify the product D.

$$\begin{array}{c}
COOH \\
\hline
D
\end{array}$$

$$\begin{array}{c}
SOCI_2 \\
Br_2
\end{array}$$

$$\begin{array}{c}
NH_3 \\
Br_2
\end{array}$$

$$\begin{array}{c}
NaOH \\
Br_2
\end{array}$$

$$\begin{array}{c}
D
\end{array}$$

Sol. Answer (4)

COOH
$$COCI$$

$$CONH_{2}$$

$$NH_{2}$$

$$NH_{3}$$

$$ROH$$

$$Br$$

$$ROH$$

$$Br$$

$$ROH$$

$$Br$$

$$ROH$$

$$RO$$

90. In Dumas' method of estimation of nitrogen 0.35 g of an organic compound gave 55 mL of nitrogen collected at 300 K temperature and 715 mm pressure. The percentage composition of nitrogen in the compound would be

(Aqueous tension at 300 K = 15 mm)

(1) 14.45

(2) 15.45

(3) 16.45

(4) 17.45


Sol. Answer (3)

$$\frac{P_1V_1}{T_1} = \frac{P_2V_2}{T_2}, \text{on calculation}$$

$$V_2 = 46.09 \text{ ml}$$

% of N =
$$\frac{28 \times 46.09 \times 100}{22400 \times 0.35} = 16.45$$

91. Which one of the following is most reactive towards electrophilic reagent ?



Sol. Answer (3)

+R effect of OH > OCH₃

- 92. Which one is a nucleophilic substitution reaction among the following?
 - (1) $CH_3CHO + HCN \longrightarrow CH_3CH(OH)CN$

(2)
$$CH_3 - CH = CH_2 + H_2O \xrightarrow{H^+} CH_3 - CH - CH_3$$

OH

(3) RCHO + R'MgX
$$\rightarrow$$
 R - CH - R' | OH

$$(4) \quad CH_3 - CH_2 - CH - CH_2Br + NH_3 \longrightarrow CH_3 - CH_2 - CH - CH_2NH_2$$

Sol. Answer (4)

Fact

- 93. Considering the state of hybridization of carbon atoms, find out the molecule among the following which is linear?
 - (1) $CH_3 CH_2 CH_2 CH_3$

(2) $CH_3 - CH = CH - CH_3$

(3) $CH_3 - C \equiv C - CH_3$

(4) $CH_2 = CH - CH_2 - C \equiv CH$

Sol. Answer (3)

94. In the following reactions,

(a)
$$CH_3 - CH - CH - CH_3 \xrightarrow{H^+/Heat} A + B \xrightarrow{Major (product)} CH$$

(b)
$$A \xrightarrow{\text{HBr, dark}} C \xrightarrow{\text{Major}} + D \xrightarrow{\text{Minor Product}} + \left(\begin{array}{c} D \\ \text{Minor Product} \end{array} \right)$$

The major products (A) and (C) are respectively

(3)
$$CH_3 - C = CH - CH_3$$
 and $CH_3 - C - CH_2 - CH_3$

$$\begin{array}{c|c} CH_3 & CH_3 \\ | & | \\ (4) & CH_3 - C = CH - CH_3 \text{ and } CH_3 - CH - CH - CH_3 \\ | & | \\ Br \end{array}$$

Sol. Answer (3)

- The Lassaigne's extract is boiled with conc. HNO₃ while testing for halogens. By doing so it.

 - (1) Increase the concentration of NO_3^- ions (2) Decomposes Na_2S and NaCN, if formed
 - (3) Helps in the precipitation of AgCI
- (4) Increases the solubility product of AgCl

Sol. Answer (2)

Fact

96. The correct IUPAC name of the compound


(1) 3-(1-ethyl propyl) hex-1-ene

(2) 4-Ethyl-3-propyl hex-1-ene

(3) 3-Ethyl-4-ethenyl heptane

(4) 3-Ethyl-4-propyl hex-5-ene

Sol. Answer (2)



4-Ethyl-3-propylhex-1-ene

- 97. Clemmensen reduction of a ketone is carried out in the presence of which of the following?
 - (1) H₂ and Pt as catalyst

(2) Glycol with KOH

(3) Zn-Hg with HCl

(4) LiAIH

Sol. Answer (3)

 $C = O \xrightarrow{Z_1 \cdot H_2} CH_2$; clemmensen reduction

- 98. Which one of the following is employed as Antithistamine?
 - (1) Omeprazole

(2) Chloramphenicol

(3) Diphenyl hydramine

(4) Norothindrone

Sol. Answer (3)

Diphenyl hydramine is employed as Antihistamine

- 99. Which one of the following statements is not true regarding (+) Lactose?
 - (1) (+) Lactose, C₁₂H₂₂O₁₁ contains 8-OH groups
 - (2) On hydrolysis (+) Lactose gives equal amount of D(+) glucose and D(+) galactose
 - (3) (+) Lactose is a β -glycoside formed by the union of a molecule of D(+) glucose and a molecule of D(+) galactose
 - (4) (+) Lactose is reducing sugar and does not exhibit mutarotation

Sol. Answer (4)

- (+) lactose is a reducing sugar and it exhibit mutarotation
- 100. Which one of the following statement is no true?
 - (1) Oxides of sulphur, nitrogen and carbon are the most widespread air pollutant
 - (2) pH of drinking water should be between 5.5 9.5
 - (3) Concentration of DO below 6 ppm is good for the growth of fish
 - (4) Clean water would have a BOD value of less than 5 ppm

Sol. Answer (3)

Fact

Aaka:	sh Institute		CBSE Prelim. 2011 (Hints & Solutions) - Code A
101.	The "Eyes" of the potato tuber are		
	(1) Axillary buds	(2)	Root buds
	(3) Flower buds	(4)	Shoot buds
Sol.	Answer (1) Axillary buds developing at nodes/notch/eyes.		
102.	Organisms called Methanogens are most abundant	in a	
	(1) Hot spring	(2)	Sulphur rock
	(3) Cattle yard	(4)	Polluted stream
Sol.	Answer (3) Methanogens are archaebacteria abundant in cattle	yard	l, and paddy fields.
103.	Which one of the following have the highest number	r of s	pecies in nature?
	(1) Angiosperms	(2)	Fungi
	(3) Insects	(4)	Birds
Sol.	Answer (3) The largest phylum in animal kingdom is arthropoda	a, an	d the largest class is insecta with 7,50,000 species.
104.	Archegoniophore is present in		
	(1) Funaria	(2)	Marchantia
	(3) Chara	(4)	Adiantum
Sol.	Answer (2) Stalk bearing archegonial cluster at tip in Marchanti	a tha	allus.
105.	Compared with the gametophytes of the bryophytes	the	gametophytes of vascular plants tend to be
	(1) Smaller and to have smaller sex organs	(2)	Smaller but to have larger sex organs
	(3) Larger but to have smaller sex organs	(4)	Larger and to haver larger sex organs
Sol.	Answer (1) Gametophytes are reduced and few celled in all vas	scula	r plants.
106.	The gametophyte is not an independent, free-living g	genei	ration in
	(1) Pinus	(2)	Polytrichum
	(3) Adiantum	(4)	Marchantia
Sol.	Answer (1) In gymnosperms and angiosperms gametophytes an	e de	pendent on sporophyte.
107.	Important site for formation of glycoproteins and gly	colip	ids is
	(1) Lysosome	(2)	Vacuole
	(3) Golgi apparatus	(4)	Plastid
Sol.	Answer (3) Golgi complex performs glycosyl transferase activity	/ for	addition of glycans on lipids and proteins.

CBSE	Prelim. 2011 (Hints & Solutions) - Code A		Aakash Institute
	Peptide synthesis inside a cell takes place in		
	(1) Ribosomes	(2)	Chloroplast
	(3) Mitochondria	(4)	Chromoplast
Sol.	Answer (1) Ribosomes are site of peptide bond formation.		
109.	In eubacteria, a cellular component that resembles	euka	aryotic cell is
	(1) Cell wall	(2)	Plasma membrane
	(3) Nucleus	(4)	Ribosomes
Sol.	Answer (2) Lipoprotein cell membrane is found in both but ribos	some	es are of different kinds.
110.	Mutations can be induced with		
	(1) Gamma radiations	(2)	Infra Red radiations
	(3) I A A	(4)	Ethylene
Sol.	Answer (1) Mutation can be induced with high energy radiation structure of DNA.	ns lik	ke UV rays, gamma rays, which cause change in the
111.	A collection of plants and seeds having diverse alle	eles c	of all the genes of a crop is called
	(1) Genome		
	(2) Herbarium		
	(3) Germplasm		
	(4) Gene library		
Sol.	Answer (3) Germplasm can be selected as seed or plantlets for	or the	eir superior traits.
112.	Which one of the following also acts as a catalyst	in a l	bacterial cell?
	(1) 23 sr RNA	(2)	5 sr RNA
	(3) sn RNA	(4)	hn RNA
Sol.	Answer (1) 23 S rRNA is catalytic RNA.		

- 113. Which one of the following statements is **correct**?
 - (1) Flower of tulip is a modified shoot
 - (2) In tomato, fruit is a capsule
 - (3) Seeds of orchids have oil-rich endosperms
 - (4) Placentation in primose is basal


Sol. Answer (1)

Tomato — Berry, Orchid seed — no endosperm formation, Primrose — Free central placentation

Aakash Institute

CBSE Prelim. 2011 (Hints & Solutions) - Code A

114. The correct floral formula of chilli is



(2)
$$\bigoplus \Phi K_{(5)}C_5A_5G_{(2)}$$

(4) $\bigoplus \Phi K_{(5)}C_{(5)}A_{(5)}G_2$

(3)
$$\bigoplus \bigoplus_{i=1}^{4} K_{(5)} C_{(5)} A_{5} G_{(2)}$$

(4)
$$\oplus \Phi^{\bullet}_{K_{(5)}}C_{(5)}A_{(5)}G_{2}$$

Sol. Answer (3)

Chilli belongs to Solanaceae.

115. Nitrifying bacteria

- (1) Reduce nitrates to free nitrogen
- (2) Oxidize ammonia to nitrates
- (3) Convert free nitrogen to nitrogen compounds
- (4) Convert proteins into ammonia

Sol. Answer (2)

$$NH_3 \xrightarrow{Step (1)} NO_2^- \xrightarrow{Step (2)} NO_3^-$$

Step (1) — Nitrification by Nitrosomonas

Step (2) — Nitratification by *Nitrocystis*

116. The function of leghaemoglobin in the root nodules of legumes is

(1) Expression of nif gene

Inhibition of nitrogenase activity

(3) Oxygen removal

Nodule differentiation

Sol. Answer (3)

LHB is O₂ scavanger.

117. Which one of the following elements in plants is not remobilised?

(1) Sulphur

Phosphorus

(3) Calcium

(4)Potassium

Sol. Answer (3)

Calcium is not remobilised, as it is a structural component in cell.

118. A drupe develops in

(1) Tomato

Mango

(3) Wheat

Pea

Sol. Answer (2)

Tomato — Berry, Wheat — Caryopsis, Pea — Legume

119. Ground tissue includes

- (1) All tissues internal to endodermis
- (2) All tissues external to endodermis
- (3) All tissues except epidermis and vascular bundles
- (4) Epidermis and cortex

Sol. Answer (3)

Ground tissue system includes — cortex, endoderm, pericycle and pith.

120. In land plants the guard cells differ from other epidermal cells in having

(1) Chloroplasts

(2) Cytoskeleton

(3) Mitochondria

Endoplasmic reticulum

Sol. Answer (1)

Guard cells are specialised chlorophyllous epidermal cells.

CBSE	Prelim. 2011 (Hints & Solutions) - Code A			Aakash Institute
121.	The ovary is half inferior in flowers of			
	(1) Guava	(2)	Peach	
	(3) Cucumber	(4)	Cotton	
Sol.	Answer (2) Ovary is half inferior in perigynous flowers.			
122.	The cork cambium, cork and secondary cortex are	colle	ctively called	
	(1) Phellem	(2)	Phelloderm	
	(3) Phellogen	(4)	Periderm	
Sol.	Answer (4) Phellem, phellogen and phelloderm are collectively	called	d periderm.	
123.	Which one of the following is wrongly matched?			
	(1) Cassia – Imbricate aestivation	(2)	Root pressure – Guttation	
	(3) Puccinia – Smut	(4)	Root – Exarch protoxylem	
Sol.	Answer (3) Puccinia — rust fungi.			
124.	Flowers are Zygomorphic in			
	(1) Datura	(2)	Mustard	
	(3) Gulmohur	(4)	Tomato	
Sol.	Answer (3) Datura, mustard and tomato have actinomorphic flor	wers.		
125.	CAM helps the plants in			
	(1) Reproduction	(2)	Conserving water	
	(3) Secondary growth	(4)	Disease resistance	
Sol.	Answer (2) These are succulent plants with water storing cells.			
126.	Of the total incident solar radiation the proportion o	f Paf	R is	
	(1) More than 80%	(2)	About 70%	
	(3) About 60%	(4)	Less than 50%	
Sol.	Answer (4) Plants capture 2-10% of PAR.			
127.	A prokaryotic autotrophic nitrogen fixing symbiont for	ound	in	
	(1) Pisum	(2)	Alnus	
	(3) Cycas	(4)	Cicer	
Sol.	Answer (3) Anabaena cycadae is a BGA found in coralloid root	s of	Cycas.	
128.	Nucellar polyembryony is reported in species of			
	(1) Brassica	(2)	Citrus	
	(3) Gossypium	(4)	Triticum	
Sol.	Answer (2) Nucellus polyembryony is common in Citrus, mange	o and	d Opuntia.	

Aaka	sh Institute		CBSE Prelim. 2011 (Hints & Solutions) - Code A
129.	Filiform apparatus is a characteristic feature of		
	(1) Zygote	(2)	Suspensor
	(3) Egg	(4)	Synergid
Sol.	Answer (4) These are fingure like projections at micropylar end	of s	ynergids.
130.	What would be the number of chromosomes of the tip cells?	aleu	rone cells of a plant with 42 chromosomes in its roots
	(1) 21	(2)	42
	(3) 63	(4)	84
Sol.	Answer (3) Aleurone is triploid and root tip is diploid.		
131.	Wind pollination is common in		
	(1) Orchids	(2)	Legumes
	(3) Lilies	(4)	Grasses
Sol.	Answer (4) Wind pollination is common in grasses and gymnos	sperr	ns.
132.	In which one of the following pollination is autogame	ous?	
	(1) Cleistogamy	(2)	Geitonogamy
	(3) Xenogamy	(4)	Chasmogamy
Sol.	Answer (1) Self pollination is favoured by cleistogamy.		
133.	Mass of living matter at a trophic level in an area a	t any	time is called
	(1) Standing state	(2)	Standing crop
	(3) Detritus	(4)	Humus
Sol.	Answer (2) Standing state represent all non-living matter in an a	area	at a given time.
134.	Which one of the following statements is wrong in o	case	of Bhopal tragedy?
	(1) It took place in the night of December 2/3/1984		
	(2) Methyl Isocyanate gas leakage took place		
	(3) Thousands of human beings died		
	(4) Radioactive fall out engulfed Bhopal		
Sol.	Answer (4) It was not a tragedy related to radioactivity.		
135.	Secondary sewage treatment is mainly a		
	(1) Biological process	(2)	Physical process
	(3) Mechanical process	(4)	Chemical process
Sol.	Answer (1)		
	Secondary sewage treatment involves aerobic and a	anaeı	rodic microdes.

	E Prelim. 2011 (Hints & Solutions) - Code A		Aakash Institute
130	Eutrophication is often seen in (1) Mountains	(2)	Deserts
		(2)	
0.1	(3) Fresh water lakes	(4)	Ocean
Soi	 Answer (3) It is process of enrichment of lakes by phosphates 	s, nitr	ates etc.
137	. Large Woody Vines are more commonly found in		
	(1) Alpine forests	(2)	Temperate forests
	(3) Mangroves	(4)	Tropical rainforests
Sol	. Answer (4)	()	'
	Lianas and epiphytes are more common in tropica	l rain	forest.
138	. Which one of the following expanded forms of the	follow	vings acronyms is correct?
	(1) IUCN = International Union for Conservation of	Natu	re and Natural Resources
	(2) IPCC = International Panel for Climate Change		
	(3) UNEP = United Nations Environmental Policy		
	(4) EPA = Environmental Pollution Agency		
Sol	. Answer (1)		
	IPCC — Intergovernmental Panel for Climate Chan	ge.	
139	. Which one of the following statements is correct for	or sec	condary succession?
	(1) It is similar to primary succession except that	it has	s a relatively fast pace
	(2) It begins on a bare rock		
	(3) It occurs on a deforested site		
	(4) It follows primary succession		
Sol	Answer (3) Secondary biotic succession occurs in abandoned flooded.	farm	lands, burned or cut forests and lands that have been
140	. Which one of the following shows maximum genet	ic div	ersity in India?
	(1) Mango	(2)	Groundnut
	(3) Rice	(4)	Maize
Sol	Answer (3) Rice has more than 50,000 genetically different str	ains,	while mango has 1000 varieties in India.
141	. Which one of the following is not a biofertilizer?		
	(1) Mycorrhiza	(2)	Agrobacterium
	(3) Rhizobium	(4)	Nostoc
Sol	Answer (2)		

Agrobacterium is a gene transfer agent.

Aakas	sn in	stitute		CBSE Prelim. 2011 (Hints & Solutions) - Code A
142.	Wh	ich one of the following acts as a physiological b	oarrie	er to the entry of microorganisms in human body?
	(1)	Skin	(2)	Epithelium of Urogenial tract
	(3)	Tears	(4)	Monocytes
Sol.	Phy	swer (3) visiological barriers to the entry of micro-organism in stomach.	ms in	human body are tears in eyes, saliva in mouth and
143.	Wh	ich one of the following helps in absorption of ph	nosph	norus from soil by plants?
	(1)	Anabaena	(2)	Glomus
	(3)	Rhizobium	(4)	Frankia
Sol.		swer (2) nmus is a endomycorrhiza for phosphorus absorp	otion.	
144.	'Hin of	ngiri' developed by hybridisation and selection f	or dis	sease resistance against rust pathogens is a variety
	(1)	Wheat	(2)	Chilli
	(3)	Maize	(4)	Sugarcane
Sol.		swer (1) s variety is resistant against leaf and stripe rust,	hill b	ount.
145.	Wh	ich of the followings is mainly produced by the a	ctivity	y of anaerobic bacteria on sewage?
	(1)	Marsh gas	(2)	Laughing gas
	(3)	Propane	(4)	Mustard gas
Sol.		swer (1) s by the activity of methanogens.		
146.	Aga	arose extracted from sea weeds finds use in		
	(1)	Gel electrophoresis	(2)	Spectrophotometry
	(3)	Tissue culture	(4)	PCR
Sol.		swer (1) arose extracted from sea weeds finds use in gel	elect	rophoresis.
147.	Max	ximum number of existing transgenic animals is	of	
	(1)	Pig	(2)	Fish
	(3)	Mice	(4)	Cow
Sol.		swer (3) % of the existing transgenic animals are mice.		
148.	Cor	ntinuous addition of sugars in 'fed batch' ferment	ation	is done to
	(1)	Degrade sewage	(2)	Produce methane
	(3)	Obtain antibiotics	(4)	Purify enzymes
Sol.		swer (4) https://doi.orgiv.org	tion is	s done to purify enzymes.

Aakash Institute

- 149. The process of RNA interference has been used in the development of plants resistant to
 - (1) Insects

(2) Nematodes

(3) Fungi

(4) Viruses

Sol. Answer (2)

RNAi *i.e.*, RNA interference is used in the development of plants resistant to nematode like *Meloidegyne incognita*.

- 150. "Jaya" and "Ratna" developed for green revolution in India are the varieties of
 - (1) Bajra

(2) Maize

(3) Rice

(4) Wheat

Sol. Answer (3)

Jaya and Ratna are released throughout the rice growing belts of India.

- 151. Which one of the following organisms is not an example of eukaryotic cells
 - (1) Amoeba proteus

(2) Paramecium caudatum

(3) Escherichia coli

(4) Euglena viridis

Sol. Answer (3)

E. coli is a prokaryotic bacterium.

- 152. Which one of the following animals is correctly matched with its particular named taxonomic category?
 - (1) Housefly Musca, an order

(2) Tiger - Tigris, the species

(3) Cuttlefish - Mollusca, a class

(4) Humans - Primata, the family

Sol. Answer (2)

The zoological name of tiger is Panthera tigris in which Panthera is genus and tigris is species.

153. In which one of the following the genus name, its two characters and its class/phylum are correctly matched?

	Genus name	Two characters Class/ Phylum		Class/ Phylum	
		(a)	cnidoblasts	Coelenterata	
(1)	Aurelia	(b)	Organ level of organization		
		(a)	Body segmented		
(2)	Ascaris	(b)	Males and females distinct	Annelida	
(3)	Salamandra	(a)	A tympanum represents ear	Amphibia	
	((b)	Fertilization is external	Типриющ	
(4)	Pteropus (a)		Skin possesses hair	Mammalia	
			Oviparous		

Sol. Answer (3)

Salamandra is a tailed amphibian, has tymphanum which represents ear.

- 154. Which one of the following groups of animals is correctly matched with its one characteristic feature without even a single exception?
 - (1) Mammalia: give birth to young ones
 - (2) Reptilia: possess 3-chambered heart with one incompletely divided ventricle
 - (3) Chordata: possess a mouth provided with an upper and a lower jaw
 - (4) Chondrichthyes: possess cartilaginous endoskeleton

Sol. Answer (4)


All chondrichthyes possess cartilaginous endoskeleton without exception.

- 155. What will you look for to identify the sex of the following?
 - (1) Male shark Claspers borne on pelvic fins
 - (2) Female Ascaris Sharply curved posterior end
 - (3) Male frog A copulatory pad on the first digit of the hind limb
 - (4) Female cockroach Anal cerci

Sol. Answer (1)

In class chondrichthyes males possess claspers on the pelvic fins.

156. The curve given below show enzymatic activity with relation to three conditions (pH, temperature and substrate concentration)



What do the two axises(x and y) represent?

	x-axis	y-axis
(1)	Enzymatic activity	Temeperature
(2)	Enzymatic activity	pН
(3)	Temperature	Enzyme
		Activity
(4)	Substrate concentration	Enzymatic
		Activity

Sol. Answer (3)

X-axis represent temperature and Y-axis represent enzyme activity. All enzymes act at an optimum temperature, above and below this temperature the enzyme activity declines.

- 157. The ciliated columnar epithelial cells in humans are known to occur in
 - (1) Fallopian tubes and urethra (2) Eustachian tube and stomach lining
 - (4) Dila desat and accombance
 - (3) Bronchioles and Fallopian tubes
- (4) Bile duct and oesophagus

Sol. Answer (3)

Ciliated columnar epithelium lines bronchioles and fallopian tubes.

- 158. Select the correct option with respect to mitosis
 - (1) Chromosomes move to the spindle equator and get alingned along equatorial plate in metaphase
 - (2) Chromatids separate but remain in the centre of the cell in anaphase
 - (3) Chromatids start moving towards opposite poles in telophase
 - (4) Golgi complex and endoplasmic reticulum are still visible at the end of prophase

Sol. Answer (1)

Chromatids show poleward movement in anaphase; golgi and ER disappears in late prophase.

159. Which one of the following structural formulae of two organic compounds is **correctly** identified along with its related function?

- (1) A: Lecithin a component of cell membrane
- (2) B: Adenine a nucleotide that makes up nucleic acids
- (3) A: Triglyceride major source of energy
- (4) B: Uracil a component of DNA

Sol. Answer (1)

A is the formula of Lecithin. Lecithin is a phospholipid and is the component of the cell membranes. B is the formula of adenine which is a nitrogenous base not a nucleotide.

160. What was the most significant trend in the evolution of modern man (Homo sapiens) from his ancestors?

(1) Increasing brain capacity

(2) Upright posture

(3) Shortening of jaws

(4) Binocular vision

Sol. Answer (1)

The most significant trend in the evolution of modern man (*Homo sapiens*) from the ancestors is increasing brain capacity.

- 161. Which one of the following conditions correctly describes the manner of determining the sex in the given example?
 - (1) Homozygous sex chromosomes (XX) produce make in Drosophila
 - (2) Homozygous sex chromosomes (ZZ) determine female sex in birds
 - (3) XO type of sex chromosomes determine male sex in grasshopper
 - (4) XO condition in humans as found in Turner Syndrome, determines female sex

Sol. Answer (3)

X/A ratio determines sex in *Drosophila*; \P is heterogametic (ZW) in birds. In 'XO type' the 'O' determines maleness.

- 162. A person with unknown blood group under ABO system, has suffered much blood loss in an accident and needs immediate blood transfusion. His one friend who has a valid certificate of his own blood type, offers for blood donation without delay. What would have been the type of blood group of the donor friend?
 - (1) Type A

(2) Type B

(3) Type AB

(4) Type O

Sol. Answer (4)

The person with blood group O is said to universal donor, because in this, there are no antigens on the surface of RBC.

- 163. What are those structures that appear as 'beads-on-string' in the chromosomes when viewed under electron microscope?
 - (1) Base pairs

(2) Genes

(3) Nucleotides

(4) Nucleosomes

Sol. Answer (4)

Nucleosome consist of octameric histone core wrapped by dsDNA.

164. Match the source gland with its respective hormone as well as the function

	Source gland	Hormone	Function		
(1)	Thyroid	Thyroxine	Regulates blood calcium level		
(2)	Anterior pituitary	Oxytocin	Contraction of uterus muscles during child birth		
(3)	Posterior pituitary	Vasopressin	Stimulates resorption of water in the distal tubules in the nephron		
(4)	Corpus luteum	Estrogen	Supports pregnancy		

Sol. Answer (3)

Vasopressin (ADH) is synthesised in hypothalamus but released into the blood from posterior lobe of pituitary, so it is called as a hormone of posterior lobe. It stimulates reabsorption of water in distal tubules in the nephron.

- 165. Which of the following is correctly stated as happens in the common cockroach?
 - (1) The food is ground by mandibles an gizzard
 - (2) Malpighian tubules are excretory organ projecting out from the colon
 - (3) Oxygen is transported by haemoglobin blood
 - (4) Nitrogenous excretory product is urea

Sol. Answer (1)

In cockroach the food is grinded by mandibles and gizzard. In insects there is no oxygen transporting pigment and nitrogenous excretory product is uric acid.

- 166. A large proportion of oxygen is left unused the human blood even after its uptake by the body tissues. This O₂
 - (1) Helps in releasing more O₂ to the epithelium tissues
 - (2) Acts as a reserve during muscular exercise
 - (3) Raises the pCO2 of blood to 75 mm of Hg
 - (4) Is enough to keep oxyhaemoglobin saturation at 96%

Sol. Answer (2)

Our tissues are able to utilise only 25% of O_2 carried by arterial blood. Our venous blood is still 75% saturated with O_2 . This O_2 acts as a reserve during muscular exercise.

Aakash Institute

167. Which one of the following enzymes carries on the initial step in the digestion of milk in humans?

(1) Trypsin

(2) Pepsin

(3) Rennin

(4) Lipase

Sol. Answer (2)

In humans milk protein digesting enzyme in stomach is pepsin. In calves it is rennin. Rennin is also present in small amounts in human infants but not adults. Pepsin acts on water soluble caseinogen (milk protein) to form solubles 'casein'. This combines with calcium salts to form insoluble calcium paracaseinate, which gets readily digested enzymatically.

168. Which one of the following is not a part of a renal pyramid?

(1) Loops of Henle

(2) Peritubular capillaries

(3) Convoluted tubules

(4) Collecting ducts

Sol. Answer (3)

In Bowman's capsule PCT and DCT are in renal cortex, whereas, loops of Henle are in medullary pyramids.

- 169. One very special feature in the earthworm pheretima is that
 - (1) It has a long dorsal tubular heart
 - (2) Fertilisation of eggs occurs inside the body
 - (3) The typhlosole greatly increases the effective absorption area of the digested food in the intestine
 - (4) The S-shaped setae embedded in the integument are the defensive weapons used against the enemies

Sol. Answer (3)

In earthworm, mid dorsal villi typhlosole greatly increases the effective absorption area of the digested food in the intestine.

170. Two friends are eating together on a dining table. One of them suddenly starts coughing while swallowing some food. This coughing would have been due to improper movement of

(1) Tongue

(2) Epiglottis

(3) Diaphragm

(4) Neck

Sol. Answer (2)

If a person suddenly starts coughing while swallowing food, it is due to improper movement of epiglottis. If the glottis is not properly closed some food can enter respiratory tract.

- 171. Arteries are best defined as the vessels which
 - (1) Carry blood from one visceral organ to another visceral organ
 - (2) Supply oxygenated blood to the different organs
 - (3) Carry blood away from the heart to different organs
 - (4) Break up into capillaries which reunite to form a vein

Sol. Answer (3)

Arteries are best defined as vessels which carry blood away from the heart to different organs.

172. 'Bundle of His' is a part of which one of the following organs in humans?

(1) Pancreas

(2) Brain

(3) Heart

(4) Kidney

Sol. Answer (3)

'Bundle of His' is a part of conducting system of human heart.

173. The purplish red pigment rhodopsin contained in the rods type of photoreceptor cells of the human eye, is a derivative of

(1) Vitamin A

(2) Vitamin B₁

(3) Vitamin C

(4) Vitamin D

Sol. Answer (1)

Vitamin A is the precursor of the purplish red pigment rhodopsin contained in the rods (photoreceptor) cells of human eye.

Aakash Institute CBSE Prelim. 2011 (Hints & Solutions) - Code A

174. Which one of the following plasma proteins is involved in the coagulation of blood?

(1) Fibrinogen

(2) An albumin

(3) Serum amylase

(4) A globulin

Sol. Answer (1)

Fibrinogen is a plasma protein involved in clotting of blood.

- 175. When a neuron is in resting state i.e. not conducting any impulse, the axonal membrane is
 - (1) Comparatively more permeable to K⁺ ions and nearly impermeable to Na⁺ ions
 - (2) Comparatively more permeable to Na⁺ ions and nearly impermeable to K⁺ ions
 - (3) Equally permeable to both Na⁺ and K⁺ ions
 - (4) Impermeable to both Na⁺ and K⁺ ions

Sol. Answer (1)

When a neuron is in resting state *i.e.*, not conducting any impulse, the axonal membrane is comparatively more permeable to K⁺ ions and nearly impermeable to Na⁺ ions.

- 176. Which one of following correctly explains the function of a specific part of a human nephron?
 - (1) Afferent arteriole: Carries the blood away from the glomerulus towards renal vein
 - (2) Podocytes: Create minute spaces (slit pores) for the filtration of blood into the Bowman's capsule
 - (3) Henle's loop: Most reabsorption of the major substances from the glomerular filtrate
 - (4) Distal convoluted tubule: Reabsorption of K⁺ ions into the surrounding blood capillaries

Sol. Answer (2)

Podocytes are specialised squamous epithelial cells in the inner wall of Bowman's capsule. They give rise to foot like processes which form filtration slits for the filtration of blood into the Bowman's capsule.

177. Given below is an incomplete table about certain hormones, their source glands and one major effect of each on the body in humans. Identify the correct option for the three blanks A, B and C

GLANDS	SECRETION	EFFECT ON BODY
А	Oestrogen	Maintenance of secondary sexual characters
Alpha cells of Islets of Langerhans	В	Raises blood sugar level
Anterior pituitary	С	Over secretion leads to gigantism

Options

	Α	В	С
(1)	Placenta	Glucagon	Calcitonin
(2)	Ovary	Glucagon	Growth hormone
(3)	Placenta	Insulin	Vasopressin
(4)	Ovary	Insulin	Calcitonin

Sol. Answer (2)

- A. Ovary secretes oestrogen for maintenance of secondary sexual characters.
- B. Alpha cells of Islets of Langerhans secrete glucagon which raises blood sugar level.
- C. Anterior lobe of pituitary secretes growth hormone. Its over secretion leads to gigantism.

- 178. Uricotelic mode of passing out nitrogenous wastes in found in
 - (1) Insects and Amphibians

(2) Reptiles and Birds


(3) Birds and Annelids

(4) Amphibians and Reptiles

Sol. Answer (2)

Reptiles and birds are uricotelic.

179. The figure given below shows a small part of human lung where exchange of gases takes place. In which one of the options given below, the one part **A**, **B**, **C** or **D** is **correctly** identified along with its function?



Options

- (1) **B**: Red blood cell transport of CO₂ mainly
- (2) C: Arterial capillary passes oxygen to tissues
- (3) A: alveolar cavity main site of exchange of respiratory gases
- (4) **D**: Capillary wall exchange of O₂ and CO₂ takes place here

Sol. Answer (3)

A is the alveolar cavity which is the main site of exchange of respiratory gases.

- 180. Which one of the following statements is correct regarding blood pressure?
 - (1) 190/110 mmHg may harm vital organs like brain and kidney
 - (2) 130/90 mmHg is considered high and requires treatment
 - (3) 100/55 mmHg is considered an ideal blood pressure
 - (4) 105/50 mmHg makes one very active

Sol. Answer (1)

Hypertension occurs if the blood pressure is 190/110. This can harm the vital organs like brain and kidneys.


- 181. Which one of the following statements is **correct** with respect to kidney function regulation?
 - (1) During summer when body loses lot of water by evaporation, the release of ADH is suppressed
 - (2) When someone drinks lot of water, ADH release is suppressed
 - (3) Exposure to cold temperature stimulates ADH release
 - (4) An increase in glomerular blood flow stimulates formation of Angiotensin II

Sol. Answer (2)

When someone drinks lot of water which is not required by his body, the osmolarity of the blood will decrease. The decrease in osmolarity will inhibit the release of ADH. ADH not released DCT becomes less permeable to water, and excess of water is eliminated.

Aakash Institute

182. The figure given below depicts a diagrammatic sectional view of the female reproductive system of humans. Which one set of three parts out of I – VI have been **correctly** identified?



- (1) (I) Perimetrium, (II) Myometrium, (III) Fallopian tube
- (2) (II) Endometrium, (III) Infundibulum, (IV) Fimbriae
- (3) (III) Infundibulum, (IV) Fimbriae, (V) Cervix
- (4) (IV) Oviducal funnel, (V) Uterus, (VI) Cervix

Sol. Answer (3)

III is infundibulum

IV is fimbriae

V is cervix

- 183. The testes in humans are situated outside the abdominal cavity inside a pouch called scrotum. The purpose served is for
 - (1) Providing a secondary sexual feature for exhibiting the male sex
 - (2) Maintaining the scrotal temperature lower than the internal body temperature
 - (3) Escaping any possible compression by the visceral organs
 - (4) Providing more space for the growth of epididymis

Sol. Answer (2)

The tests in humans are situated outside the abdominal cavity in scrotal sacs. This is because the temperature of scrotal sacs is 2.5°C lesser than internal body temperature.

- 184. Which one of the following is the most widely accepted method of contraception in India, as at present?
 - (1) IUDs' (Intra uterine devices)

(2) Cervical caps

(3) Tubectomy

(4) Diaphragms

Sol. Answer (1)

The most widely accepted method of contraception in India is IUDs.

- 185. If for some reason, the vasa efferentia in the human reproductive system get blocked, the gametes will not be transported from
 - (1) Vagina to uterus

(2) Testes to epididymis

(3) Epididymis to vas deferens

(4) Ovary to uterus

Sol. Answer (2)

The path of transport of gametes is Seminiferous tubules \rightarrow rete testis \rightarrow vasa efferentia \rightarrow epididymis. So, if vasa efferentia are blocked the gametes from testes will not enter epididymis.

Aakash Institute

- 186. Medical Termination of Pregnancy (MTP) is considered safe up to how many weeks of pregnancy?
 - (1) Six weeks

(2) Eight weeks

(3) Twelve weeks

(4) Eighteen weeks

Sol. Answer (3)


MTPs are considered safe upto twelve weeks of pregnancy.

- 187. Which one of the following is categorised as a parasite in true sense?
 - (1) The cuckoo (koel) lays its egg in crow's nest
 - (2) The female Anopheles bites and sucks blood from humans
 - (3) Human foetus developing inside the uterus draws nourishment from the mother
 - (4) Head louse living on the human scalp as well as laying eggs on human hair

Sol. Answer (4)

Head louse living on the human scalp as well as laying eggs on human hair is a parasite in true sense. Female mosquito is not considered as a parasite, though it needs human blood for reproduction. Koel that lays in crow's nest is just a brood parasite.

188. What type of human population is represented by the following pyramid?



(1) Expanding population

(4) Declining population

(3) Stable population

Sol. Answer (4)

It is an Urn shaped pyramid with least number of pre-reproductive individuals.

- 189. Which one of the following statements for pyramid of energy is incorrect, whereas the remaining three are correct?
 - (1) It is upright in shape
 - (2) Its base is broad
 - (3) It shows energy content of different trophic level organisms
 - (4) It is inverted in shape

Sol. Answer (4)

It is never inverted.

- 190. Ethanol is commercially produced through a particular species of
 - (1) Aspergillus

(2) Saccharomyces

(3) Clostridium

(4) Trichoderma

Sol. Answer (2)

Yeast species.

Aakash Institute CBSE Prelim. 2011 (Hints & Solutions) - Code A 191. Consider the following four conditions (a - d) and select the correct pair of them as adaptation to environment in desert lizards. The conditions (a) Burrowing in soil to escape high temperature (b) Losing heat rapidly from the body during high temperature (c) Bask in sun when temperature is low (d) Insulating body due to thick fatty dermis (1) (a), (b) (2) (c), (d) (3) (a), (c) (4) (b), (d) Sol. Answer (3) The adaptations in desert lizard are (i) burrowing in soil to escape high temperature (ii) bask in sun when temperature is low 192. Which one of the following pairs of gases are the major cause of "Greenhouse Effect"? (1) CO₂ and N₂O (2) CO_2 and O_3 (4) CFCs and SO₂ (3) CO₂ and CO Sol. Answer (1) CO₂, CH₄, N₂O and CFC are common green house gases. 193. Where will you look for the sporozoites of the malarial parasite? (1) Salivary glands of freshly moulted female Anopheles mosquito (2) Saliva of infected female Anopheles mosquito (3) Red blood corpuscles of humans suffering from malaria (4) Spleen of infected humans Sol. Answer (2) Sporozoites are the infective stage of malarial parasite. They present in the saliva of infected female Anopheles mosquito. 194. When two unrelated individuals or lines are corssed, the performance of F1 hybrid is often superior to both its parents. This phenomenon is called (1) Metamorphosis (2) Heterosis (3) Transformation (4) Spheing Sol. Answer (2) Heterosis is equivalent to hybrid vigour. 195. A certain patient is suspected to be suffering from Acquired Immuno Deficiency Syndrome. Which diagnostic technique will you recommend for its detection? (1) WIDAL (2) ELISA

Sol. Answer (2)

(3) MRI

ELISA is a diagnostic test for AIDS.

Ultra sound

CBSE	Prelim. 2011 (Hints & Solutions) - Code A			Aakash Institute	
196.	At which stage of HIV infection does one usually sho	ow s	symptoms of AIDS?		
	(1) Within 15 days of sexual contact with an infected person				
	(2) When the infecting retrovirus enters host cells				
	(3) When viral DNA is produced by reverse transcript	tase)		
	(4) When HIV replicates rapidly in helper T-lymphocy	ytes	and damages large number of these		
Sol.	Answer (4)				
	Symptoms of AIDs appear when there is depletion of	hel	per T-cells.		
197.	Given below is a sample of a portion of DNA strand is so special shown in it?	givir	ng the base sequence on the opposite	strands. What	
	5′ GAATTC 3′				
	3′ 5′				
	(1) Palindromic sequence of base pairs	(2)	Replication completed		
	(3) Deletion mutation ((4)	Start codon at the 5' end		
Sol.	Answer (1)				
	5' — GAATTC — 3'				
	3' — CTTAAG — 5'				
	is the palindromic sequence, recognised by EcoRI.				
198.	The most common substrate used in distilleries for the	he p	roduction of ethanol is		
	(1) Molasses ((2)	Corn meal		
	(3) Soya meal	(4)	Ground gram		
Sol.	Answer (1)				
	Molasses are used commonly in distilleries for ethan-	ol p	roduction		
199.	An organism used as biofertilizer for raising soyabea	ın cr	op is		
	(1) Nostoc	(2)	Azotobacter		
	(3) Azospirillum (-	(4)	Rhizobium		
Sol.	Answer (4)				
	Soyabean is a legume associated symbiotically with	Rh	izobium.		
200.	There is a restriction endonuclease called EcoRI. When	hat (does "co" part in it stand for?		
	(1) coli (5)	(2)	colon		
	(3) coelom ((4)	coenzyme		
Sol.	Answer (1)				
	In EcoRI, 'co' stands for coli (species of bacteria, from	m w	here it is obtained)		
)			
		-	_		