A

Register Number

# Part III - MATHEMATICS

(English Version)

Time Allowed : 3 Hours ]

[ Maximum Marks : 200

3521

## SECTION - A

N. B.: i) All questions are compulsory.

ii) Each question carries one mark.

iii) Choose the most suitable answer from the given four alternatives.  $40 \times 1 = 40$ 

1. If 
$$|\vec{a} + \vec{b}| = |\vec{a} - \vec{b}|$$
, then  
a)  $\vec{a}$  is parallel to  $\vec{b}$  b)  $\vec{a}$  is perpendicular to  $\vec{b}$   
c)  $|\vec{a}| = |\vec{b}|$  d)  $\vec{a}$  and  $\vec{b}$  are unit vectors.  
2. The shortest distance of the point (2, 10, 1) from the plane  
 $\vec{r} \cdot (3\vec{t} - \vec{j} + 4\vec{k}) = 2\sqrt{26}$  is

a)  $2\sqrt{26}$  b)  $\sqrt{26}$ c) 2 d)  $\frac{1}{\sqrt{26}}$ 

3. The point of intersection of the lines  $\frac{x-6}{-6} = \frac{y+4}{4} = \frac{z-4}{-8}$  and

2

$$\frac{x+1}{2} = \frac{y+2}{4} = \frac{z+3}{-2}$$
 is

(0,0,-4) a)

(1, 0, 0) b)

(0, 2, 0) c)

(1, 2, 0). d)

The projection of  $3\vec{i} + \vec{j} - \vec{k}$  on  $4\vec{i} - \vec{j} + 2\vec{k}$  is 4.

a)	$\frac{9}{\sqrt{21}}$	mark	Ties one	b)	$\frac{-9}{\sqrt{21}}$
c)	$\frac{81}{\sqrt{21}}$			d)	$\frac{-81}{\sqrt{21}}.$

The centre and radius of the sphere  $|\vec{r} - (2\vec{\iota} - \vec{j} + 4\vec{k})| = 5$  are 5.

(2, -1, 4) and 5 a)

(2, 1, 4) and 5 b)

(-2, 1, 4) and 6 c)

(2, 1, -4) and 5. d)

6.

The distance between the foci of the ellipse  $9x^2 + 5y^2 = 180$  is

4 a)

8

c)

A

2. d)

b)

6

of the holes (2. FO. L. L. Son

7. The directrices of the hyperbola  $x^2 - 4(y - 3)^2 = 16$  are

a)  $y = \pm \frac{8}{\sqrt{5}}$ b)  $x = \pm \frac{8}{\sqrt{5}}$ c)  $y = \pm \frac{\sqrt{5}}{8}$ d)  $x = \pm \frac{\sqrt{5}}{8}$ .

8. The point of intersection of tangents at  $t_1$  and  $t_2$  to the parabola  $y^2 = 4ax$  is

- a)  $\left[ a \left( t_1 + t_2 \right), a t_1 t_2 \right]$
- b)  $\begin{bmatrix} at_1 t_2, a(t_1 + t_2) \end{bmatrix}$
- c)  $[at^2, 2at]$

A

d)  $\begin{bmatrix} at_1 t_2, a(t_1 - t_2) \end{bmatrix}$ .

9. The slope of the tangent to the curve  $y = 3x^2 + 3 \sin x$  at x = 0 is

a) 3 b) 2 c) 1 d) -1.

10. The function  $f(x) = x^2$  is decreasing in

a)  $(-\infty, \infty)$ b)  $(-\infty, 0)$ c)  $(0, \infty)$ d)  $(-2, \infty)$ .

3521

11. The area between the ellipse  $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$  and its auxiliary circle is

4

- a)  $\pi b(a-b)$
- b)  $2\pi a(a-b)$
- c)  $\pi a(a-b)$
- d)  $2\pi b(a-b)$ .

12. The volume generated by rotating the triangle with vertices at (0, 0), (3, 0) and(3, 3) about *x*-axis is

a) 18π

b) 2π

d)

9π.

c) 36π

13.  $\int_{a}^{b} f(x) \, \mathrm{d}x$  is

A

a) 
$$2\int_{0}^{a} f(x) dx$$
  
b) 
$$\int_{a}^{b} f(a-x) dx$$
  
c) 
$$\int_{a}^{b} f(b-x) dx$$
  
d) 
$$\int_{a}^{b} f(a+b-x) dx$$

14. The integrating factor of  $\frac{dy}{dx} + 2\frac{y}{x} = e^{4x}$  is a)  $\log x$ **b**)  $x^2$ c) ex d) x. 15. The complementary function of  $(D^2 + 1)y = e^{2x}$  is  $(Ax + B)e^{x}$ a) b)  $A \cos x + B \sin x$  $(Ax+B)e^{2x}$ c) d)  $(Ax + B)e^{-x}$ . 16. If p is true and q is unknown, then a)  $\sim p$  is true b)  $p V (\sim p)$  is false c)  $p \wedge (\sim p)$  is true d) p V q is true. 17. If  $f(x) = \begin{cases} kx^2 & ; 0 < x < 3 \\ 0 & ; elsewhere. \end{cases}$ is a probability density function, then the value of k is  $\frac{1}{3}$ b)  $\frac{1}{6}$ a) c)  $\frac{1}{9}$ d)  $\frac{1}{12}$ . 18. Given E(x+c) = 8 and E(x-c) = 12, then the value of c is - 2 a) **b**) 4 c) - 4 **d**) 2.

A

[ Turn over

3521

5

19. In a Poisson distribution, if P(x=2) = P(x=3) then the value of its parameter  $\lambda$ -is

6

- 2 b) a) 6 d) 0. c) 3

d)

A

20. Which of the following are correct ? E(aX+b)=aE(X)+bΙ.  $\mu_2 = \mu_2' - (\mu_1')^2$ II.  $\mu_2 = Variance$ III. IV.  $Var(aX+b) = a^2 var(X)$ . I, II and III b) All a) d) I and IV. c) II and III 21. If  $A = \begin{bmatrix} 2 & 0 & 1 \end{bmatrix}$ , then the rank of  $AA^T$  is 2 b) a) 1 d) 0. 3 c) 22. If A is a matrix of order 3, then det ( kA ) is  $k^3 \det(A)$ a)  $k^2 \det(A)$ b) k det (A) c) det ( A ).

23.	If .	A = [	0 0	], then $A^{12}$	is	ndanu:	28igo		
•	a)	0	0 60			b)	0	0 5 <sup>12</sup> ]	ile
	cl	0	0 0			d)	[1	0	
	0,	Lo	0 ]				Lo	ıſ	

24. In a homogeneous system  $\rho$  ( *A* ) < ( the number of unknowns ) then the system has

a) only trivial solution

b) trivial solution and infinitely many non-trivial solutions

c) only non-trivial solutions

d) no solution.

25. If  $\overrightarrow{a}$  and  $\overrightarrow{b}$  include an angle 120° and their magnitudes are 2 and  $\sqrt{3}$ , then  $\overrightarrow{a}$ ,  $\overrightarrow{b}$  is equal to

a) √3

b) -√3

c) 2

d)  $-\frac{\sqrt{3}}{2}$ .

26. If  $x^2 + y^2 = 1$  then the value of  $\frac{1 + x + iy}{1 + x - iy}$  is

a) x - iy b) 2x

c) -2iy d) x + iy.

A

27. The polar form of the complex number  $(i^{25})^3$  is

8

- a)  $\cos\frac{\pi}{2} + i\sin\frac{\pi}{2}$
- b)  $\cos \pi + i \sin \pi$
- c)  $\cos \pi i \sin \pi$
- d)  $\cos\frac{\pi}{2} i\sin\frac{\pi}{2}$ .

28. The value of  $i + i^{22} + i^{23} + i^{24} + i^{25}$  is

- a) i b) i
  - c) 1 d) 1.

29. Which of the following is incorrect regarding  $n^{\text{th}}$  roots of unity ?

- a) The number of distinct roots is n
- b) The roots are in G.P. with common ratio  $cis\left(\frac{2\pi}{n}\right)$

c) The arguments are in A.P. with common difference  $\frac{2\pi}{n}$ 

d) Product of the roots is 0 and the sum of the roots is  $\pm 1$ .

30. If the line 4x + 2y = c is a tangent to the parabola  $y^2 = 16x$  then c is

a) -1 b) -2

c)

A

4

d) - 4.

31. Which of the following curves is concave down?

a)  $y = -x^2$ b)  $y = x^2$ c)  $y = e^x$ d)  $y = x^2 + 2x - 3$ .

32. One of the conditions of Rolle's theorem is

- a) f is defined and continuous on (a, b)
- b) f is differentiable on [a, b]

c) f(a) = f(b)

- d) f is differentiable on (a, b].
- 33. If  $u = \frac{1}{\sqrt{x^2 + y^2}}$ , then  $x \frac{\partial u}{\partial x} + y \frac{\partial u}{\partial y}$  is equal to a)  $\frac{1}{2}u$  b) uc)  $\frac{3}{2}u$  d) -u.

34. If  $x = r \cos \theta$ ,  $y = r \sin \theta$ , then  $\frac{\partial r}{\partial x} =$ 

- a)  $\sec \theta$  b)  $\sin \theta$
- c)  $\cos \theta$  d)  $\csc \theta$ .  $\pi/4$
- 35. The value of  $\int_{0}^{\pi/4} \cos^3 2x \, dx$  is
  - a)  $\frac{2}{3}$
  - c) 0

A

b)  $\frac{1}{3}$ d)  $\frac{2\pi}{3}$ .

3521

36. Solution of  $\frac{dx}{dy} + mx = 0$ , where m < 0 is

a)  $x = ce^{-my}$  b)  $x = ce^{-my}$ 

c) x = my + c d)

37. The order and degree of the differential equation  $\frac{d^2 y}{dx^2} - y + \left(\frac{dy}{dx} + \frac{d^3 y}{dx^3}\right)^{3/2} = 0$ 

x = c.

are

- a) 2, 3b) 3, 3c) 3, 2d) 2, 2.
- 38. Which of the following is a tautology?
  - a)  $p \vee q$  b)  $p \wedge q$
  - c)  $p \lor \sim p$  d)  $p \land \sim p$ .

39. Which of the following is not a binary operation on R?

- a) a \* b = ab
- b) a \* b = a b
- c)  $a * b = \sqrt{ab}$
- d)  $a * b = \sqrt{a^2 + b^2}$ .

40. The value of  $[3] + {}_{11}([5] + {}_{11}[6])$  is

a) [0] b) [1] c) [2] d) [3].

A

## SECTION - B

N. B.: i) Answer any ten questions.

- ii) Question No. 55 is compulsory and choose any nine questions from the remaining.
- iii) Each question carries six marks.  $10 \times 6 = 60$

41. Solve the following system of linear equations by determinant method.

$$2x - 3y = 7$$
,  $4x - 6y = 14$ .

42. If  $A = \begin{bmatrix} 1 & 2 \\ 1 & 1 \end{bmatrix}$  and  $B = \begin{bmatrix} 0 & -1 \\ 1 & 2 \end{bmatrix}$ , verify that  $(AB)^{-1} = B^{-1}A^{-1}$ .

- 43. Find the equation of the sphere on the join of the points A and B having position vectors  $2\overrightarrow{i} + 6\overrightarrow{j} 7\overrightarrow{k}$  and  $2\overrightarrow{i} 4\overrightarrow{j} + 3\overrightarrow{k}$  and respectively as a diameter.
- 44. i) A force of magnitude 5 units acting parallel to  $2\vec{i} 2\vec{j} + \vec{k}$  displaces the point of application from (1, 2, 3) to (5, 3, 7). Find the work done.
  - ii) The volume of a parallelopiped whose edges are represented by  $-12\vec{i} + \lambda \vec{k}, \ 3\vec{j} \vec{k}, \ 2\vec{i} + \vec{j} 15\vec{k}$  is 546. Find the value of  $\lambda$ .
- 45. Find the square root of 7 + 24*i*.
- 46. Solve the equation  $x^4 4x^3 + 11x^2 14x + 10 = 0$ , if one root is 1 + 2i.

47. Evaluate  $\lim_{x \to 1} x^{\frac{1}{x-1}}$ .

A

48. i) Verify Rolle's theorem for the function  $f(x) = \sin x$ ,  $0 \le x \le \pi$ .

ii) Prove that  $e^x$  is strictly increasing function on R.

49. If 
$$z = ye^{x^2}$$
, where  $x = 2t$ ,  $y = 1 - t$ , then find  $\frac{dz}{dt}$ .

50. Evaluate 
$$\int_{0}^{0} \frac{\sqrt{x}}{\sqrt{x} + \sqrt{3} - x} \, \mathrm{d}x.$$

3

3521

51. Solve 
$$\frac{\mathrm{d}y}{\mathrm{d}x} + 2y \tan x = \sin x$$
.

52. Prove that  $p \leftrightarrow q \equiv (p \rightarrow q) \land (q \rightarrow p)$ .

- 53. A game is played with a single fair die. A player wins Rs. 20 if a 2 turns up, Rs. 40 if a 4 turns up, loses Rs. 30 if a 6 turns up, while he neither wins nor loses if any other face turns up. Find the expected sum of money he can win.
- 54. Four coins are tossed simultaneously. What is the probability of getting
  - (i) exactly 2 heads, (ii) at least 2 heads, and (iii) at most 2 heads ?
  - 55. a) Let G be a group,  $a, b \in G$ . Then prove that  $(a * b)^{-1} = b^{-1} * a^{-1}$ .

#### OR

b) Find the equations of the tangent and normal to the parabola

 $x^{2} + 2x - 4y + 4 = 0$  at the point (0, 1).

### SECTION - C

N. B.: i) Answer any ten questions.

- ii) Question No. 70 is compulsory and choose any nine questions from the remaining.
- iii) Each question carries ten marks.  $10 \times 10 = 100$
- 56. Examine the consistency of the following system of equations. If it is consistent, solve them. (Use rank method)

x + y - z = 1, 2x + 2y - 2z = 2, -3x - 3y + 3z = -3.

- 57. Show that the lines  $\frac{x-1}{3} = \frac{y-1}{-1} = \frac{z+1}{0}$  and  $\frac{x-4}{2} = \frac{y}{0} = \frac{z+1}{3}$  intersect and hence find the point of intersection.
- 58. Find the vector and cartesian equations of the plane passing through the points with position vectors  $3\overrightarrow{i} + 4\overrightarrow{j} + 2\overrightarrow{k}$ ,  $2\overrightarrow{i} 2\overrightarrow{j} \overrightarrow{k}$  and  $7\overrightarrow{i} + \overrightarrow{k}$ .

A

- 60. On lighting a rocket cracker it gets projected in a parabolic path and reaches a maximum height of 4 m when it is 6 m away from the point of projection. Finally it reaches the ground 12 m away from the starting point. Find the angle of projection.
- 61. Find the eccentricity, centre, foci and vertices of the ellipse

 $16x^2 + 9y^2 - 32x + 36y - 92 = 0$  and draw the diagram.

13

<sup>59.</sup> Solve  $x^7 + x^4 + x^3 + 1 = 0$ .

62. Prove that the line 5x + 12y = 9 touches the hyperbola  $x^2 - 9y^2 = 9$  and find

its point of contact.

63. Find the intervals of concavity and the points of inflexion of the curve

$$u = 12x^2 - 2x^3 - x^4.$$

64. Trace the curve  $y = x^3 + 1$ .

- 65. Find the area between the curve  $y = x^2 x 2$ , x-axis and the lines x = -2and x = 4.
- 66. Prove that the curved surface area of a sphere of radius r intercepted between two parallel planes at the distances a and b from the centre of the sphere is  $2\pi r(b-a)$  and hence deduce the surface area of the sphere (b > a).
- 67. Radium disappears at a rate proportional to the amount present. If 5% of the original amount disappears in 50 years, how much will remain at the end of 100 years? [Take  $A_0$  as the initial amount ]
- 68. Show that the set { [1], [3], [4], [5], [9] } forms an Abelian group under multiplication modulo 11.
- 69. Find c,  $\mu$  and  $\sigma^2$  of the normal distribution whose probability density function is given by  $f(x) = ce^{-x^2 + 3x}, -\infty < x < \infty$ .

A

15

3521

70. a) Solve:  $(D^2 - 1)y = \cos 2x - 2 \sin 2x$ .

A

## OR

b) A car A is travelling from west to east at 50 km/hr and car B is travelling from south towards north at 60 km/hr. Both are headed for the intersection of the two roads. At what rate are the cars approaching each other when car A is 0.3 km and car B is 0.4 km from the intersection ?