#### Subject: - Engineering Chemistry and Environmental Science Section: - B CORROSION AND LUBRICANTS

- 1. Volatile oxidation corrosion product of a metal is,
  - A.  $Fe_2O_3$
  - B. MoO<sub>3</sub>
  - C. Fe<sub>3</sub>O<sub>4</sub>
  - D. FeO
- 2. Lower is P<sup>H</sup>, corrosion is,
  - A. Greater
  - B. Lower
  - C. Constant
  - D. None of above
- 3. Electrochemical corrosion takes place on,
  - A. Anodic area
  - B. Cathodic area
  - C. Near cathode
  - D. Near anode
- 4. Chemical formula of Rust is,
  - A.  $Fe_2O_3$
  - B. FeO
  - C.  $Fe_3O_4$
  - D. Fe<sub>2</sub>O<sub>3-x</sub>H<sub>2</sub>O
- 5. Which of following metals could provide cathodic protection to Fe?
  - A. Al & Cu
  - B. Al & Zn
  - C. Zn & Cu
  - D. Al & Ni
- 6. Smaller the grain size, corrosion is,
  - A. Greater
  - B. Lower
  - C. Constant
  - D. Doesn't affected
- 7. Process of corrosion enhanced by,
  - A. AIR & Moisture
  - B. Electrolytes in water
  - C. Metallic impurities
  - D. Gases like CO<sub>2</sub> & SO<sub>2</sub>
  - E. All of above.

- 8. Standard electrode potential of hydrogen is,
  - A. 1.00 V
  - B. 0.00 V
  - C. 0.01 V
  - D. 0.001 V
- 9. Standard electrode potential of Al / Al<sup>3+</sup> is,
  - A. +0.66V
  - B. 0.66 V
  - C. 1.66 V
  - D. + 1.66 V
- 10. Standard electrode potential of Zn<sup>2+</sup>/ Zn is,
  - A. 0.76 V
  - B. +0.76 V
  - C. 2.76 V
  - D. + 2.76 V
- 11. Which of the following gases accelerates rusting of iron?
  - A.  $CO_2$
  - B. SO<sub>2</sub>
  - C. NO<sub>2</sub>
  - D. All of above
- 12. Standard electrode potential of Fe<sup>2+</sup>/Fe is,
  - A. + 0.44 V
  - B. 0.44 V
  - C. + 1.44 V
  - D. 1.44 V
- 13. Which of the following metal does not resists the corrosion process?
  - A. Ni
  - B. Cu
  - C. Pb
  - D. Fe
- 14. Viscosity index can be defined as,
  - A. (L-U/L-H) X 100
  - B.  $(L+U/L-H) \times 100$
  - C.  $(L-H/L-U) \times 100$
  - D.  $(L-H/L+U) \times 100$
- 15. Identify liquid lubricant from the following,
  - A. Graphite
  - **B.** Lubricating oil
  - C. Mo Disulphide
  - D. Soapstone

- 16. Identify the odd type of lubricant from following,
  - A. Semi solid
  - B. Liquid
  - C. Gaseous
  - D. Solid
- 17. Graphite is \_\_\_\_\_ lubricant.
  - A. Solid
  - B. Liquid
  - C. Semi solid
  - D. None of above.
- 18. Antiwear additive in extreme pressure additive is,
  - A. Soapstone
  - **B.** Tricresyl phosphate
  - C. Mica
  - D. Graphite
- 19. Calcium soap base grease is also called as,
  - A. Soap grease
  - B. Greases
  - C. Axle grease
  - D. Cup grease
- 20. Ethanol is used as,
  - A. Polymeric thickener
  - B. Polymer
  - C. Polymeric thinner
  - D. None of above.
- 21. What type of oil is suitable for thick film lubrication?
  - A. Hydrocarbon oil
  - B. Mineral oil
  - C. Polymeric oil
  - D. None of above.
- 22. What type of lubrication is used in delicate machines like watches, sewing machines, etc?
  - A. Fluid film lubrication
  - B. Extreme lubrication
  - C. Boundary lubrication
  - D. Thin film lubrication
- 23. Minimum separating distance in hydrodynamic lubrication is,
  - A.  $100 \text{ A}^0$
  - B. 1000 A<sup>0</sup>
  - C.  $10000 \text{ A}^0$
  - D.  $10 \, \text{A}^0$

- 24. Coefficient of friction in fluid film or hydrodynamic lubrication is,
  - A. 0.1 to 0.3
  - B. 0.01 to 0.03
  - C. 0.001 to 0.03
  - D. None of above.
- 25. Coefficient of friction for boundary lubrication is,
  - A. 0.5 to 15
  - B. 0.005 to 0.05
  - C. 0.5 to 0.15
  - D. 0.05 to 0.15
- 26. Special additives added to mineral oils are known as,
  - A. Extreme pressure additives
  - B. Special additives
  - C. Mineral additives
  - D. Lubricating additives
- 27. Important extreme pressure additives contains functional group from following,
  - A. Chlorides
  - B. Sulphides
  - C. Phosphides
  - D. All of above.
- 28. Animal and vegetable oils possess,
  - A. Good oiliness
  - B. Poor oiliness
  - C. Optimum oiliness
  - D. None of above.
- 29. Animal and vegetable oils are also used as,
  - A. Oiliness carrier
  - **B.** Blending agent
  - C. Solvent agent
  - D. Extreme pressure additives
- 30. Length of hydrocarbon chain in petroleum oils varies between about,
  - A. 0 to 12 Carbon atoms
  - B. 0 to 50 Carbon atoms
  - C. 21 to 30 Carbon atoms
  - D. 12 to 50 Carbon atoms
- 31. Shorter the chain of petroleum oil,
  - A. Lower viscosity
  - B. Higher viscosity
  - C. Softer
  - D. None of above.

- 32. Example of mineral / petroleum oil is or are,
  - A. Oleic acid
  - B. Stearic acid
  - C. Oxalic acid
  - D. Acetic acid
- 33. Purification of petroleum oil contains following sequence as,
  - A. Dewaxing, solvent refining, acid refining
  - B. Dewaxing, acid refining, solvent refining
  - C. Acid refining, dewaxing, solvent refining
  - D. Solvent refining, acid refining, dewaxing.
- 34. Acid refining step used to purify petroleum oil contains the use of,
  - A. Conc. H<sub>2</sub>SO<sub>4</sub>
  - B. Conc. HCl
  - C. Dil. H<sub>2</sub>SO<sub>4</sub>
  - D. Conc. HNO<sub>3</sub>
- 35. In acid refining step which used to purify petroleum oil, comprises removal of excess Conc. H<sub>2</sub>SO<sub>4</sub> with addition of calculated quantity of,
  - A. KOH
  - B. NH<sub>4</sub>OH
  - C. Ca(OH)<sub>2</sub>
  - D. NaOH
- 36. Solvent refining step in purification of petroleum oil comprises use of solvent,
  - A. Furfural
  - B. Dichloroethyl ether
  - C. Nitrobenzene
  - D.  $SO_2$  + benzene
  - E. All of above
- 37. In solvent refining we can observe following layers in purification of petroleum oil,
  - A. Oil
  - **B.** Solvent
  - C. Impurity
  - D. None of above.
- 38. Blended oils can be prepared by mixing petroleum oil with,
  - A. Waxy substance
  - B. Different solvents
  - C. Specific additives
  - D. None of above

39. To increase oiliness of petroleum oil following which acid is not used,

	A.	Palmitic acid
	B.	Stearic acid
	C.	Acetic acid
	D.	Oleic acid
40. T	o inc	crease viscosity index of petroleum oil which one of the following is used?
	A.	Hexanol
	B.	Methanol
	C.	Ethanol
	D.	2-methoxy phenol
41. T	To de	crease pour point of petroleum oil which is used from following?
	A.	Ether
	В.	Acetone
	C.	Decane
	D.	Phenol
42.		is used as abrasion inhibitor.
		Tribromyl sulphate
		Tricresyl napthanate
		Tricresyl phosphate
		None of above
43.		is used as emulsifier.
		Sodium salt of sulphonic acid
		Sodium salt of sulphuric acid
		Sodium salt of sulphurous acid
		Sodium salt of di-sulphonic acid
44.		ioxidants may be,
		Aromatic
		Phenolic
		Amino compounds
		All of above
45. 7	-	event corrosion phenomenon, combination of is used,
		Phosphorous or Antimony
		Phosphorous or Chlorine
		Phosphorous or Nitrogen
		Antimony or Sodium
	_	revent heat resistance property of grease, inorganic solid thickening agents are used from the
f	ollov	
		Finely divided clay
		Bentonite
		Colloidal silica
	D.	Carbon black

E. All of above
47. Main function of soap is,
A. Thickening agent
B. Soapy agent
C. Lathering agent
D. Corrosion preventer
48. Axle greases can be formed by adding to resin and fatty oils.
A. Lime
B. Calcium
C. soda
D. Lithium
49. Graphite consists of structure.
A. Multitude layered
B. Trigonal
C. Orthorhombic
D. Spiral
50. Graphite doesn't oxidize in air below temperature
A. 375°C
B. 300°C
C. 500 <sup>0</sup> C
D. 1057 <sup>0</sup> C
51. When graphite is dispersed in water, is called,
A. Oil dug
B. Aqua dug
C. Graphia-water
D. None of above
52. Mo disulphide possesses very low
A. Coefficient of friction
B. Lubricating property
C. Frictional resistance
D. Conductivity
53. Mo disulphide is stable in air up to temperature.
A. 200° C
B. 300 <sup>0</sup> C
C. 400° C
D. 375 <sup>0</sup> C
54. Unit of viscosity is,
A. Ohm
B. Ohm <sup>-1</sup>
C. Poise
D. Cm <sup>-1</sup>

- 55. Low viscosity standard oil is,
  - A. Petroleum oil
  - B. Gulf oil
  - C. Blended oil
  - D. Mineral oil
- 56. High viscosity standard oil is,
  - A. Blended oil
  - B. Gulf oil
  - C. Pennylsylvanian oil
  - D. Petroleum oil
- 57. Viscosities of lubricating oil can be measured by
  - A. Ostwald' viscometer
  - B. Redwood viscometer
  - C. Saybolt viscometer
  - D. Ubbelhode viscometer
- 58. If viscosity of lubricating oil is measured by redwood viscometer, the flow time can be expressed as,
  - A. 100 Redwood seconds at 20<sup>0</sup> C
  - B. 100 Redwood viscometer working at  $T = 20^{\circ}$  C
  - C. At  $T = 20^{\circ}$  C, flow time = 100 seconds
  - D. At 20<sup>o</sup> C, 100 Redwood seconds
- 59. Redwood viscometer No. 1 is used to determine viscosities of,
  - A. Lubricating Oils
  - B. Thin lubricating oils
  - C. Highly viscous oils
  - D. None of above
- 60. Apparatus used to determine flash and fire points of lubricating oil is known as,
  - A. Bomb calorimeter
  - B. Spectrophotometer
  - C. Redwood viscometer
  - D. Pensky Martin Apparatus
- 61. In pour point apparatus the freezing mixture used is,
  - A.  $Ice + CaCl_2$
  - B.  $Ice + BaCl_2$
  - C.  $Ice + MgCl_2$
  - D.  $Ice + AlCl_3$
- 62. Lubricating oil should possess acid value,
  - A. More than 0.1
  - B. 0.1
  - C. Less than 0.1
  - D. None of above

63. Acid value is also called as,

A. Acid number
B. Neutralization number
C. Base number
D. Basic number
64. In differential aeration corrosion, poor oxygenated parts are,
A. Anodic
B. Cathodic
C. Corroded
D. None of above
65. Pitting corrosion is a accelerated attack.
A. Non-localized
B. Localized
C. Diverse
D. None of above
66. In waterline corrosion highly oxygenated parts acts as,
A. Cathodic
B. Anodic
C. Corroded
D. None of above
67. Paints which are used to restrict corrosion are known as,
A. Anticorrosion paints
B. Antifouling paints
C. Special paints
D. Marine paints
68. Caustic embrittlement, a type of stress corrosion contains deposition of corrosion products,
A. $Na_2FeO_2$
B. NaFeO <sub>2</sub>
C. $K_2FeO_2$
D. $Fe_3O_4$
69. In galvanic series, a metal high in series is more
A. Anodic
B. Cathodic
C. Corroded
D. None f above
70. Hydrogen overvoltage is
A. + 0.33 V
B 0.33 V
C 1.33 V
D. $+ 1.33 \text{ V}$

- I	the state of the s
71 Dot	tio of volumes of metal oxides to metal is known as,
	Specific mass ratio
	Volume ratio
	Specific ratio
	Specific volume ratio
	ative humidity above which atmospheric corrosion rate of metal increases sharply is known as,
	Critical humidity
	Humidity rate
	Environmental humidity
	Atmospheric humidity
	ore active metal used in sacrificial anodic protection method is known as,
	Sacrificial anode
В.	Sacrificial cathode
C.	Active anode
D.	Active cathode
74. Ide	ntify the metal which is not employed as Sacrificial anode,
A.	Mg
B.	Zn
C.	Al
D.	Na
75. Idea	ntify the group which is not used as anodic inhibitor,
A.	Chromates
B.	Phosphates
C.	Sulphates
D.	Tungstates
76. Coa	ating of Zn, Al and Cd on steel are, because their electrode potentials are lower.
A.	Cathodic
В.	Anodic
C.	Not affecting
D.	None of above
77	is used for producing a coating of low melting metal such as Zn, Sn, Pb, Al on Fe, steel
and	l Cu.
	Hot dipping
	Anodic coating
	Cathodic coating
	Galvanizing
	is the process of coating Fe or steel with a zinc coating.
	Tinning
В.	Hot dipping

C. GalvanizingD. None of above

79.		is the process of coating of tin over Fe or steel.
	A.	Tinning
	B.	Galvanizing
	C.	Metal cladding
	D.	Sheardizing
80.		coating is non toxic in nature.
	A.	Sn
	B.	Zn
	C.	Fe
	D.	Cu
81.	Oxi	dation potential of Ni / Ni <sup>2+</sup> is,
	A.	+ 0.284 V
	B.	- 0.284 V
	C.	+ 2.284 V
	D.	- 2.284 V
82.	Fe o	or steel is with respect to copper.
	A.	Anodic
	B.	Cathodic
	C.	Corrosive
	D.	Non corrosive
83.	Al is	s than Zn.
	A.	Less anodic
	B.	More anodic
	C.	Less Cathodic
	D.	More Cathodic
84.	Zn i	s more than Fe.
		Electronegative
		Corrosive
	C.	Electropositive
	D.	None of above
85.	Req	uired potential for protecting metal / alloy can be obtained from its
	A.	Potential current curve
	B.	Protecting curve
	C.	Potential curve
	D.	None of above
86.	Coa	ting applied must be chemically to the environment.
	A.	Inert
	B.	Reactive
	C.	Soluble
	D.	Non reactive

87. Aci	dic media are more corrosive than and neutral media.	
A.	Less acidic	
В.	Alkaline	
C.	Inert	
D.	Non reactive	
88. Reduction in over voltage of corroding metal / alloy accelerates the		
A.	Reactivity	
В.	Inertness	
C.	Corrosion	
D.	Reduction	
89. Pre	sence of silicate anions leads to formation of reaction products.	
A.	Soluble	
B.	Poisonous	
C.	Corrosive	
D.	Insoluble	
90. Evo	olution of hydrogen type corrosion occurs in environment.	
A.	Acidic	
B.	Neutral	
C.	Basic	
D.	Alkaline	
91. And	odic reaction involves dissolution of metal as corresponding metallic ions with liberation of	
A.	Pair of electron	
В.	Free electron	
C.	Ions	
D.	Current in electrolytic solution.	
92. Des	struction of metal starts	
A.	At the surface	
B.	Just on layer below from surface	
C.	In the middle	
D.	At the bottom	
93. Co	rosion is a process reverse of of metal.	
A.	Destruction	
В.	Extraction	
C.	Rusting	
D.	Galvanizing	
94. Green film of basic carbonate on surface of Cu contains CuCO <sub>3</sub> and		
A.	$BaCO_3$	
B.	$Ba(OH)_2$	
C.	$Cu(OH)_2$	
D.	CuO	

95. Red	dish scale of iron oxide has molecular formula
A.	$Fe(OH)_3$
B.	$Fe_2O_3$
C.	$Fe_3O_4$
D.	FeO
96. Fro	n the following, which is inert to oxidation?
A.	Cu
B.	Fe
C.	Steel
D.	
97. Con	version of Fe to Fe <sup>2+</sup> is,
A.	Oxidation
	Reduction
	Corrosion
	None of above
	rosion process is nothing but
	Reduction
	Oxidation
	Protection
	None of above
	s above and closely adjacent to waterline are
	Protected
	Anodic
	Cathodic
	Inert to environment
	usting of iron is catalyzed by which of the following?
	Iron
	Oxygen
	Zinc
D.	$\mathbf{H}^{+}$
	ENVIRONMENTAL CHEMISTRY
101.	Pollutant is
	Undesirable foreign matter
	Desirable foreign matter
	Required foreign matter
	Useful foreign matter
102.	Smog is mixture of fog and
	Gases
	Smoke
	Pollutants

	D.	Oxides	
103.		From the following, identify the gas which is not responsible for acid rain.	
	A.	$SO_2$	
	В.	$NO_2$	
	C.	HCl	
	D.	Ozone	
104.		Photochemical smog is formed by combination of nitrogen oxide and	
	A.	Hydrocarbons	
	B.	Smoke	
	C.	Fog	
	D.	$SO_2$	
105.			
	A.	0.05 <sup>0</sup> C / Year	
	B.	$0.05^0 \mathrm{C}$ / Month	
	C.	$0.15^0 \mathrm{C}$ / Year	
	D.	0.05 <sup>0</sup> C / Decade	
106.		From the following, identify the metal which can't pollute environment.	
	A.	As	
	B.	Pb	
	C.	Pt	
	D.	Hg	
107.		How much air does a man normally inhale in a day?	
	A.	16 Kg	
	B.	15 Kg	
	C.	10 Kg	
	D.	Vary according to season	
108.		Effect of increasing CO <sub>2</sub> in air is	
	A.	Heating	
	B.	Cooling	
		Increasing pollution	
	D.	None of above	
109.		Naturally CO <sub>2</sub> is removed from air by	
	A.	Metallic reaction	
		Photosynthesis	
	C.	Forestation	
	D.	Deforestation	
110.		Most dangerous pollutant emitted in air during incomplete combustion of fuels is	
	A.	CO	
	B.	$\mathrm{CO}_2$	
		$NO_2$	
	D	CH.	

111.	The disease caused by presence of particulate pollutants in atmosphere is		
A.	Leukemia		
В.	Bronchial asthma		
C.	Anemia		
D.	Lung's cancer		
112.	A part of atmosphere from where ozone concentration has been depleted is known as		
A.	Biosphere		
B.	Ozone layer		
C.	Ozone hole		
D.	Ozone rich layer		
113. A measure of oxidisable impurities present in the sewage is			
A.	COD		
B.	BOD		
C.	Atmospheric oxygen		
D.	Ozone		
114.	When anaerobic decomposition continues the sewage is called as		
A.	Septic		
В.	Stale		
C.	Dirty		
D.	Offensive		
115.	When aerobic decomposition continues the sewage is called as		
	Septic		
	Stale		
	Dirty		
	None of above		
116.	The depth in soil below which soil particles are filled with water only known as		
	Water table		
	Water layer		
	Water quantity		
	Depth of water		
117.	Skin cancer is also called as		
	Melanoma		
	Anemia		
	Leukemia		
	None of above		
118.	Radioactive wastes disposed off in salt mines, because		
	Salt absorbs it		
	Salt reflects it		
	Salt decreases the intensity of it		
D.	None of above		

119.	Environment includes air, water, land and
A.	Gases
В.	Biota
C.	Segments of Environment
D.	None of above
120.	Soil, air and water come under environment.
A.	Physical
B.	Biotic
C.	Living
D.	Meteorological
121.	Climatic factors like temperature, sunlight, humidity may also be referred as,
A.	Climatic Environment
B.	Physical Environment
C.	Biotic Environment
D.	Meteorological Environment
122.	Water can be placed in segment of environment.
A.	Lithosphere
	Hydrosphere
	Biosphere
D.	Atmosphere
123.	
A.	
	2.4
	0.4
	0.6
124.	Z 1 1
	0.5
	6.0
	0.06
125.	The part of earth, upwards at least to a height of 10 Km is
	Atmosphere
	Biosphere
	Hydrosphere Name of charge
	None of above  Port of earth surrounding our to nearly 500 Km shows from earth's surface is
126.	Part of earth surrounding up to nearly 500 Km above from earth's surface is
	Atmosphere
	Biosphere
	Lithosphere
<b>D</b> .	Environment

127.		Region 20 - 40 Km above earth's surface is
	A.	Atmosphere
	B.	Biosphere
	C.	Environment
	D.	Ozonosphere
128.		Weight of atmosphere is about tones.
	A.	$5 \times 10^{10}$
	B.	$5 \times 10^9$
	C.	$5 \times 10^{11}$
	D.	$5 \times 10^{12}$
129.		At earth's surface density of air is $g / cm^3$ .
	A.	0.13
		0.0013
	C.	0.013
	D.	0.3100
130.	•	Region at about $40 - 100$ Km above earth's surface which contains charged particles is called as
-		Ozonosphere
		Ionosphere
		Charge sphere
		Atmosphere
131.		A protective layer which absorbs harmful ionizing radiations like cosmic and x-rays is
131.		Ozonosphere
		Atmosphere
		Protectosphere
		Ionosphere
132.		80 % of earth's surface is covered by water, so it called as
102.		Hydro planet
		Aqua planet
		Blue planet
		Special planet
133.		Layer which absorbs harmful UV radiations falling on earth from sun is known as
		Ionosphere
		Environment
	C.	Ozonosphere
	D.	Atmosphere
134.		By volume CO <sub>2</sub> is present in atmosphere is only %.
	A.	0.93
	B.	0.03
		0.3
	D.	0.02

135.	Saline water present in hydrosphere segment of environr	nent is about	%.
A.	95		
B.	2.4		
C	97		
D.	96		
136.	Radiations received from sun are nothing but	energy.	
$\mathbf{A}$ .	Solar		
В.	Photochemical		
C.	Natural		
D.	Environmental		
137.	The layer which is just up to 6.4 Km above the earth is _	·	
A.	Biosphere		
B.	Atmosphere		
C.	Ozonosphere		
D.	Troposphere		
138.	Upper portion of troposphere is known as,		
A	Stratosphere		
В.	Biosphere		
C.	Ozonosphere		
D.	Ionosphere		
139.	Air pollution mainly concerns state of		
A.	Atmosphere		
B.	Ozonosphere		
C.	Troposphere		
D.	Stratosphere		
140.	Presence of SO <sub>2</sub> doesn't responsible for the	_ disease.	
A.	Cardiac		
B.	Respiratory		
C.	Leukemia		
D.	Pulmonary		
141.	From the following, identify the acid which is not respon	nsible for acid rain.	
A.	Acetic acid		
B.	Sulphurous acid		
C.	Nitrous acid		
D.	Nitric acid		
142.	Which acid is responsible for acid rain?		
A.	$H_2SO_3$		
B.	HCl		
C.	$C_2H_2O_4$		
D.	Salicylic acid		

143.		Quantity of oxygen available to body cells when reduced it is generally known as,
	A.	Hemophilia
	B.	Leukemia
	C.	Anemia
	D.	Anoxia
144.		Carbon monoxide directly attacks on the
	A.	Chlorophyll
	В.	Hemoglobin
	C.	Haaecyanin
	D.	Haemerithrin
145.		Ozone is of oxygen.
	A.	Allotrope
	B.	Isomer
	C.	Isotope
	D.	Homomer
146.		$O_2$ is converted into $O_3$ by absorption of
		Cosmic rays
		UV light
		Sun light
		IR radiations
147.		Emission ofoxide by high flying supersonic aircrafts responsible for ozone depletion.
		Sulphuric
		Ferric
		Nitric
		Zinc
148.		are used in refrigeration and air conditioning.
		CFC's
		CH4
		Ozone
		Hydrocarbons
149.		DNA breakage, inhibition, alteration, replication and formation of DNA adduct arises due to,
		Acid rain
		Green house effect
		Global warming
		Ozone layer depletion
150.		Fine ash from pulverized fuel (coal) burned in power station is known as
		Coal ash
		Fly ash
		Pulverized ash
	D.	Fuel ash

151.		London type of smog which contains SO <sub>2</sub> , SO <sub>3</sub> and humidity is responsible for
	A.	Pollution
	В.	Acid rain
(	C.	Global warming
	D.	Ozone layer depletion
152.		is slow decaying radio nuclide.
	A.	Iodine 137
	В.	Neptunium 93
(	C.	Radium 88
	D.	Technetium 43
153.		Human ear can tolerate decibel noise.
	A.	100
	В.	140
	C.	120
	D.	15
		ELECTROCHEMISTRY
154.		$P^{OH}$ of pure water at $25^0$ C is
	A.	7
	В.	
•	C.	14
	D.	None of above
155.		Relationship between equivalent and molar conductance is,
	A.	$(\Lambda_{\rm m} / \Lambda_{\rm eq}) = (Normality / 1)$
	В.	$(\Lambda_{\rm m} / \Lambda_{\rm eq}) = (1 / Molarity)$
•	C.	$(\Lambda_{\rm m} / \Lambda_{\rm eq}) = (Molarity / Normality)$
	D.	$(\Lambda_{\rm m} / \Lambda_{\rm eq}) = (Normality / Molarity)$
156.		On dilution conductivity decreases.
	A.	Specific
	В.	Equivalent
•	C.	Molar
	D.	Molal
157.		Electrochemical cell is a device which is used to get energy.
	A.	Thermal
		Electrical
		Chemical
	D.	Mechanical
158.		Battery is a type of cell.
		Electrolytic
		Electrochemical
		Chemo electric
	D.	Thermal

159.	Dry cell is an example ofcell.
	A. Primary
	B. Secondary
	C. Electrochemical
	D. Electrolytic
160.	is acts as anode in dry / laclanche cell.
	A. Carbon rod
	B. Mn
	C. Zn
	D. Fe
161.	On anode reaction occurs.
	A. Oxidation
	B. Reduction
	C. Redox
1 - 0	D. None of above
162.	<u> </u>
	A. 0.8 to 1.0 V
	B. 1.0 to 1.8 V
	C. 0.5 to 1.0 V
162	D. 0.5 to 1.8 V
103.	Specific conductance is denoted by
	Α. κ
	Β. ρ
	C. σ
1.64	D. Ω
164.	Unit of cell constant is
	Α. Ω
	B. $\Omega^{-1}$
	C. mho
1.0	D. cm <sup>-1</sup>
165.	
	A. $\Omega^{-1}$ cm <sup>-1</sup>
	<b>B. S cm<sup>-1</sup></b> C. Ω cm <sup>-1</sup>
	D. $\Omega^{-1}$ cm
166.	
	A. $\Lambda_{eq} = (1000 \text{ K/N})$
	B. $\Lambda_{eq} = (1000 \text{ G/N})$
	C. $\Lambda_{\rm eq} = (1000  \kappa  \sigma / N)$
	D. $\Lambda_{\rm eq} = (1000\kappa / 2N)$

167. Unit of equivalent conductance is
$A. \Omega^{-1} cm^2 eq^{-1}$
B. $\Omega$ cm <sup>2</sup> eq <sup>-1</sup>
$C. \Omega cm^2 eq$
D. $\Omega^{-1}$ cm <sup>2</sup> eq <sup>-2</sup>
168. Unit of molar conductance is
A. $\Omega^{-1}$ cm <sup>2</sup> mol <sup>-1</sup>
B. $\Omega$ cm <sup>2</sup> mol <sup>-1</sup>
C. $\Omega^{-1} \text{ cm}^{-1} \text{ mol}^{-2}$
D. $\Omega^{-1}  \mathrm{cm}^{-2}  \mathrm{mol}^{-1}$
169. Fuel cells are cells.
A. Primary
B. Secondary
C. Galvanic
D. none of the above
170. Aq solution is used in Ni-metal hydride battery.
A. NaOH
В. КОН
C. NH <sub>4</sub> OH
D. $Ca(OH)_2$
171. Cell potential of Ni - Metal hydride battery is,
A. 1.25 to 1.35 V
B. 0.25 to 1.3 V
C. 0.25 to 0.35 V
D. 0.025 to 1.35 V
172. Hydrogen in metal-hydride is used or acts as anode in battery
A. Lithium Battery
B. Lithium MnO <sub>2</sub> cell
C. Lead Accumulator
D. Ni - metal Hydride battery
173. From conventional method, in steps chemical energy is converted into electrical energy
A. 3
B. 2
C. 1
D. 0
174. In fuel cell net energy profit is %.
A. 20
B. 30
C. 40
D. 45

175. Theoretically any cell shows% efficiency.
A. 75
B. 50
C. 90
D. 100
176. Aqueous solution is used as electrolytic solution in H <sub>2</sub> -O <sub>2</sub> fuel cell.
A. NaOH
В. КОН
C. NH <sub>4</sub> OH
D. Mg (OH) $_2$
177. Practically emf of fuel cell is V.
A. 0.8 to 1.0
B. 0.8 to 0.1
C. 0.5 to 1.0
D. 0.7 to 1.7
178. In methanol - O <sub>2</sub> fuel cell cathode is Ni- sheet with as catalyst.
A. Pt
B. Pd
C. Ag
D. Au
In methanol - $O_2$ fuel cell anode is Ni- sheet with as catalyst.
A. Pt
B. Pd
C. Ag
D. Au
180. Efficiency of methanol - O <sub>2</sub> fuel cell is%.
A. 50 - 80
B. 100
C. 20
D. None of the above
181. In $H_2$ - $O_2$ fuel cell $O_2$ gas acts as
A. Anode
B. Cathode
C. Reductant
D. Oxidant
182. In Li- MnO <sub>2</sub> cell acts as separator.
A. Polyethene
B. Polypropylene
C. Polybutylene
D. Polymeric Li

183. In Li- MnO <sub>2</sub> cell used as electrolytic solution
A. LiO
B. LiX in organic solvent
C. LiX in inorganic solvent
D. LiO in organic solvent
184. In Lithium batteries, electrolytes can't acts as aqueous solution because,
A. High reactivity of Li with H <sub>2</sub> O
B. Low density of Li w. r. t. H <sub>2</sub> O
C. High electro positivity of Li <sup>+</sup>
D. None of above
185. A fresh dry cell has potentialV
A. 1.0
B. 1.5
C. 2.0
D. 3.5
186. In dry cell in secondary reactions the complex formed is
A. $Zn[NH_3]_2Cl_2$
B. $\mathbf{Zn[NH_3]Cl_2}$
C. ZnCl <sub>2</sub>
D. $\mathbf{Zn[NH_3]_2Cl}$
187. Conductivity of 1cm <sup>3</sup> of solution is
A. Equivalent
B. Moral
C. Molal
D. Specific
188. $\Lambda_0 = \lambda_0^+ + \lambda_0^-$ , is the mathematical relation of,
A. Ohm's Law
B. Debye Huckel Limiting Law
C. Ostwald's Law
D. Kohlrausch Law
189. The battery used in automobiles is
A. Lead accumulator
B. Ni-metal hydride
C. Li-MnO <sub>2</sub>
D. Laclanche cell
190. Cell constant can be defined as
A. (Specific / Observed) Conductance
B. (Observed / Specific) Conductance
C. (1 / Specific) Conductance
D. (1 / Observed) Conductance

<u>_</u> .

199.	Aquatic animals survive at dissolved oxygen more than mg / it.
<b>A.</b> 1	5
B. 2	2.5
<b>C.</b> 3	3
D. (	0.25
200. A	pplication of Kohlrausch law,
A. I	Determination of equivalent conductance of weak electrolyte
B. I	Determination of degree of dissociation
C. I	Determination of ionic product of water
<b>D.</b> A	All of above.