

SECOND YEAR HIGHER SECONDARY EXAMINATION MARCH 2019

SUBJECT : MATHEMATICS (COMMERCE)

CODE. NO: SY 51

Qn No	Sub Qns	Answer Key/Value Points	Score	Total
1.	i)	$f(x_1) = f(x_2) \Rightarrow 3 - 4x_1 = 3 - 4x_2$ $\Rightarrow -4x_1 = -4x_2$ $\Rightarrow x_1 = x_2$ <p>$\therefore f$ is one-one.</p> <p>Let $y \in \mathbb{R}$, $y = f(x)$</p> $\Rightarrow y = 3 - 4x$ $\Rightarrow x = \frac{3-y}{4} \in \mathbb{R}$ <p>$\therefore f$ is onto.</p>	$\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$	
	ii)	$y = 3 - 4x$ $\Rightarrow x = \frac{3-y}{4}$ <p>$\therefore f^{-1}(x) = \frac{3-x}{4}$, or $f^{-1} = \frac{3-y}{4}$.</p> <p><u>Remark.</u> for definition of 1-1 and onto give $\frac{1}{2}$ score each. For any alternate correct method give full score.</p>	$\frac{1}{2}$ $\frac{1}{2}$	3
2.		$[a_{ij}] = \begin{bmatrix} a_{11} & a_{12} & a_{13} & a_{14} \\ a_{21} & a_{22} & a_{23} & a_{24} \\ a_{31} & a_{32} & a_{33} & a_{34} \end{bmatrix}$ $= \begin{bmatrix} 1 & 0 & -1 & -2 \\ 3 & 2 & 1 & 0 \\ 5 & 4 & 3 & 2 \end{bmatrix}$ <p>Remark: For any 9 correct entries give full score.</p>	2 1	3

Qn No	Sub Qns	Answer Key/Value Points	Score	Total
3.		$\text{Area} = \frac{1}{2} \begin{vmatrix} x_1 & y_1 & 1 \\ x_2 & y_2 & 1 \\ x_3 & y_3 & 1 \end{vmatrix}$ $= \frac{1}{2} \begin{vmatrix} 0 & 3 & 1 \\ 2 & 0 & 1 \\ 4 & 5 & 1 \end{vmatrix}$ $= \frac{1}{2} 0 - 3(-2) + 1(10) $ $= 8 \text{ sq. units}$	1 1 $\frac{1}{2}$ $\frac{1}{2}$	3
4.	(i) (ii)	$f(2) = 3.$ Since f is continuous at $x=2$ $\lim_{x \rightarrow 2^-} f(x) = \lim_{x \rightarrow 2^+} f(x) = f(2)$ $4k = 3$ $k = \frac{3}{4}$ <p><u>Remark:</u> For definition of continuity give 1 score.</p>	1 1 $\frac{1}{2}$ $\frac{1}{2}$	3
5	(i) (ii)	$\frac{1}{a} \tan^{-1}\left(\frac{x}{a}\right) + C.$ $\int \frac{dx}{x^2 - 6x - 7} = \int \frac{dx}{(x-3)^2 - 4^2}$ $\int \frac{dx}{x^2 - a^2} = \frac{1}{2a} \log \left \frac{x-a}{x+a} \right + C$ $= \frac{1}{8} \log \left \frac{x-7}{x+1} \right + C$	1 $\frac{1}{2}$ 1 $\frac{1}{2}$	3.


Qn No	Sub Qns	Answer Key/Value Points	Score	Total
6	(i) (ii)	$\vec{a} \cdot \vec{b} = 0$ $\text{Let } \vec{a} = \hat{i} - \hat{j} + 3\hat{k}, \vec{b} = 3\hat{i} - \hat{j} + \hat{k}$ $\vec{a} \cdot \vec{b} = 3 + 1 + 3 = 7$ $ \vec{a} = \sqrt{1+1+9} = \sqrt{11}$ $ \vec{b} = \sqrt{9+1+1} = \sqrt{11}$ $\cos \theta = \frac{\vec{a} \cdot \vec{b}}{(\vec{a} \vec{b})} = \frac{7}{\sqrt{11} \cdot \sqrt{11}}$ $= \frac{7}{11}$ <p><u>Remark:</u> For $\cos \theta = \frac{\vec{a} \cdot \vec{b}}{(\vec{a} \vec{b})}$ give $\frac{1}{2}$ score</p>	1 $\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$	3
7.		$\vec{a}_1 = \hat{i} + \hat{j} + \hat{k}, \vec{b}_1 = \hat{i} + \hat{j}$ $\vec{a}_2 = 2\hat{i} + \hat{j} + 2\hat{k}, \vec{b}_2 = \hat{j} + \hat{k}$ $\vec{a}_2 - \vec{a}_1 = \hat{i} + \hat{k}$ $\vec{b}_1 \times \vec{b}_2 = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 1 & 1 & 0 \\ 0 & 1 & 1 \end{vmatrix} = \hat{i} - \hat{j} + \hat{k}$ $(\vec{a}_2 - \vec{a}_1) \cdot (\vec{b}_1 \times \vec{b}_2) = 1 + 1 = 2$ $ \vec{b}_1 \times \vec{b}_2 = \sqrt{3}$ $S.D = \frac{ (\vec{a}_2 - \vec{a}_1) \cdot (\vec{b}_1 \times \vec{b}_2) }{ \vec{b}_1 \times \vec{b}_2 } = \frac{2}{\sqrt{3}}$ <p><u>Remark:</u> For formula give 1 score</p>	1 $\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$	3

Qn No	Sub Qns	Answer Key/Value Points	Score	Total
8	(i) (ii) (iii)	$2 * 3 = 2 \times 3^2 = 18$ $a * b = ab^2, b * a = ba^2$ $\Rightarrow a * b \neq b * a$ $\Rightarrow *$ is not commutative $a * (b * c) = a * (bc^2)$ $= a(bc^2)^2 = ab^2c^4$ $(a * b) * c = (ab^2) * c$ $= ab^2c^2$ $a * (b * c) \neq (a * b) * c$ $\Rightarrow *$ is not associative <u>Remark</u> (i) For direct answer or example give full score (ii) For direct answer give 1 score, with example give full score.	1 $\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$	4
9	i ii)	(i) $\pi/6$ (ii) Let $\sin^{-1}(\frac{5}{13}) = x \Rightarrow \sin x = \frac{5}{13}$ $\Rightarrow \tan x = \frac{5}{12}$ $\therefore x = \tan^{-1} \frac{5}{12}$ (iii) $\sin^{-1} \frac{3}{5} = \tan^{-1} \frac{3}{4}$	1 1 1	

(5)

Qn No	Sub Qns	Answer Key/Value Points	Score	Total
		$\therefore \tan^{-1} \frac{5}{12} + \tan^{-1} \frac{3}{4} = \tan^{-1} \left(\frac{\frac{5}{12} + \frac{3}{4}}{1 - \frac{5}{12} \cdot \frac{3}{4}} \right)$ $= \tan^{-1} \frac{56}{33}$ <p><u>Remark</u>: For any correct alternate method give full score. For any suitable formula give 1 score.</p>	1	4
10	<p>i) 0</p> <p>ii) $R_1 \rightarrow R_1 + R_2 + R_3$</p>	$\Rightarrow \begin{vmatrix} 3x+k & 3x+k & 3x+k \\ x & x+k & x \\ x & x & x+k \end{vmatrix}$ $\Rightarrow (3x+k) \begin{vmatrix} 1 & 1 & 1 \\ x & x+k & x \\ x & x & x+k \end{vmatrix}$ <p>$C_2 \rightarrow C_2 - C_1$ and $C_3 \rightarrow C_3 - C_1$</p> $\Rightarrow (3x+k) \begin{vmatrix} 1 & 0 & 0 \\ x & k & 0 \\ x & 0 & k \end{vmatrix}$ $\Rightarrow (3x+k) \cdot 1 \cdot (k^2 - 0) = k^2 (3x+k)$ <p><u>Remark</u> For correct expansion give one score.</p>	<p>1</p> <p>1</p> <p>$\frac{1}{2}$</p> <p>1</p> <p>$\frac{1}{2}$</p>	<p>4</p>

Qn No	Sub Qns	Answer Key/Value Points	Score	Total
11	(i) (ii)	$f'(x) = 2x - 4$ $f(x)$ is continuous in $[1, 3]$ $f'(x) = 2x - 4$ exist in $(1, 3)$ $f(1) = 0$ and $f(3) = 0$ $\Rightarrow f(1) = f(3)$ Conditions satisfied $\therefore f'(c) = 0$ $\Rightarrow 2c - 4 = 0 \Rightarrow c = 2 \in (1, 3)$ Hence Rolle's Theorem Verified	1 $\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$ 1 $\frac{1}{2}$	4
12	(i) (ii)	(b) 0 Let $I = \int_0^{\pi/2} \frac{\cos^n x}{\sin^n x + \cos^n x} dx \rightarrow (1)$ we have $\int_0^a f(x) dx = \int_0^a f(a-x) dx$ $\therefore I = \int_0^{\pi/2} \frac{\cos^n(\pi/2 - x)}{\sin^n(\pi/2 - x) + \cos^n(\pi/2 - x)} dx$ $= \int_0^{\pi/2} \frac{\sin^n x}{\cos^n x + \sin^n x} dx \rightarrow (2)$ $\textcircled{1} + \textcircled{2}$ $\Rightarrow 2I = \int_0^{\pi/2} 1 dx = (x)_0^{\pi/2} = \frac{\pi}{2}$ $\therefore I = \frac{\pi}{4}$	1 1 $\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$	4.

Qn No	Sub Qns	Answer Key/Value Points	Score	Total
13.		<p>Area under a curve = $\int_a^b y \, dx$</p>  <p>Area of the shaded portion = $\int_0^2 \sqrt{4-x^2} \, dx$</p> $\int \sqrt{a^2-x^2} \, dx = \frac{x}{2} \sqrt{a^2-x^2} + \frac{a^2}{2} \sin^{-1} \frac{x}{a} + C$ $\therefore \text{Area} = \left[\frac{x}{2} \sqrt{4-x^2} + \frac{4}{2} \cdot \sin^{-1} \frac{x}{2} \right]_0^2$ $= (0 + 2 \sin^{-1}(1)) - (0 + 0)$ $= 2 \cdot \frac{\pi}{2} = \pi \text{ sq. units}$ <p>\therefore Required Area = 4π sq. units</p> <p><u>Remark:</u> For correct figure give $\frac{1}{2}$ score.</p>	<p>$\frac{1}{2}$</p> <p>1</p> <p>$\frac{1}{2}$</p> <p>$\frac{1}{2}$</p> <p>$\frac{1}{2}$</p> <p>$\frac{1}{2}$</p>	<p>4.</p>
14	(i)	Order - 1 degree - 1	$\frac{1}{2} + \frac{1}{2}$	
	(ii)	<p>$P = \frac{1}{x}, Q = x^2$</p> <p>Integrating factor = $e^{\int P \, dx} = e^{\int \frac{1}{x} \, dx} = e^{\log x} = x$</p>	$\frac{1}{2}$	
			$\frac{1}{2}$	

Qn No	Sub Qns	Answer Key/Value Points	Score	Total
16	(c)	$P(A \cap B) = P(A) + P(B) - P(A \cup B)$ $= 0.6 + 0.5 - 0.8 = 0.3$ $P(A/B) = \frac{P(A \cap B)}{P(B)}$ $= \frac{0.3}{0.5} = \frac{3}{5} = 0.6$	1 1/2 1 1/2	4
	(c) (d)	$P(E \cap F)$	1	
17		<p>Let food M be x kg and food N be y kg.</p> <p>Minimise $Z = 50x + 70y$</p> <p>Subject to $3x + 4y \geq 9$ $5x + 2y \geq 11$ $x, y \geq 0$</p>	1 1 1 1	4
18	(c)	$A^T = \begin{bmatrix} 2 & 2 & 1 \\ 0 & 1 & -1 \\ 1 & 3 & 0 \end{bmatrix}$	1	
	(c)	<p>Let $P = \frac{A + A^T}{2}$</p> $= \frac{1}{2} \begin{bmatrix} 4 & 2 & 2 \\ 2 & 2 & 2 \\ 2 & 2 & 0 \end{bmatrix} = \begin{bmatrix} 2 & 1 & 1 \\ 1 & 1 & 1 \\ 1 & 1 & 0 \end{bmatrix}$	1/2 1/2	
		$P^T = P \Rightarrow P$ is symmetric.		










Qn No	Sub Qns	Answer Key/Value Points	Score	Total
		$Q = \frac{A - A^T}{2}$ $= \frac{1}{2} \begin{bmatrix} 0 & -2 & 0 \\ 2 & 0 & 4 \\ 0 & -4 & 0 \end{bmatrix} = \begin{bmatrix} 0 & -1 & 0 \\ 1 & 0 & 2 \\ 0 & -2 & 0 \end{bmatrix}$ $Q^T = -Q \Rightarrow Q \text{ is skew-symmetric.}$ <p>Also $P + Q = A$</p>	$\frac{1}{2}$ $\frac{1}{2}$ 1	
(cc)		$A \cdot A^T = \begin{bmatrix} 2 & 0 & 1 \\ 2 & 1 & 3 \\ 1 & -1 & 0 \end{bmatrix} \begin{bmatrix} 2 & 2 & 1 \\ 0 & 1 & -1 \\ 1 & 3 & 0 \end{bmatrix}$ $= \begin{bmatrix} 5 & 7 & 2 \\ 7 & 14 & 1 \\ 2 & 1 & 2 \end{bmatrix}$	1 1	6.
19 (c)		$ A = 3(2-3) + 2(4+4) + 3(-10)$ $= -3 + 16 - 30 = -17$	1	
(c)		$\text{Cofactor matrix} = \begin{bmatrix} -1 & -8 & -10 \\ -5 & -6 & 1 \\ -1 & 9 & 7 \end{bmatrix}$	2	
		$\text{adj } A = \begin{bmatrix} -1 & -5 & -1 \\ -8 & -6 & 9 \\ -10 & 1 & 7 \end{bmatrix}$	1	

Qn No	Sub Qns	Answer Key/Value Points	Score	Total
	(cc)	$Ax = B, A = \begin{bmatrix} 3 & -2 & 3 \\ 2 & 1 & -1 \\ 4 & -3 & 2 \end{bmatrix}$ $x = \begin{bmatrix} x \\ y \\ z \end{bmatrix} \quad B = \begin{bmatrix} 2 \\ 3 \\ 0 \end{bmatrix}$ $x = A^{-1}B$ $= \frac{1}{-17} \begin{bmatrix} -1 & -5 & -1 \\ -8 & -6 & 9 \\ -10 & 17 \end{bmatrix} \begin{bmatrix} 2 \\ 3 \\ 0 \end{bmatrix} = \begin{bmatrix} 1 \\ 2 \\ 1 \end{bmatrix}$ <p>$\therefore x=y \quad x=1, y=2, z=1$</p>	$\frac{1}{2}$ $\frac{1}{2}$ 1	6.
20	(c)	$x^y = y^x$ <p>Take log on both sides</p> $\log x^y = \log y^x$ $y \log x = x \log y$ <p>differentiating</p> $y \cdot \frac{1}{x} + \log x \cdot \frac{dy}{dx} = x \cdot \frac{1}{y} \frac{dy}{dx} + \log y \cdot 1$ $\frac{dy}{dx} (\log x - \frac{x}{y}) = \log y - \frac{y}{x}$ $\therefore \frac{dy}{dx} = \frac{\log y - y/x}{\log x - x/y}$	1 1 1	

Qn No	Sub Qns	Answer Key/Value Points	Score	Total
	(cc)	$\frac{dx}{dt} = 4at, \quad \frac{dy}{dt} = 4at^3$ $\frac{dy}{dx} = \frac{dy/dt}{dx/dt} = \frac{4at^3}{4at}$ $= \underline{\underline{t^2}}$	1+1 $\frac{1}{2} + \frac{1}{2}$	6
21	(c)	$y^2 = x$ $2y \frac{dy}{dx} = 1$ $\frac{dy}{dx} = \frac{1}{2y}$ <p>Slope, $m = \frac{dy}{dx}$ at (1,1)</p> $= \frac{1}{2}$ <p>Equation of tgt. is</p> $y - y_1 = m(x - x_1)$ $\Rightarrow y - 1 = \frac{1}{2}(x - 1)$ $\Rightarrow x - 2y + 1 = 0$ <hr/> $f'(x) = 6x^2 - 6x - 36$ $f'(x) = 0 \Rightarrow 6(x^2 - x - 6) = 0$ $\Rightarrow (x - 3)(x + 2) = 0$ $\Rightarrow x = -2, 3$ <p>for increasing $f'(x) > 0$, decreasing $f'(x) < 0$ increasing in the interval $(-\infty, -2) \cup (3, \infty)$</p> <p>and decreasing in $(-2, 3)$</p>	$\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$ 1 $\frac{1}{2}$ 1 $\frac{1}{2}$ $\frac{1}{2}$	6

Qn No	Sub Qns	Answer Key/Value Points	Score	Total
22.	(i)	<p>Since vectors are coplanar</p> $[\vec{a} \ \vec{b} \ \vec{c}] = 0$ $\therefore \begin{vmatrix} 2 & -1 & 1 \\ 3 & 1 & 2 \\ 1 & \lambda & -3 \end{vmatrix} = 0$ $\Rightarrow 2(-3-2\lambda) + 1(-9-2) + 1(3\lambda-1) = 0$ $\Rightarrow \lambda = -18$	<p>1</p> <p>1</p> <p>1/2</p> <p>1/2</p>	
	(ii)	$[\vec{a}+\vec{b} \ \vec{b}+\vec{c} \ \vec{c}+\vec{a}]$ $= (\vec{a}+\vec{b}) \cdot [(\vec{b}+\vec{c}) \times (\vec{c}+\vec{a})]$ $= (\vec{a}+\vec{b}) \cdot \left[\vec{b} \times \vec{c} + \vec{b} \times \vec{a} + \vec{c} \times \vec{c} + \vec{c} \times \vec{a} \right]$ $\equiv (\vec{a}+\vec{b}) \cdot \left[\vec{b} \times \vec{c} + \vec{b} \times \vec{a} + \vec{c} \times \vec{a} \right]$ $= \vec{a} \cdot (\vec{b} \times \vec{c}) + \vec{a} \cdot (\vec{b} \times \vec{a}) + \vec{a} \cdot (\vec{c} \times \vec{a}) + \vec{b} \cdot (\vec{b} \times \vec{c}) + \vec{b} \cdot (\vec{b} \times \vec{a}) + \vec{b} \cdot (\vec{c} \times \vec{a})$ $= [\vec{a} \ \vec{b} \ \vec{c}] + 0 + 0 + 0 + 0 + [\vec{b} \ \vec{c} \ \vec{a}]$ $= 2 [\vec{a} \ \vec{b} \ \vec{c}]$	<p>1</p> <p>1/2</p> <p>1/2</p> <p>1/2</p> <p>1/2</p> <p>1/2</p>	<p>6</p>

Qn No	Sub Qns	Answer Key/Value Points	Score	Total									
23	(c)	$2x + y = 6.$ <table border="1" data-bbox="750 257 1029 392"> <tr><td>x</td><td>0</td><td>3</td></tr> <tr><td>y</td><td>6</td><td>0</td></tr> </table>	x	0	3	y	6	0	$\frac{1}{2}$				
x	0	3											
y	6	0											
		$3x + 4y = 12$ <table border="1" data-bbox="734 425 1013 548"> <tr><td>x</td><td>0</td><td>4</td></tr> <tr><td>y</td><td>3</td><td>0</td></tr> </table>	x	0	4	y	3	0	$\frac{1}{2}$				
x	0	4											
y	3	0											
			3.										
	ii)	Corner points are $A(3, 0)$ $B(4, 0)$ $C(\frac{12}{5}, \frac{6}{5})$	$\frac{1}{2}$	6									
		<table border="0"> <tr> <td>A</td> <td>(3, 0)</td> <td>$z = 30$</td> </tr> <tr> <td>B</td> <td>(4, 0)</td> <td>$z = 40$ → Maximum.</td> </tr> <tr> <td>C</td> <td>$(\frac{12}{5}, \frac{6}{5})$</td> <td>$z = 28.8 = \frac{144}{5}$</td> </tr> </table> <p>Maximum at $x = 4, y = 0.$</p>	A	(3, 0)	$z = 30$	B	(4, 0)	$z = 40$ → Maximum.	C	$(\frac{12}{5}, \frac{6}{5})$	$z = 28.8 = \frac{144}{5}$	$\frac{1}{2}$	
A	(3, 0)	$z = 30$											
B	(4, 0)	$z = 40$ → Maximum.											
C	$(\frac{12}{5}, \frac{6}{5})$	$z = 28.8 = \frac{144}{5}$											
Remark:		For each correct line give 1 score. {Incorrect shaded region } give $3\frac{1}{2}$ score. and correct graph											

1. Preeti. K.R, 9495331511
G.H.S.S. Wadakkanchery, (8021) 
2. Ganga Devi. K. 9497853974
SVVHSS Miyapadavu. (14095) 
3. JyothiKunshna.S 9446433842
H.S.S Thiruvampady. Atappuzha (04068) 
4. Leena P.V. 9495216599
VCSHSS, Puthenvelikkasa (07041) 
5. Jincy P. Geia 9847777424
St. Dominic's HSS Kanjirapally (05062) 
6. HAREESH S 9447451513
St. Mary's HSS Pariyapuram (11045) 
7. J. John Victor 9446171748
New HSS Nellikood TVM (01069) 
8. MARIA DANIEL FERNANDEZ
KRISPAS HSS ; KOLLAM. 
9. Raheeq. K 9745108689. 
Chennamangallur HSS. Kozhikode. (10044).
10. Jose Mathew , 9495214997
st. Mary's H.S.S , Vellavankunnu
Idukki. 