

# Study Material

Downloaded from Vedantu

## About Vedantu

Vedantu is India's largest **LIVE online teaching platform** with best teachers from across the country.

Vedantu offers Live Interactive Classes for **JEE, NEET, KVPY, NTSE, Olympiads, CBSE, ICSE, IGCSE, IB & State Boards** for Students Studying in **6-12th Grades** and Droppers.

**FREE** LIVE ONLINE

### MASTER CLASSES

FREE Webinars by Expert Teachers



Register for **FREE**

## Awesome Master Teachers



**Anand Prakash**  
B.Tech, IIT Roorkee  
Co-Founder, Vedantu



**Pulkit Jain**  
B.Tech, IIT Roorkee  
Co-Founder, Vedantu



**Vamsi Krishna**  
B.Tech, IIT Bombay  
Co-Founder, Vedantu



“My mentor is approachable and **guides me in my future aspirations as well.**”

Student - **Ayushi**



“My son loves the sessions and **I can already see the change.**”

Parent - **Sreelatha**



**10,04,600+**  
Hours of LIVE Learning



**9,49,900+**  
Happy Students



**95%**  
Top Results

**95%** Students of Regular Tuitions on Vedantu scored above **90%** in exams!

Vedantu

## FREE MASTER CLASS SERIES

- ✓ For **Grades 6-12th** targeting **JEE, CBSE, ICSE** & much more
- ✓ **Free 60 Minutes Live Interactive** classes everyday
- ✓ Learn from the **Master Teachers** - India's best

Register for **FREE**

Limited Seats!

# Download Vedantu's App & Get



All Study Material  
with Solution




LIVE  
Doubt Solving



Daily  
LIVE Classes



FREE Tests and  
Reports



GET IT ON  
**Google Play**

**DOWNLOAD THE APP**

# DIFFERENTIAL EQUATION

## 1. INTRODUCTION

Differential equation constitute a very important part of mathematics as it has many applications in real life. Various laws of physics are often in the form of equations involving rate of change of one quantity with respect to another. As the mathematical equivalent of a rate is a derivative, differential equation arise very naturally in real life and methods for solving them acquire paramount importance.

### Definition

An equation involving the dependent variable and independent variable and also the derivatives of the dependable variable is known as differential equation.

For example:

$$(i) \frac{dy}{dx} = \frac{x}{y^{1/3}(1+x^{1/3})} \quad (ii) \frac{d^2y}{dx^2} = -p^2y$$

$$(iii) \left[1 + \left(\frac{dy}{dx}\right)^2\right]^{3/2} = 3 \frac{d^2y}{dx^2} \quad (iv) x^2 \left(\frac{dy}{dx}\right)^2 = y^2 + 1$$

Differential equations which involve only one independent variable are called ordinary differential equation.

## 2. ORDER AND DEGREE OF DIFFERENTIAL EQUATIONS

### 2.1 Order

The order of a differential equation is the order of the highest derivative involved in the differential equation

For example:

$$(i) \left(\frac{dy}{dx}\right)^3 + \left(\frac{dy}{dx}\right)^2 + 4x = 0 \text{ is the differential equation of the first order because maximum derivative of } y \text{ with respect to } x \text{ is } \frac{dy}{dx}$$

$$(ii) \frac{d^2y}{dx^2} = -p^2y \text{ is the differential equation of the second order because maximum derivative of } y \text{ w.r.t } x \text{ is } \frac{d^2y}{dx^2}$$

$$(iii) \left(\frac{d^3y}{dx^3}\right)^2 - 3\left(\frac{dy}{dx}\right)^3 + 2 = 0 \text{ is the differential equation of the third order because maximum derivative of } y \text{ w.r.t } x \text{ is } \frac{d^3y}{dx^3}$$

### 2.2 Degree

The degree of a differential equation is the degree of the highest differential coefficient when the equation has been made rational and integral as far as the differential coefficients are concerned.

For example:

$$(i) \frac{dy}{dx} = \frac{x}{y^{1/3}(1+x^{1/3})} \text{ is the differential equation of first degree, because power of the highest order derivative } \frac{dy}{dx} \text{ is } 1.$$

$$(ii) \left(\frac{d^3y}{dx^3}\right)^2 - 3\left(\frac{dy}{dx}\right)^3 + 2 = 0 \text{ is the differential equation of second degree, because power of highest order derivative } \frac{d^3y}{dx^3} \text{ is } 2.$$

$$(iii) \left[1 + \left(\frac{dy}{dx}\right)^2\right]^{2/3} = 3 \frac{d^2y}{dx^2} \text{ is the differential equation of third degree, because power of highest order derivative } \frac{d^2y}{dx^2} \text{ is } 3 \text{ (after cubing)}$$

# Vedantu

## Study Materials

[NCERT Solutions for Class 6 to 12 \(Math & Science\)](#)

[Revision Notes for Class 6 to 12 \(Math & Science\)](#)

[RD Sharma Solutions for Class 6 to 12 Mathematics](#)

[RS Aggarwal Solutions for Class 6, 7 & 10 Mathematics](#)

[Important Questions for Class 6 to 12 \(Math & Science\)](#)

[CBSE Sample Papers for Class 9, 10 & 12 \(Math & Science\)](#)

[Important Formula for Class 6 to 12 Math](#)

[CBSE Syllabus for Class 6 to 12](#)

[Lakhmir Singh Solutions for Class 9 & 10](#)

[Previous Year Question Paper](#)

[CBSE Class 12 Previous Year Question Paper](#)

[CBSE Class 10 Previous Year Question Paper](#)

[JEE Main & Advanced Question Paper](#)

[NEET Previous Year Question Paper](#)

## DIFFERENTIAL EQUATION

**Illustration 1:** Find the order and degree of the following differential equations.

$$(i) \sqrt{\frac{d^2y}{dx^2}} = \sqrt[3]{\frac{dy}{dx}} + 3$$

$$(ii) \frac{d^2y}{dx^2} = \left\{ 1 + \left( \frac{dy}{dx} \right)^4 \right\}^{5/3}$$

$$(iii) y = px + \sqrt{a^2p^2 + b^2} \text{ where } p = \frac{dy}{dx}$$

**Sol.** (i) The given differential equation can be written as

$$\left( \frac{d^2y}{dx^2} \right)^3 = \left( \frac{dy}{dx} + 3 \right)^2$$

Hence order = 2, degree = 3

(ii) The given differential equation can be written as

$$\left( \frac{d^2y}{dx^2} \right)^3 = \left[ 1 + \left( \frac{dy}{dx} \right)^4 \right]^{5/3}$$

Hence order = 2, degree = 3

(iii) The given differential equation can be written as

$$\left( y - x \frac{dy}{dx} \right)^2 = a^2 \left( \frac{dy}{dx} \right)^2 + b^2$$

Hence order = 1, degree = 2

### 3. FORMATION OF ORDINARY DIFFERENTIAL EQUATION

An ordinary differential equation is formed in an attempt to eliminate certain arbitrary constants from a relation in the variables and constants. Consider an equation containing  $n$  arbitrary constants. Differentiating this equation  $n$  times we get  $n$  additional equations containing  $n$  arbitrary constants and derivatives. Eliminating  $n$  arbitrary constants from the above  $(n + 1)$  equations, we obtain differential equation involving  $n$ th derivative.

Thus if an equation contains  $n$  arbitrary constants, the resulting differential equation obtained by eliminating these constants will be a differential equation of  $n$ th order. i.e., an equation of the form

$$\phi \left( x, y, \frac{dy}{dx}, \frac{d^2y}{dx^2}, \dots, \frac{d^ny}{dx^n} \right) = 0$$

**Illustration 2 :** Find the differential equation of the family of all circles which pass through the origin and whose centre lie on  $y$ -axis

**Sol.** Let the equation of the circle be

$$x^2 + y^2 + 2gx + 2fy + c = 0$$

If it passes through  $(0, 0)$ , then  $c = 0$

$\therefore$  The equation of circle is  $x^2 + y^2 + 2gx + 2fy = 0$

Since the centre of the circle lies on  $y$ -axis then  $g = 0$

$\therefore$  The equation of the circle is

$$x^2 + y^2 + 2fy = 0 \quad \dots(i)$$

This represents family of circles.

Differentiating, we get

$$2x + 2y \frac{dy}{dx} + 2f \frac{dy}{dx} = 0 \quad \dots(ii)$$

From (i) and (ii), we get

or,  $(x^2 - y^2) \frac{dy}{dx} - 2xy = 0$  Which is the required differential equation.

### 4. SOLUTION OF A DIFFERENTIAL EQUATION

The solution of the differential equation is a relation is a relation between the independent and dependent variable free from derivatives satisfying the given differential equation.

Thus the solution of  $dy/dx = m$  could be obtained by simply integrating both sides i.e.,  $y = mx + c$ , where  $c$  is arbitrary constant.

(a) **General solution (or complete primitive)**

The general solution of a differential equation is the relation between the variables (not involving the derivatives) which contain the same number of the arbitrary constants as the order of the differential equation.

Thus the general solution of the differential equation

$$\frac{d^2y}{dx^2} = 4y \text{ is } y = A \sin 2x + B \cos 2x, \text{ where } A \text{ and } B \text{ are the constants.}$$

(b) **Particular solution or Integral**

A solution which is obtained by giving particular values to the arbitrary constants in the general solution is called a particular solution.

**Illustration 3:** Show that  $v = \frac{A}{r} + B$  is the general solution of

the second order differential equation  $\frac{d^2v}{dr^2} + \frac{2}{r} \frac{dv}{dr} = 0$ ,

where A and B are arbitrary constant.

**Sol.** Given  $v = \frac{A}{r} + B$

Differentiating twice  $\frac{d^2v}{dr^2} = \frac{2A}{r^3}$  ... (i)

From (i)  $\frac{d^2v}{dr^2} + \frac{2}{r} \frac{dv}{dr} = \frac{2A}{r^3} + \frac{2}{r} \left( -\frac{A}{r^2} \right) = \frac{2A}{r^3} - \frac{2A}{r^3} = 0$

Putting  $A = 4$ ,  $B = 5$  in  $v = \frac{A}{r} + B$  we get a particular solution of the differential equation

$\frac{d^2v}{dr^2} + \frac{2}{r} \frac{dv}{dr} = 0$  is  $v = \frac{4}{r} + 5$ .

**Illustration 4:** Show that  $y = ae^x + be^{2x} + ce^{-3x}$  is a solution of

the equation  $\frac{d^3y}{dx^3} - 7 \frac{dy}{dx} + 6y = 0$

**Sol.** We have

$y = ae^x + be^{2x} + ce^{-3x}$  ... (i)

Differentiating, we get

$y_1 = ae^x + 2be^{2x} - 3ce^{-3x}$  ... (ii)

$\therefore y_1 - y = be^{2x} - 4ce^{-3x}$  ... (iii)

Differentiating (ii), we get

$y_2 = ae^x + 4be^{2x} + 9ce^{-3x}$

$\therefore y_2 - y_1 = 2be^{2x} + 12ce^{-3x}$  ... (iv)

Now, (iv) - 2 (iii)  $\Rightarrow y_2 - y_1 - 2(y_1 - y) = 20ce^{-3x}$

or,  $y_2 - 3y_1 + 2y = 20ce^{-3x}$  ... (v)

Differentiating,  $y_2 = ae^x + 4be^{2x} + 9ce^{-3x}$ , we get

$y_3 = ae^x + 8be^{2x} - 27ce^{-3x}$

Now  $y_3 - 3y_2 + 2y_1 = -60ce^{-3x}$  ... (vi)

And (vi) + 3(v)  $\Rightarrow y_3 - 3y_2 + 2y_1 + 3(y_2 - 3y_1 + 2y) = 0$

or,  $y_3 - 7y_1 + 6y = 0$

i.e.,  $\frac{d^3y}{dx^3} - 7 \frac{dy}{dx} + 6y = 0$ , which is the required differential equation

## 5. METHOD OF SOLVING AN EQUATION OF THE FIRST ORDER AND FIRST DEGREE

A differential equation of the first order and first degree can be written in the form

$$\frac{dy}{dx} = f(x, y)$$

or,  $M dx + N dy = 0$ , where M and N are functions of x and y

### Method - 1

#### (i) Variable Separation:

The general form of such an equation is

$$f(x)dx + f(y)dy = 0 \quad \dots (i)$$

Integrating, we get

$$\int f(x)dx + \int f(y)dy = c \text{ which is the solution of (i)}$$

#### (ii) Solution of differential equation of the type

$$\frac{dy}{dx} = f(ax + by + c) :$$

Consider the differential equation  $\frac{dy}{dx} = f(ax + by + c)$

... (i)

Where  $f(ax + by + c)$  is some function of  $ax + by + c$ .

Let  $z = ax + by + c$

$$\therefore \frac{dz}{dx} = a + b \frac{dy}{dx}$$

$$\text{or, } \frac{dy}{dx} = \frac{\frac{dz}{dx} - a}{b}$$

$$\text{From (i) } \frac{\frac{dz}{dx} - a}{b} = f(z)$$

$$\text{or, } \frac{dz}{dx} = b f(z) + a$$



# DIFFERENTIAL EQUATION

$$\text{or, } \frac{dz}{b f(z) + a} = dx \quad \dots(ii)$$

In the differential equation (ii), the variables  $x$  and  $z$  are separated.

Integrating, we get

$$\int \frac{dx}{b f(z) + a} = \int dx + c$$

$$\text{or, } \int \frac{dx}{b f(z) + a} = x + c, \text{ where } z = ax + by + c$$

This represents the general solution of the differential equation (i)

**Illustration 5.** Solve  $(x - y)^2 \frac{dy}{dx} = a^2$

**Sol.** Putting  $x - y = v$

$$\Rightarrow \frac{dy}{dx} = 1 - \frac{dv}{dx} \Rightarrow dx = \frac{v^2}{v^2 - a^2} dv, \text{ variable have been separated}$$

$$\text{Integrating, we get } \int dx = \int \frac{v^2}{v^2 - a^2} dv$$

$$\text{or, } 2y + k = a \log \frac{x - y - a}{x - y + a}$$

**Illustration 6.** Solve,  $\frac{dy}{dx} = \sin(x + y) + \cos(x + y)$

**Sol.** Let  $z = x + y$

$$\therefore \frac{dz}{dx} = 1 + \frac{dy}{dx} \Rightarrow \frac{dy}{dx} = \frac{dz}{dx} - 1$$

$$\frac{dz}{dx} - 1 = \sin z + \cos z$$

$$\text{or, } dx = \frac{dz}{\sin z + \cos z + 1}$$

Integrating, we get

$$\int dx = \int \frac{dz}{\sin z + \cos z + 1} = \int \frac{dt}{t + 1}, \text{ putting } t = \tan \frac{z}{2}$$

i.e.,  $x + c = \log |t + 1|$  This is the required general solution.

**(iii) Solution of differential equation of the type**

$$\frac{dy}{dx} = \frac{a_1 x + b_1 y + c_1}{a_2 x + b_2 y + c_2}, \text{ where } \frac{a_1}{a_2} = \frac{b_1}{b_2} \neq \frac{c_1}{c_2}$$

$$\text{Here } \frac{dy}{dx} = \frac{a_1 x + b_1 y + c_1}{a_2 x + b_2 y + c_2} \text{ where } \frac{a_1}{a_2} = \frac{b_1}{b_2} \neq \frac{c_1}{c_2} \quad \dots(i)$$

$$\text{Let } \frac{a_1}{a_2} = \frac{b_1}{b_2} = \lambda \text{ (say)}$$

$$\therefore a_1 = \lambda a_2, b_1 = \lambda b_2$$

$$\text{From (i), } \frac{dy}{dx} = \frac{\lambda a_2 x + \lambda b_2 y + c_1}{a_2 x + b_2 y + c_2} = \frac{\lambda(a_2 x + b_2 y) + c_1}{a_2 x + b_2 y + c_2} \quad \dots(ii)$$

$$\text{Let } z = a_2 x + b_2 y$$

$$\therefore \frac{dz}{dx} = a_2 + b_2 \frac{dy}{dx} \Rightarrow \frac{dy}{dx} = \frac{\frac{dz}{dx} - a_2}{b_2} \quad \dots(iii)$$

$$\text{From (ii) and (iii), we get } \frac{\frac{dz}{dx} - a_2}{b_2} = \frac{\lambda z + c_1}{z + c_2}$$

$$\text{or, } \frac{dz}{dx} = \frac{b_2(\lambda z + c_1)}{z + c_2} + a_2 = \frac{\lambda b_2 z + b_2 c_1 + a_2 z + a_2 c_2}{z + c_2}$$

$$\text{or, } dx = \frac{z + c_2}{\lambda b_2 + a_2)z + b_2 c_1 + a_2 c_2} dz, \text{ where } x \text{ and } z \text{ are separated}$$

Integrating, we get

$$x + c = \int \frac{z + c_2}{\lambda b_2 + a_2)z + b_2 c_1 + a_2 c_2} dz \text{ where } z = a_2 x + b_2 y$$

## Method – 2

## (i) Homogeneous differential equation:

A function  $f(x, y)$  is called homogeneous function of degree  $n$  if

$$f(\lambda x, \lambda y) = \lambda^n f(x, y)$$

For example:

- (a)  $f(x, y) = x^2y^2 - xy^3$  is a homogeneous function of degree four, since

$$\begin{aligned} f(\lambda x, \lambda y) &= (\lambda^2 x^2)(\lambda^2 y^2) - (\lambda x)(\lambda^3 y^3) \\ &= \lambda^4 (x^2y^2 - xy^3) \\ &= \lambda^4 f(x, y) \end{aligned}$$

- (b)  $f(x, y) = x^2e^{x/y} + \frac{x^3}{y} + y^2 \log\left(\frac{y}{x}\right)$  is a homogeneous function of degree two, since

$$\begin{aligned} f(\lambda x, \lambda y) &= (\lambda^2 x^2)e^{\lambda x/\lambda y} + \frac{\lambda^3 x^3}{\lambda y} + (\lambda^2 y^2) \log\left(\frac{\lambda y}{\lambda x}\right) \\ &= \lambda^2 \left[ x^2 e^{x/y} + \frac{x^3}{y} + y^2 \log\left(\frac{y}{x}\right) \right] \\ &= \lambda^2 f(x, y) \end{aligned}$$

A differential equation of the form  $\frac{dy}{dx} = f(x, y)$ , where

$f(x, y)$  is a homogeneous polynomial of degree zero is called a homogeneous differential equation. Such

equations are solved by substituting  $v = \frac{y}{x}$  or  $\frac{x}{y}$  and

then separating the variables.

**Illustration 7 :** Solve  $\frac{dy}{dx} = \frac{y(2y-x)}{x(2y+x)}$  ... (i)

**Sol.** Since each of the functions  $y(2y-x)$  and  $x(2y+x)$  is a homogeneous function of degree 2, so the given equation is a homogeneous differential equation.

$\therefore$  Putting  $y = vx$

Differentiating w.r.t  $x$ , we get  $\frac{dy}{dx} = v + x \frac{dv}{dx}$

From (i),

$$v + x \frac{dv}{dx} = \frac{vx(2vx-x)}{x(2vx+x)} = \frac{v(2v-1)}{2v+1}$$

$$\Rightarrow 2dv + \frac{dx}{x} + 2\frac{dx}{x} = 0$$

Integrating, we get

$$2v + \log v + \log x^2 = \log k \quad \text{or,} \quad xy = ke^{-2y/x}$$

## (ii) Differential equation reducible to homogeneous forms:

Equation of the form  $\frac{dy}{dx} = \frac{ax+by+c}{a'x+b'y+c'}$  where  $\frac{a}{a'} \neq \frac{b}{b'}$

can be reduced to homogeneous form by changing the variables  $x, y$  to  $x', y'$  by equations  $x = x' + h$  and  $y = y' + k$  where  $h$  and  $k$  are constants to be chosen so as to make the given equation homogeneous, we have

$$dx = dx' \quad \text{and} \quad dy = dy'$$

$\therefore$  The given equation becomes

$$\begin{aligned} \frac{dy'}{dx'} &= \frac{a(x'+h) + b(y'+k) + c}{a'(x'+h) + b'(y'+k) + c'} \\ &= \frac{ax' + by' + (ah + bk + c)}{a'x' + b'y' + (a'h + b'k + c')} \end{aligned}$$

Now, we choose  $h$  and  $k$  so that

$$ah + bk + c = 0$$

$$\text{and} \quad a'h + b'k + c' = 0$$

From these equation we get the values of  $h$  and  $k$  in terms of the coefficients.

Then the given equation reduces to

$$\frac{dy'}{dx'} = \frac{ax' + by'}{a'x' + b'y'}$$

Which is the homogeneous form.



## DIFFERENTIAL EQUATION

## Method – 3

## (i) Linear differential equation:

A differential equation is said to be linear if the dependent variable  $y$  and its derivative occur in the first degree.

An equation of the form  $\frac{dy}{dx} + P y = Q$  ... (i)

where  $P$  and  $Q$  are functions of  $x$  only or constant is called a linear equation of the first order

Similarly  $\frac{dx}{dy} + P x = Q$  is a linear differential equation

where  $P$  and  $Q$  are functions of  $y$  only. To get the general solution of the above equations we shall determine a function  $R$  of  $x$  called Integrating function (I.F). We shall multiply both sides of the given equation by  $R$

$\therefore$  where,  $R = e^{\int P dx} = \text{I.F}$  ... (iii)

From (i) and (iii), we get

$$e^{\int P dx} \cdot \frac{dy}{dx} + P y e^{\int P dx} = Q e^{\int P dx} \quad \text{or,}$$

$$\frac{d}{dx} \left( y e^{\int P dx} \right) = Q e^{\int P dx}$$

Integrating, we get

$$y e^{\int P dx} = \int Q e^{\int P dx} dx + c \text{ is the required solution.}$$

**Note:** We remember the solution of the above equation as

$$y(\text{I.F}) = \int Q(\text{I.F}) dx + c$$

**Illustration 8 :** Solve  $2x \frac{dy}{dx} = y + 6x^{5/2} - 2\sqrt{x}$

**Sol.** The given equation can be written as

$$\frac{dy}{dx} + \left( \frac{-1}{2x} \right) y = 3x^{3/2} - \frac{1}{\sqrt{x}} \quad \dots (i)$$

This is the form of  $\frac{dy}{dx} + P y = Q$

$$\text{Hence I.F} = e^{\int \frac{-1}{2x} dx} = e^{-\frac{1}{2} \log x} = \frac{1}{\sqrt{x}} \quad \dots (ii)$$

$$\text{From (i) and (ii), we get } y = \frac{3}{2} x^{5/2} - \sqrt{x} \log x + c\sqrt{x}$$

## (ii) Differential equation reducible to the linear form:

Sometimes equations which are not linear can be reduced to the linear form by suitable transformation.

$$\text{Here, } f'(y) \frac{dy}{dx} + f(y) P(x) = Q(x) \quad \dots (i)$$

$$\text{Let, } f(y) = u \Rightarrow f'(y) dy = du$$

Then (i) reduces to

$$\frac{du}{dx} + u P(x) = Q(x) \text{ Which is of the linear differential equation form.}$$

**Illustration 9:** Solve  $\sec^2 \theta d\theta + \tan \theta (1 - r \tan \theta) dr = 0$

**Sol.** The given equation can be written as

$$\frac{d\theta}{dr} + \frac{\tan \theta}{\sec^2 \theta} = \frac{r \tan^2 \theta}{\sec^2 \theta}$$

$$\text{or, } \left( \frac{\sec^2 \theta}{\tan^2 \theta} \right) \frac{d\theta}{dr} + \frac{1}{\tan \theta} = r$$

$$\text{or, } \cos \sec^2 \theta \frac{d\theta}{dr} + \cot \theta = r \quad \dots (i)$$

$$\text{Let } \cot \theta = u$$

$$\Rightarrow -\cos \sec^2 \theta d\theta = du$$

Then (i) reduces to

$$-\frac{du}{dr} + u = r \quad \text{or,} \quad \frac{du}{dr} - u = -r \quad \dots (ii)$$

Which is a linear differential equation.

$$\text{So, } \text{I.F} = e^{\int -1 dr} = e^{-r} \quad \dots (iii)$$

Form (ii) and (iii), we get

$$u e^{-r} = -\int r e^{-r} dr = r e^{-r} + \int e^{-r} dr, \text{ by parts}$$

$$= re^{-r} - e^{-r} + c \quad \text{or,} \quad u = r - 1 + ce^r$$

$$\text{or,} \quad \cot \theta = r - 1 + ce^r$$

### (iii) Extended form of linear equations :

#### Bernoulli's equation:

An equation of the form  $\frac{dy}{dx} + P y = Q y^n$ , where P and

Q are function of x alone or constants and n is constant, other than 0 and 1, is called a Bernoulli's equation.

$$\text{Here } \frac{dy}{dx} + P y = Q y^n$$

Dividing by  $y^n$ , we get

$$\frac{1}{y^n} \frac{dy}{dx} + P \cdot \frac{1}{y^{n-1}} = Q$$

$$\text{Putting } \frac{1}{y^{n-1}} = v \quad \text{and differentiating w.r.t } x,$$

$$\text{we get } -\frac{(n-1)}{y^n} \frac{dy}{dx} = \frac{dv}{dx}$$

$$\text{or,} \quad \frac{1}{y^n} \frac{dy}{dx} = \frac{-1}{n-1} \frac{dv}{dx}$$

$$\text{or,} \quad \frac{dv}{dx} = (1-n) y^{-n} \frac{dy}{dx}, \text{ the equation becomes}$$

$$\frac{dv}{dx} + (1-n) P v = Q(1-n)$$

Which is a linear equation with v as independent variable.

**Illustration 10 :** Solve  $\cos^2 x \frac{dy}{dx} - y \tan 2x = \cos^4 x$ , where

$$|x| < \frac{\pi}{4} \text{ and } y\left(\frac{\pi}{4}\right) = \frac{3\sqrt{3}}{8}$$

**Sol.** The given equation can be written as

$$\frac{dy}{dx} - y \tan 2x \sec^2 x = \cos^2 x$$

$$\text{This is the form of } \frac{dy}{dx} + P y = Q$$

$$\text{Here } P = -\tan 2x \sec^2 x, Q = \cos^2 x$$

$$\therefore \int P dx = -\int \tan 2x \sec^2 x dx$$

$$= -\int \frac{2 \tan x}{1 - \tan^2 x} \sec^2 x dx$$

$$= \int \frac{dt}{t} \quad \text{Putting } 1 - \tan^2 x = t$$

$$\therefore -2 \tan x \sec^2 x dx = dt$$

$$= \log t = \log (1 - \tan^2 x)$$

$$\therefore \text{I. F.} = e^{\int P dx} = e^{\log (1 - \tan^2 x)} = 1 - \tan^2 x$$

$\therefore$  The solution is

$$y (1 - \tan^2 x) = \int (1 - \tan^2 x) \cos^2 x dx + c$$

$$= \int \cos 2x dx + c = \frac{\sin 2x}{2} + c \quad \dots(i)$$

$$\text{Given, At } x = \frac{\pi}{4}, y = \frac{3\sqrt{3}}{8}, \text{ then } \frac{3\sqrt{3}}{8} \left(1 - \frac{1}{3}\right) = \frac{\sqrt{3}}{4} + c$$

$$\text{or } c = 0$$

Hence from (i),

$$y (1 - \tan^2 x) = \frac{\sin 2x}{2}$$

$$\text{or } y = \frac{\sin 2x}{2 (1 - \tan^2 x)}$$

# DIFFERENTIAL EQUATION

## Method – 4

### Exact differential equation:

A differential equation is said to be exact if it can be derived from its solution (primitive) directly by differentiation, without any elimination, multiplication etc.

For example, the differential equation  $x dy + y dx = 0$  is an exact differential equation as it is derived by direct differentiation for its solution, the function  $xy = c$

**Illustration 11 :** Solve  $(1 + xy) y dx + (1 - xy) x dy = 0$

**Sol.** The given equation can be written as

$$y dx + xy^2 dx + x dy - x^2y dy = 0$$

$$\text{or, } (y dx + x dy) + xy (y dx - x dy) = 0$$

$$\text{or, } d(xy) + xy (y dx - x dy) = 0$$

Dividing by  $x^2y^2$ , we get

$$\frac{d(xy)}{x^2y^2} + \frac{y dx - x dy}{xy} = \text{or,}$$

$$\frac{d(xy)}{x^2y^2} + \frac{dx}{x} - \frac{dy}{y} =$$


Integrating, we get

$$-\frac{1}{xy} + \log x - \log y = c$$

Which is the required solution.

### Application of differential equations :

In solving some geometrical problems, the following results are very helpful.



Let PT and PN be the tangent and the normal at  $P(x, y)$ . Let the tangent at P makes an angle  $\theta$  with the x-axis.

$$\text{Then the slope of the tangent at } P = \tan \theta = \left( \frac{dy}{dx} \right)_P$$

$$\text{and the slope of the normal at } P = -\frac{1}{\left( \frac{dy}{dx} \right)_P}$$

Equation of the tangent at  $P(x, y)$  is

$$Y - y = \left( \frac{dy}{dx} \right)_P (X - x)$$

Equation of the normal at  $P(x, y)$  is

$$Y - y = -\frac{1}{\left( \frac{dy}{dx} \right)_P} (X - x)$$

$$\text{From } \Delta PGT \quad \sin \theta = \frac{PG}{PT} = \frac{y}{PT}$$

$$\therefore PT = y \operatorname{cosec} \theta \text{ (length of the tangent)}$$

$$= y \frac{\sqrt{1 + \tan^2 \theta}}{\tan \theta} = y \frac{\sqrt{1 + \left( \frac{dy}{dx} \right)^2}}{\frac{dy}{dx}}$$

$$\text{And, } \tan \theta = \frac{PG}{TG} = \frac{y}{TG}$$

$$\Rightarrow \boxed{TG = y \cot \theta \text{ (length of the sub tangent)} = \frac{y}{\frac{dy}{dx}}}$$

$$\text{From } \Delta PGN \quad \cos \theta = \frac{PG}{PN} = \frac{y}{PN}$$


$$\Rightarrow \boxed{PN = y \sec \theta \text{ (length of the normal)}} \\ = y \sqrt{1 + \tan^2 \theta} = y \sqrt{1 + \left(\frac{dy}{dx}\right)^2}$$

$$\tan \theta = \frac{GN}{y}$$

$$\Rightarrow \boxed{GN = y \tan \theta = y \frac{dy}{dx} \text{ (length of the sub normal)}}$$

**Illustration 12 :** If the length of the sub-normal at any point P on the curve is directly proportional to  $OP^2$ , where O is the origin, then form the differential equation of the family of curves and hence find the family of curves.

**Sol.** Here  $AB = y \tan \theta = y \frac{dy}{dx}$



$$\text{Also } OP^2 = x^2 + y^2$$

Given, length of the subnormal = k.  $OP^2$

$$\text{or, } y \frac{dy}{dx} = k(x^2 + y^2)$$

$$\text{or, } 2y \frac{dy}{dx} - 2ky^2 = 2kx^2 \quad \dots(i)$$

$$\text{Let } y^2 = t \Rightarrow 2y \frac{dy}{dx} = \frac{dt}{dx} \quad \dots(ii)$$

$$\text{From (i) and (ii), we get } \frac{dt}{dx} - 2kt = 2kx^2$$

Which is a linear differential equation.

$$\therefore I.F = e^{\int -2k dx} = e^{-2kx}$$

$\therefore$  The solution is

$$t \cdot e^{-2kx} = \int 2kx^2 e^{-2kx} dx + c$$

$$= 2k \left[ x^2 \frac{e^{-2kx}}{-2k} + \frac{2}{2k} \int x e^{-2kx} dx \right]$$

$$= 2k \left[ x^2 \frac{e^{-2kx}}{-2k} + \frac{1}{k} \left\{ x \frac{e^{-2kx}}{-2k} + \frac{1}{2k} \int e^{-2kx} dx \right\} \right]$$

$$= -x^2 e^{-2kx} - \frac{x e^{-2kx}}{k} + \frac{1}{k} \frac{e^{-2kx}}{2k} + c \text{ or,}$$

$$y^2 = -x^2 - \frac{x}{k} + \frac{1}{2k^2} + c e^{2kx}$$

**Thank You**  
for downloading the PDF

**FREE** LIVE ONLINE

# MASTER CLASSES

FREE Webinars by Expert Teachers



Vedantu

## FREE MASTER CLASS SERIES

- ✓ For **Grades 6-12th** targeting **JEE, CBSE, ICSE** & much more
- ✓ **Free 60 Minutes Live Interactive** classes everyday
- ✓ Learn from the **Master Teachers** - India's best

Register for **FREE**

Limited Seats!