



1

COMPLEX NUMBERS

We started our study of number systems with the set of natural numbers, then the number zero was included to form the system of whole numbers; negative numbers were defined. Thus, we extended our number system to whole numbers and integers.

To solve the problems of the type $p \div q$ we included rational numbers in the system of integers. The system of rational numbers have been extended further to irrational numbers as all lengths cannot be measured in terms of lengths expressed in rational numbers. Rational and irrational numbers taken together are termed as real numbers. But the system of real numbers is not sufficient to solve all algebraic equations. There are no real numbers which satisfy the equation $x^2 + 1 = 0$ or $x^2 = -1$. In order to solve such equations, i.e., to find square roots of negative numbers, we extend the system of real numbers to a new system of numbers known as complex numbers. In this lesson the learner will be acquainted with complex numbers, its representation and algebraic operations on complex numbers.



OBJECTIVES

After studying this lesson, you will be able to:

- describe the need for extending the set of real numbers to the set of complex numbers;
- define a complex number and cite examples;
- identify the real and imaginary parts of a complex number;
- state the condition for equality of two complex numbers;
- recognise that there is a unique complex number $x + iy$ associated with the point $P(x, y)$ in the Argand Plane and vice-versa;
- define and find the conjugate of a complex number;
- define and find the modulus and argument of a complex number;

MODULE - I
Algebra

Notes

- represent a complex number in the polar form;
- perform algebraic operations (addition, subtraction, multiplication and division) on complex numbers;
- state and use the properties of algebraic operations (closure, commutativity, associativity, identity, inverse and distributivity) of complex numbers; and
- state and use the following properties of complex numbers in solving problems:

$$(i) \quad |z| = 0 \Leftrightarrow z = 0 \text{ and } z_1 = z_2 \Rightarrow |z_1| = |z_2|$$

$$(ii) \quad |z| = |-z| = |\bar{z}|$$

$$(iii) \quad |z_1 + z_2| \leq |z_1| + |z_2|$$

$$(iv) \quad |z_1 \cdot z_2| = |z_1| \cdot |z_2|$$

$$(v) \quad \left| \frac{z_1}{z_2} \right| = \frac{|z_1|}{|z_2|} \quad (z_2 \neq 0)$$

EXPECTED BACKGROUND KNOWLEDGE

- Properties of real numbers.
- Solution of linear and quadratic equations
- Representation of a real number on the number line
- Representation of point in a plane.

1.1 COMPLEX NUMBERS

Consider the equation $x^2 + 1 = 0$(A)

This can be written as $x^2 = -1$
or $x = \pm\sqrt{-1}$

But there is no real numbers which satisfy $x^2 = -1$. In other words, we can say that there is no real numbers whose square is -1 . In order to solve such equations, let us imagine that there exist a number 'i' which equal to $\sqrt{-1}$.

In 1748, a great mathematician, L. Euler named a number 'i' as *Iota* whose square is -1 . This *Iota* or 'i' is defined as imaginary unit. With the introduction of the new symbol 'i', we can interpret the square root of a negative number as a product of a real number with i.

Therefore, we can denote the solution of (A) as $x = \pm i$

Thus, $-4 = 4(-1)$

$$\therefore \sqrt{-4} = \sqrt{(-1)(4)} = \sqrt{i^2 \cdot 2^2} = 2i$$

Conventionally written as $2i$.

$$\text{So, we have } \sqrt{-4} = 2i, \quad \sqrt{-7} = \sqrt{7}i$$

$\sqrt{-4}$, $\sqrt{-7}$ are all examples of complex numbers.

Consider another quadratic equation:

$$x^2 - 6x + 13 = 0$$

This can be solved as under:

$$(x - 3)^2 + 4 = 0$$

$$\text{or, } (x - 3)^2 = -4$$

$$\text{or, } x - 3 = \pm 2i$$

$$\text{or, } x = 3 \pm 2i$$

We get numbers of the form $x + yi$ where x and y are real numbers and $i = \sqrt{-1}$.

Any number which can be expressed in the form $a + bi$ where a, b are real numbers and $i = \sqrt{-1}$, is called a complex number.

A complex number is, generally, denoted by the letter z .

i.e. $z = a + bi$, ' a ' is called the real part of z and is written as $\text{Re}(a + bi)$ and ' b ' is called the imaginary part of z and is written as $\text{Imag}(a + bi)$.

If $a = 0$ and $b \neq 0$, then the complex number becomes bi which is a purely imaginary complex number.

$-7i$, $\frac{1}{2}i$, $\sqrt{3}i$ and πi are all examples of purely imaginary numbers.

If $a \neq 0$ and $b = 0$ then the complex number becomes ' a ' which is a real number.

5 , 2.5 and $\sqrt{7}$ are all examples of real numbers.

If $a = 0$ and $b = 0$, then the complex number becomes 0 (zero). Hence the real numbers are particular cases of complex numbers.

Example 1.1 Simplify each of the following using ' i '.

$$(i) \quad \sqrt{-36} \qquad (ii) \quad \sqrt{25} \cdot \sqrt{-4}$$

Solution: (i) $\sqrt{-36} = \sqrt{36(-1)} = 6i$

$$(ii) \quad \sqrt{25} \cdot \sqrt{-4} = 5 \times 2i = 10i$$

1.2 POSITIVE INTEGRAL POWERS OF i

We know that

$$i^2 = -1$$

$$i^3 = i^2 \cdot i = -1 \cdot i = -i$$

$$i^4 = (i^2)^2 = (-1)^2 = 1$$



MODULE - I
Algebra

Notes

$$i^5 = (i^2)^2 \cdot i = 1 \cdot i = i$$

$$i^6 = (i^2)^3 = (-1)^3 = -1$$

$$i^7 = (i^2)^3(i) = -i$$

$$i^8 = (i^2)^4 = 1$$

Thus, we find that any higher powers of 'i' can be expressed in terms of one of four values i, -1, -i, 1

If n is a positive integer such that $n > 4$, then to find i^n , we first divide n by 4.

Let m be the quotient and r be the remainder.

Then $n = 4m + r$, where $0 \leq r < 4$.

Thus,

$$\begin{aligned} i^n &= i^{(4m+r)} = i^{4m} \cdot i^r \\ &= (i^4)^m \cdot i^r \\ &= i^r \quad (\because i^4 = 1) \end{aligned}$$

Note : For any two real numbers a and b, $\sqrt{a} \times \sqrt{b} = \sqrt{ab}$ is true only when atleast one of a and b is either 0 or positive.

$$\begin{aligned} \text{If fact } &\sqrt{-a} \times \sqrt{-b} \\ &= i\sqrt{a} \times i\sqrt{b} = i^2 \sqrt{ab} \\ &= -\sqrt{ab} \quad \text{where a and b are positive real numbers.} \end{aligned}$$

Example 1.2 Find the value of $1 + i^{10} + i^{20} + i^{30}$

Solution: $1 + i^{10} + i^{20} + i^{30}$

$$\begin{aligned} &= 1 + (i^2)^5 + (i^2)^{10} + (i^2)^{15} \\ &= 1 + (-1)^5 + (-1)^{10} + (-1)^{15} \\ &= 1 + (-1) + 1 + (-1) \\ &= 1 - 1 + 1 - 1 \\ &= 0 \end{aligned}$$

Thus, $1 + i^{10} + i^{20} + i^{30} = 0$.



Example 1.3 Express $8i^3 + 6i^{16} - 12i^{11}$ in the form of $a + bi$

Solution: $8i^3 + 6i^{16} - 12i^{11}$ can be written as $8(i^2).i + 6(i^2)^8 - 12(i^2)^5.i$

$$= 8(-1).i + 6(-1)^8 - 12(-1)^5.i$$

$$= -8i + 6 - 12(-1).i$$

$$= -8i + 6 + 12i$$

$$= 6 + 4i$$

which is of the form of $a + bi$ where 'a' is 6 and 'b' is 4.



CHECK YOUR PROGRESS 1.1

- Simplify each of the following using 'i'.
(a) $\sqrt{-27}$ (b) $-\sqrt{-9}$ (c) $\sqrt{-13}$
- Express each of the following in the form of $a + bi$
(a) 5 (b) $-3i$ (c) 0
- Simplify $10i^3 + 6i^{13} - 12i^{10}$
- Show that $i^m + i^{m+1} + i^{m+2} + i^{m+3} = 0$ for all $m \in \mathbb{N}$.

1.3 CONJUGATE OF A COMPLEX NUMBER

Consider the equation:

$$x^2 - 6x + 25 = 0 \quad \dots \quad (i)$$

or, $(x - 3)^2 + 16 = 0$

or, $(x - 3)^2 = -16$

or, $(x - 3) = \pm\sqrt{-16} = \pm\sqrt{16.(-1)}$

or, $x = 3 \pm 4i$

The roots of the above equation (i) are $3 + 4i$ and $3 - 4i$.

Consider another equation:

$$x^2 + 2x + 2 = 0 \quad \dots \quad (ii)$$

or, $(x + 1)^2 + 1 = 0$

or, $(x + 1)^2 = -1$

or, $(x + 1) = \pm\sqrt{-1} = \pm i$

or, $x = -1 \pm i$

MODULE - I
Algebra

Notes

The roots of the equation (ii) are $-1+i$ and $-1-i$.

Do you find any similarity in the roots of (i) and (ii)?

The equations (i) and (ii) have roots of the type $a + bi$ and $a - bi$. Such roots are known as conjugate roots and read as $a + bi$ is conjugate to $a - bi$ and vice-versa.

The complex conjugate (or simply conjugate) of a complex number $z = a + bi$ is defined as the complex number $a - bi$ and is denoted by \bar{z} .

Thus, if $z = a + bi$ then $\bar{z} = a - bi$.

Note : The conjugate of a complex number is obtained by changing the sign of the imaginary part.

Following are some examples of complex conjugates:

- (i) If $z = 2 + 3i$, then $\bar{z} = 2 - 3i$
- (ii) If $z = 1 - i$, then $\bar{z} = 1 + i$
- (iii) If $z = -2 + 10i$, then $\bar{z} = -2 - 10i$

1.3.1 PROPERTIES OF COMPLEX CONJUGATES

- (i) If z is a real number then $z = \bar{z}$ i.e., the conjugate of a real number is the number itself.

For example, let $z = 5$

This can be written as

$$z = 5 + 0i$$

$$\therefore \bar{z} = 5 - 0i = 5$$

$$\therefore z = 5 = \bar{z}.$$

- (ii) If z is a purely imaginary number then $\bar{z} = -z$

For example, if $z = 3i$

This can be written as

$$z = 0 + 3i$$

$$\therefore \bar{z} = 0 - 3i = -3i$$

$$= -z$$

$$\therefore \bar{z} = -z.$$



(iii) Conjugate of the conjugate of a complex number is the number itself.

i.e., $\overline{(\overline{z})} = z$

For example, if $z = a + bi$ then

$$\overline{z} = a - bi$$

Again $\overline{(\overline{z})} = \overline{(a - bi)} = a + bi$

$$= z$$

$$\therefore \overline{(\overline{z})} = z$$

Example 1.4 Find the conjugate of each of the following complex number:

(i) $3 - 4i$

(ii) $2i$

(iii) $(2 + i)^2$

(iv) $\frac{i + 1}{2}$

Solution : (i) Let $z = 3 - 4i$

then $\overline{z} = \overline{(3 - 4i)} = 3 + 4i$

Hence, $3 + 4i$ is the conjugate of $3 - 4i$.

(ii) Let $z = 2i$ or $0 + 2i$

then $\overline{z} = \overline{(0 + 2i)} = 0 - 2i$

Hence, $-2i$ is the conjugate of $2i$.

(iii) Let $z = (2 + i)^2$

i.e. $z = (2)^2 + (i)^2 + 2(2)(i)$

$$= 4 - 1 + 4i$$

$$= 3 + 4i$$

Then $\overline{z} = \overline{(3 + 4i)} = 3 - 4i$

Hence, $3 - 4i$ is the conjugate of $(2 + i)^2$

(iv) Let $z = \frac{i + 1}{2} = \frac{1}{2} + \frac{1}{2}i$

MODULE - I
Algebra

Notes

$$\text{then } \bar{z} = \overline{\left(\frac{1}{2} + \frac{1}{2}i\right)} = \frac{1}{2} - \frac{1}{2}i$$

Hence, $\frac{1}{2} - \frac{i}{2}$ or $\frac{-i+1}{2}$ is the conjugate of $\frac{i+1}{2}$

1.4 GEOMETRICAL REPRESENTATION OF A COMPLEX NUMBER

Let $z = a + bi$ be a complex number. Let two mutually perpendicular lines xox' and yoy' be taken as x-axis and y-axis respectively, O being the origin.

Let P be any point whose coordinates are (a,b). We say that the complex $z = a + bi$ is represented by the point P (a, b) as shown in Fig. 1.1




Fig. 1.1

If $b = 0$, then z is real and the point representing complex number $z = a + 0i$ is denoted by (a, 0). This point (a, 0) lies on the x-axis.

So, xox' is called the real axis. In the Fig. 1.2 the point Q (a, 0) represent the complex number $z = a + 0i$.




Fig. 1.2

If $a = 0$, then z is purely imaginary and the point representing complex number $z = 0 + bi$ is denoted by (0, b). The point (0, b) lies on the y-axis.




Fig. 1.3

So, yoy' is called the imaginary axis. In Fig. 1.3, the point R (0, b) represents the complex number $z = 0 + bi$.

The plane of two axes representing complex numbers as points is called the complex plane or Argand Plane.

The diagram which represents complex number in the Argand Plane is called Argand Diagram.

Example 1.5

Represent complex numbers $2 + 3i$ and $3 + 2i$ in the same Argand Plane.

Solution :

1. $2 + 3i$ is represented by the point A (2,3)
2. $3 + 2i$ is represented by the point B (3, 2) Clearly, the points A and B are different




Fig. 1.4

Example 1.6

Represent complex numbers $2 + 3i$ and $-2 - 3i$ in the same Argand Plane.

Solution:

- $2 + 3i$ is represented by the point $P(2, 3)$
- $-2 - 3i$ is represented by the point $Q(-2, -3)$




Fig. 1.5

Points P and Q are different and lie in the I quadrant and III quadrant respectively.

Example 1.7

Represent complex numbers $2 + 3i$ and $2 - 3i$ in the same Argand Plane

Solution:

- $2 + 3i$ is represented by the point $R(2, 3)$
- $2 - 3i$ is represented by the point $S(2, -3)$




Fig. 1.6

Example 1.8

Represent complex numbers $2 + 3i$, $-2 - 3i$, $2 - 3i$ in the same Argand Plane

Solution:

- $2 + 3i$ is represented by the point $P(2, 3)$
- $-2 - 3i$ is represented by the point $Q(-2, -3)$
- $2 - 3i$ is represented by the point $R(2, -3)$




Fig. 1.7

1.5 MODULUS OF A COMPLEX NUMBER

We have learnt that any complex number $z = a + bi$ can be represented by a point in the Argand Plane. How can we find the distance of the point from the origin? Let $P(a, b)$ be a point in the plane representing $a + bi$. Draw perpendiculars PM and PL on x-axis and y-axis respectively. Let $OM = a$ and $MP = b$. We have to find the distance of P from the origin.



MODULE - I
Algebra

Notes

$$\begin{aligned}\therefore OP &= \sqrt{OM^2 + MP^2} \\ &= \sqrt{a^2 + b^2}\end{aligned}$$

OP is called the modulus or absolute value of the complex number $a + bi$.

\therefore Modulus of any complex number z such that $z = a + bi$, $a \in \mathbb{R}$, $b \in \mathbb{R}$ is denoted by

$$|z| \text{ and is given by } \sqrt{a^2 + b^2}$$

$$\therefore |z| = |a + ib| = \sqrt{a^2 + b^2}$$

1.5.1 Properties of Modulus

$$(a) \quad |z| = 0 \Leftrightarrow z = 0.$$

Proof : Let $z = a + bi$, $a \in \mathbb{R}$, $b \in \mathbb{R}$

$$\text{then } |z| = \sqrt{a^2 + b^2}$$

$$|z| = 0 \Leftrightarrow a^2 + b^2 = 0$$

$$\Leftrightarrow a = 0 \text{ and } b = 0 \text{ (since } a^2 \text{ and } b^2 \text{ both are positive)}$$

$$\Leftrightarrow z = 0$$

$$(b) \quad |z| = |\bar{z}|$$

Proof : Let $z = a + bi$

$$\text{then } |z| = \sqrt{a^2 + b^2}$$

$$\text{Now, } \bar{z} = a - bi$$

$$\therefore |\bar{z}| = \sqrt{a^2 + (-b)^2} = \sqrt{a^2 + b^2}$$

$$\text{Thus, } |z| = \sqrt{a^2 + b^2} = |\bar{z}| \quad \dots(i)$$

$$(c) \quad |z| = |-z|$$

Proof : Let $z = a + bi$ then $|z| = \sqrt{a^2 + b^2}$




Fig. 1.8



$$\begin{aligned} -z = -a - bi \text{ then } |-z| &= \sqrt{(-a)^2 + (-b)^2} \\ &= \sqrt{a^2 + b^2} \end{aligned}$$

$$\text{Thus, } |z| = \sqrt{a^2 + b^2} = |-z| \quad \dots(\text{ii})$$

By (i) and (ii) it can be proved that

$$|z| = |-z| = |\bar{z}| \quad \dots(\text{iii})$$

Now, we consider the following examples:

Example 1.9 Find the modulus of z and \bar{z} if $z = -4 + 3i$

Solution : $z = -4 + 3i$, then $|z| = \sqrt{(-4)^2 + (3)^2}$

$$= \sqrt{16 + 9} = \sqrt{25} = 5$$

and $\bar{z} = -4 - 3i$

$$\text{then, } |\bar{z}| = \sqrt{(-4)^2 + (-3)^2} = \sqrt{16 + 9} = \sqrt{25} = 5$$

Thus, $|z| = 5 = |\bar{z}|$

Example 1.10 Find the modulus of z and $-z$ if $z = 5 + 2i$

Solution: $z = 5 + 2i$, then $-z = -5 - 2i$

$$|z| = \sqrt{5^2 + 2^2} = \sqrt{29} \text{ and } |-z| = \sqrt{(-5)^2 + (-2)^2} = \sqrt{29}$$

$$\text{Thus, } |z| = \sqrt{29} = |-z|$$

Example 1.11 Find the modulus of z , $-z$ and \bar{z} where $z = 1 + 2i$

Solution : $z = 1 + 2i$ then $-z = -1 - 2i$ and $\bar{z} = 1 - 2i$

$$|z| = \sqrt{1^2 + 2^2} = \sqrt{5}$$

$$|-z| = \sqrt{(-1)^2 + (-2)^2} = \sqrt{5}$$

$$\text{and } |\bar{z}| = \sqrt{(1)^2 + (-2)^2} = \sqrt{5}$$

$$\text{Thus, } |z| = |-z| = \sqrt{5} = |\bar{z}|$$

MODULE - I
Algebra

Notes

Example 1.12 Find the modulus of :

- (i) $1 + i$ (ii) 2π (iii) 0 (iv) $\frac{1}{2}i$

Solution: (i) Let $z = 1 + i$

$$\text{then } |z| = \sqrt{1^2 + 1^2} = \sqrt{2}$$

$$\text{Thus, } |1 + i| = \sqrt{2}$$

(ii) Let $z = 2\pi$ or $2\pi + 0i$

$$\text{Then } |z| = \sqrt{(2\pi)^2 + (0)^2} = 2\pi$$

$$\text{Thus, } |2\pi| = 2\pi.$$

If z is real then $|z| = z$ (iii) $z = 0$ or $0 + 0i$

$$\text{then } |z| = \sqrt{(0)^2 + (0)^2} = 0$$

$$\text{Thus, } |z| = 0$$

If z is 0 then $|z| = 0$ (iv) Let $z = -\frac{1}{2}i$ or $0 - \frac{1}{2}i$

$$\text{then } |z| = \sqrt{0^2 + \left(-\frac{1}{2}\right)^2} = \frac{1}{2}$$

$$\text{Thus, } \left| -\frac{1}{2}i \right| = \frac{1}{2}.$$

If z is purely imaginary number, then $z \neq |z|$.**Example 1.13** Find the absolute value of the conjugate of the complex number

$$z = -2 + 3i$$

Solution : Let $z = -2 + 3i$ then $\bar{z} = -2 - 3i$

Absolute value of $\bar{z} = |\bar{z}| = |-2 - 3i| = \sqrt{(-2)^2 + (-3)^2} = \sqrt{4 + 9} = \sqrt{13}$

Example 1.14 Find the modulus of the complex numbers shown in an Argand Plane (Fig. 1.9)

Solution: (i) P(4, 3) represents the complex number $z = 4 + 3i$

$$\therefore |z| = \sqrt{4^2 + 3^2} = \sqrt{25}$$

$$\text{or } |z| = 5$$

(ii) Q(-4, 2) represents the complex number $z = -4 + 2i$

$$\therefore |z| = \sqrt{(-4)^2 + 2^2} = \sqrt{16 + 4} = \sqrt{20}$$

$$\text{or } |z| = 2\sqrt{5}$$

(iii) R(-1, -3) represents the complex number $z = -1 - 3i$

$$\therefore |z| = \sqrt{(-1)^2 + (-3)^2} = \sqrt{1 + 9}$$

$$\text{or } |z| = \sqrt{10}$$

(iv) S(3, -3) represents the complex number $z = 3 - 3i$

$$\therefore |z| = \sqrt{(3)^2 + (-3)^2} = \sqrt{9 + 9}$$

$$\text{or } |z| = \sqrt{18} = 3\sqrt{2}$$




Fig. 1.9



Notes



CHECK YOUR PROGRESS 1.2

1. Find the conjugate of each of the following:

- (a) $-2i$ (b) $-5 - 3i$ (c) $-\sqrt{2}$ (d) $(-2 + i)^2$

2. Represent the following complex numbers on Argand Plane :

- (a) (i) $2 + 0i$ (ii) $-3 + 0i$ (iii) $0 - 0i$ (iv) $3 - 0i$

MODULE - I
Algebra

Notes

- (b) (i) $0 + 2i$ (ii) $0 - 3i$ (iii) $4i$ (iv) $-5i$
- (c) (i) $2 + 5i$ and $5 + 2i$ (ii) $3 - 4i$ and $-4 + 3i$
- (iii) $-7 + 2i$ and $2 - 7i$ (iv) $-2 - 9i$ and $-9 - 2i$
- (d) (i) $1 + i$ and $-1 - i$ (ii) $6 + 5i$ and $-6 - 5i$
- (iii) $-3 + 4i$ and $3 - 4i$ (iv) $4 - i$ and $-4 + i$
- (e) (i) $1 + i$ and $1 - i$ (ii) $-3 + 4i$ and $-3 - 4i$
- (iii) $6 - 7i$ and $6 + 7i$ (iv) $-5 - i$ and $-5 + i$

3. (a) Find the modulus of following complex numbers :

- (i) 3 (ii) $(i + 1)(2 - i)$ (iii) $2 - 3i$ (iv) $4 + \sqrt{5}i$
- (b) For the following complex numbers, verify that $|z| = |\bar{z}|$
- (i) $-6 + 8i$ (ii) $-3 - 7i$
- (c) For the following complex numbers, verify that $|z| = |-z|$
- (i) $14 + i$ (ii) $11 - 2i$
- (d) For the following complex numbers, verify that $|z| = |-z| = |\bar{z}|$
- (i) $2 - 3i$ (ii) $-6 - i$ (iii) $7 - 2i$

1.6 EQUALITY OF TWO COMPLEX NUMBERS

Let us consider two complex numbers $z_1 = a + bi$ and $z_2 = c + di$ such that $z_1 = z_2$

we have $a + bi = c + di$

$$\text{or } (a - c) + (b - d)i = 0 = 0 + 0i$$

Comparing real and imaginary parts on both sides, we have

$$a - c = 0, \text{ or } a = c$$

$$\Rightarrow \text{real part of } z_1 = \text{real part of } z_2$$

$$\text{and } b - d = 0 \quad \text{or } b = d$$

$$\Rightarrow \text{imaginary part of } z_1 = \text{imaginary part of } z_2$$

Therefore, we can conclude that two complex numbers are equal if and only if their real parts and imaginary parts are respectively equal.

In general $a + bi = c + di$ if and only if $a = c$ and $b = d$.



Properties: $z_1 = z_2 \Rightarrow |z_1| = |z_2|$

Let $z_1 = a + bi$ and $z_2 = c + di$

$z_1 = z_2$ gives $a = c$ and $b = d$

Now $|z_1| = \sqrt{a^2 + b^2}$ and $|z_2| = \sqrt{c^2 + d^2}$

$$= \sqrt{a^2 + b^2} \quad [\text{since } a = c \text{ and } b = d]$$

$\Rightarrow |z_1| = |z_2|$

Example 1.15 For what value of x and y , $5x + 6yi$ and $10 + 18i$ are equal?

Solution : It is given that

$$5x + 6yi = 10 + 18i$$

Comparing real and imaginary parts, we have

$$5x = 10 \quad \text{or } x = 2$$

$$\text{and } 6y = 18 \quad \text{or } y = 3$$

For $x = 2$ and $y = 3$, the given complex numbers are equal.

1.7 ADDITION OF COMPLEX NUMBERS

If $z_1 = a + bi$ and $z_2 = c + di$ are two complex numbers then their sum $z_1 + z_2$ is defined by

$$z_1 + z_2 = (a + c) + (b + d)i$$

For example, if $z_1 = 2 + 3i$ and $z_2 = -4 + 5i$,

$$\text{then } z_1 + z_2 = [2 + (-4)] + [3 + 5]i$$

$$= -2 + 8i.$$

Example 1.16 Simplify

$$(i) \quad (3 + 2i) + (4 - 3i)$$

$$(ii) \quad (2 + 5i) + (-3 - 7i) + (1 - i)$$

Solution : (i) $(3 + 2i) + (4 - 3i) = (3 + 4) + (2 - 3)i = 7 - i$

$$(ii) \quad (2 + 5i) + (-3 - 7i) + (1 - i) = (2 - 3 + 1) + (5 - 7 - 1)i$$

$$= 0 - 3i$$

$$\text{or } (2 + 5i) + (-3 - 7i) + (1 - i) = -3i$$

MODULE - I
Algebra

Notes

1.7.1 Geometrical Representation of Addition of Two Complex Numbers

Let two complex numbers z_1 and z_2 be represented by the points $P(a, b)$ and $Q(c, d)$.

Their sum, $z_1 + z_2$ is represented by the point $R(a + c, b + d)$ in the same Argand Plane.

Join OP , OQ , OR , PR and QR .

Draw perpendiculars PM , QN , RL from P , Q , R respectively on X -axis.

Draw perpendicular PK to RL

In $\triangle QON$

$$ON = c$$

$$\text{and } QN = d.$$

In $\triangle ROL$

$$RL = b + d$$

$$\text{and } OL = a + c$$

$$\text{Also } PK = ML$$

$$= OL - OM$$

$$= a + c - a$$

$$= c = ON$$

In $\triangle POM$

$$PM = b$$

$$OM = a$$




Fig. 1.10

$$RK = RL - KL$$

$$= RL - PM$$

$$= b + d - b$$

$$= d = QN.$$

In $\triangle QON$ and $\triangle RPK$,

$$ON = PK, QN = RK \text{ and } \angle QNO = \angle RKP = 90^\circ$$

$$\therefore \triangle QON \cong \triangle RPK$$

$$\therefore OQ = PR \text{ and } OQ \parallel PR$$

\Rightarrow $OPRQ$ is a parallelogram and OR its diagonal.

Therefore, we can say that the sum of two complex numbers is represented by the diagonal of a parallelogram.



Example 1.17 Prove that $|z_1 + z_2| \leq |z_1| + |z_2|$

Solution: We have proved that the sum of two complex numbers z_1 and z_2 represented by the diagonal of a parallelogram OPRQ (see fig. 1.11).

In $\triangle OPR$

$$OR \leq OP + PR$$

or $OR \leq OP + OQ$ (since $OQ = PR$)

or $|z_1 + z_2| \leq |z_1| + |z_2|$

Example 1.18 If $z_1 = 2 + 3i$ and $z_2 = 1 + i$,

verify that $|z_1 + z_2| \leq |z_1| + |z_2|$

Solution: $z_1 = 2 + 3i$ and $z_2 = 1 + i$ represented by the points (2, 3) and (1, 1) respectively. Their sum ($z_1 + z_2$) will be represented by the point (2+1, 3+1) i.e. (3, 4)

Verification

$$|z_1| = \sqrt{2^2 + 3^2} = \sqrt{13} = 3.6 \text{ approx.}$$

$$|z_2| = \sqrt{1^2 + 1^2} = \sqrt{2} = 1.41 \text{ approx.}$$

$$|z_1 + z_2| = \sqrt{3^2 + 4^2} = \sqrt{25} = 5$$

$$|z_1| + |z_2| = 3.6 + 1.41 = 5.01$$

$$\therefore |z_1 + z_2| \leq |z_1| + |z_2|$$

1.7.2 Subtraction of the Complex Numbers

Let two complex numbers $z_1 = a + bi$ and $z_2 = c + di$ be represented by the points (a, b) and (c, d) respectively.

$$\begin{aligned} \therefore (z_1) - (z_2) &= (a + bi) - (c + di) \\ &= (a - c) + (b - d)i \end{aligned}$$

which represents a point (a - c, b - d)

\therefore The difference i.e. $z_1 - z_2$ is represented by the point (a - c, b - d).

Thus, to subtract a complex number from another, we subtract corresponding real and imaginary parts separately.

MODULE - I
Algebra

Notes

Example 1.19 Find $z_1 - z_2$ in each of following if:

(a) $z_1 = 3 - 4i, \quad z_2 = -3 + 7i$

(b) $z_1 = -4 + 7i \quad z_2 = -4 - 5i$

Solution: (a) $z_1 - z_2 = (3 - 4i) - (-3 + 7i)$

$$= (3 - 4i) + (3 - 7i)$$

$$= (3 + 3) + (-4 - 7)i$$

$$= 6 + (-11i) = 6 - 11i$$

(b) $z_1 - z_2 = (-4 + 7i) - (-4 - 5i)$

$$= (-4 + 7i) + (4 + 5i)i$$

$$= (-4 + 4) + (7 + 5)i$$

$$= 0 + 12i = 12i$$

Example 1.20 What should be added to i to obtain 5 ?**Solution:** Let $z = a + bi$ be added to i to obtain $5 + 4i$

$$\therefore i + (a + bi) = 5 + 4i$$

or, $a + (b + 1)i = 5 + 4i$

Equating real and imaginary parts, we have

$$a = 5 \text{ and } b + 1 = 4 \text{ or } b = 3$$

$$\therefore z = 5 + 3i \text{ is to be added to } i \text{ to obtain } 5 + 4i$$

1.8 PROPERTIES: WITH RESPECT TO ADDITION OF COMPLEX NUMBERS.**1. Closure :** The sum of two complex numbers will always be a complex number.

Let $z_1 = a_1 + b_1i$ and $z_2 = a_2 + b_2i, \quad a_1, b_1, a_2, b_2 \in \mathbb{R}.$

Now, $z_1 + z_2 = (a_1 + a_2) + (b_1 + b_2)i$ which is again a complex number.

This proves the closure property of complex numbers.

Thus, $(1 + i) + (2 + 3i) = (1 + 2) + (1 + 3)i = 3 + 4i$, which is again a complex number.Similarly, the difference of two complex numbers will always be a complex number. For example, $(2 + 4i) - (1 - 4i) = (2 - 1) + \{4 - (-4)\}i = 1 + 8i$, which is again a complex number.



2. Commutative : If z_1 and z_2 are two complex numbers then

$$z_1 + z_2 = z_2 + z_1$$

Let $z_1 = a_1 + b_1i$ and $z_2 = a_2 + b_2i$

$$\begin{aligned} \text{Now } z_1 + z_2 &= (a_1 + b_1i) + (a_2 + b_2i) \\ &= (a_1 + a_2) + (b_1 + b_2)i \\ &= (a_2 + a_1) + (b_2 + b_1)i \quad [\text{commutative property of real numbers}] \\ &= (a_2 + b_2i) + (a_1 + b_1i) \\ &= z_2 + z_1 \end{aligned}$$

i.e. $z_1 + z_2 = z_2 + z_1$

Hence, addition of complex numbers is commutative.

For example, if $z_1 = 8 + 7i$ and $z_2 = 9 - 3i$ then

$$\begin{aligned} z_1 + z_2 &= (8 + 7i) + (9 - 3i) \quad \text{and} \quad z_2 + z_1 = (9 - 3i) + (8 + 7i) \\ &= (8 + 9) + (7 - 3)i \quad \text{and} \quad = (9 + 8) + (-3 + 7)i \end{aligned}$$

or $z_1 + z_2 = 17 + 4i \quad \text{and} \quad z_2 + z_1 = 17 + 4i$

We get, $z_1 + z_2 = z_2 + z_1$

$$\begin{aligned} \text{Now, } z_1 - z_2 &= (a_1 + b_1i) - (a_2 + b_2i) \\ &= (a_1 - a_2) + (b_1 - b_2)i \\ \text{and } z_2 - z_1 &= (a_2 + b_2i) - (a_1 + b_1i) \\ &= (a_2 - a_1) + (b_2 - b_1)i \\ &= -(a_1 - a_2) - (b_1 - b_2)i \\ &= -(a_1 + b_1i) + (a_2 + b_2i) \end{aligned}$$

$$\therefore z_1 - z_2 \neq z_2 - z_1$$

Hence, subtraction of complex numbers is not commutative.

For example, if $z_1 = 8 + 7i$ and $z_2 = 9 - 3i$ then

$$\begin{aligned} z_1 - z_2 &= (8 + 7i) - (9 - 3i) \quad \text{and} \quad z_2 - z_1 = (9 - 3i) - (8 + 7i) \\ &= (8 - 9) + (7 + 3)i \quad \text{and} \quad = (9 - 8) + (-3 - 7)i \end{aligned}$$

or $z_1 - z_2 = -1 + 10i \quad \text{and} \quad z_2 - z_1 = 1 - 10i$

$$z_1 - z_2 \neq z_2 - z_1$$

MODULE - I
Algebra

Notes

3. Associative

If $z_1 = a_1 + b_1i$, $z_2 = a_2 + b_2i$ and $z_3 = a_3 + b_3i$ are three complex numbers, then

$$z_1 + (z_2 + z_3) = (z_1 + z_2) + z_3$$

Now $z_1 + (z_2 + z_3)$

$$= (a_1 + b_1i) + \{(a_2 + b_2i) + (a_3 + b_3i)\}$$

$$= (a_1 + b_1i) + \{(a_2 + a_3) + (b_2 + b_3)i\}$$

$$= \{a_1 + (a_2 + a_3)\} + \{b_1 + (b_2 + b_3)\}i$$

$$= \{(a_1 + a_2) + (b_1 + b_2)i\} + (a_3 + b_3i)$$

$$= \{(a_1 + b_1i) + (a_2 + b_2i)\} + (a_3 + b_3i)$$

$$= (z_1 + z_2) + z_3$$

Hence, the associativity property holds good in the case of addition of complex numbers.

For example, if $z_1 = 2 + 3i$, $z_2 = 3i$ and $z_3 = 1 - 2i$, then

$$z_1 + (z_2 + z_3) = (2 + 3i) + \{(3i) + (1 - 2i)\}$$

$$= (2 + 3i) + (1 + i)$$

$$= (3 + 4i)$$

and $(z_1 + z_2) + z_3 = \{(2 + 3i) + (3i)\} + (1 - 2i)$

$$= (2 + 6i) + (1 - 2i)$$

$$= (3 + 4i)$$

Thus, $z_1 + (z_2 + z_3) = (z_1 + z_2) + z_3$

The equality of two sums is the consequence of the associative property of addition of complex numbers.

Like commutativity, it can be shown that associativity also does not hold good in the case of subtraction.

4. Existence of Additive Identity

If $x + yi$ be a complex number, then there exists a complex number $(0 + 0i)$

such that $(x + yi) + (0 + 0i) = x + yi$.

Let $z_2 = x + yi$ be the additive identity of $z_1 = 2 + 3i$ then

$$z_1 + z_2 = z_1$$

i.e. $(2 + 3i) + (x + yi) = 2 + 3i$



or $(2 + x) + (3 + y)i = 2 + 3i$

or $(2 + x) = 2$ and $3 + y = 3$

or $x = 0$ and $y = 0$

i.e. $z_2 = x + yi = 0 + 0i$ is the additive identity.

i.e. if $z = a + bi$ is any complex number, then

$$(a + bi) + (0 + 0i) = a + bi$$

i.e. $(0 + 0i)$ is the additive identity.

$$z_1 - z_2 = (2 + 3i) - (0 + 0i)$$

$$= (2 - 0) + (3 - 0)i$$

$$= 2 + 3i = z_1$$

$\therefore z_2 = 0 + 0i$ is the identity w.r.t. subtraction also.

as $(a + bi) - (0 + 0i) = a + bi$

5. Existence of Additive Inverse

For every complex number $a + bi$ there exists a unique complex number $-a - bi$ such that $(a + bi) + (-a - bi) = 0 + 0i$

e.g. Let $z_1 = 4 + 5i$ and $z_2 = x + yi$ be the additive inverse of z_1

Then, $z_1 + z_2 = 0$

or $(4 + 5i) + (x + yi) = 0 + 0i$

or $(4 + x) + (5 + y)i = 0 + 0i$

or $4 + x = 0$ and $5 + y = 0$

or $x = -4$ and $y = -5$

Thus, $z_2 = -4 - 5i$ is the additive inverse of $z_1 = 4 + 5i$

In general, additive inverse of a complex number is obtained by changing the signs of real and imaginary parts.

Consider $z_1 - z_2 = 0$

or $(4 + 5i) - (x + yi) = 0 + 0i$

or $(4 - x) + (5 - y)i = 0 + 0i$

or $4 - x = 0$ and $5 - y = 0$

MODULE - I
Algebra

Notes

or $x = 4$ and $y = 5$ i.e. $z_1 - z_2 = 0$ gives $z_2 = 4 + 5i$

Thus, in subtraction, the number itself is the inverse.

i.e. $(a + bi) - (a + bi) = 0 + 0i$ or 0 **CHECK YOUR PROGRESS 1.3****1. Simplify:**

(a) $(\sqrt{2} + \sqrt{5}i) + (\sqrt{5} - \sqrt{2}i)$

(b) $\frac{2+i}{3} + \frac{2-i}{6}$

(c) $(1 + i) - (1 - 6i)$

(d) $(\sqrt{2} - \sqrt{3}i) - (-2 - 7i)$

2. If $z_1 = (5 + i)$ and $z_2 = (6 + 2i)$, then:

(a) find $z_1 + z_2$

(b) find $z_2 + z_1$

(c) Is $z_1 + z_2 = z_2 + z_1$?

(d) find $z_1 - z_2$

(e) find $z_2 - z_1$

(f) Is $z_1 - z_2 = z_2 - z_1$?

3. If $z_1 = (1 + i)$, $z_2 = (1 - i)$ and $z_3 = (2 + 3i)$, then:

(a) find $z_1 + (z_2 + z_3)$

(b) find $(z_1 + z_2) + z_3$

(c) Is $z_1 + (z_2 + z_3) = (z_1 + z_2) + z_3$?

(d) find $z_1 - (z_2 - z_3)$

(e) find $(z_1 - z_2) - z_3$

(f) Is $z_1 - (z_2 - z_3) = (z_1 - z_2) - z_3$?

4. Find the additive inverse of the following:

(a) $12 - 7i$

(b) $4 - 3i$

5. What should be added to $(-15 + 4i)$ to obtain $(3 - 2i)$?6. Show that $\overline{\{(3 + 7i) - (5 + 2i)\}} = \overline{(3 + 7i)} - \overline{(5 + 2i)}$ **1.9 ARGUMENT OF A COMPLEX NUMBER**

Let $P(a, b)$ represent the complex number $z = a + bi$, $a \in \mathbb{R}$, $b \in \mathbb{R}$, and OP makes an angle θ with the positive direction of x -axis.

Draw $PM \perp OX$

Let $OP = r$

In right $\triangle OMP$

$$OM = a$$

$$MP = b$$

$$\therefore r \cos \theta = a$$

$$r \sin \theta = b$$




Fig. 1.11

Then $z = a + bi$ can be written as $z = r(\cos \theta + i \sin \theta)$... (i)

$$\text{where } r = \sqrt{a^2 + b^2} \text{ and } \tan \theta = \frac{b}{a}$$

$$\text{or } \theta = \tan^{-1} \left(\frac{b}{a} \right)$$

This is known as the polar form of the complex number z , and r and θ are respectively called the modulus and argument of the complex number.

1.10 MULTIPLICATION OF TWO COMPLEX NUMBERS

Two complex numbers can be multiplied by the usual laws of addition and multiplication as is done in the case of numbers.

Let $z_1 = (a + bi)$ and $z_2 = (c + di)$ then,

$$z_1 \cdot z_2 = (a + bi) \cdot (c + di)$$

$$= a(c + di) + bi(c + di)$$

$$\text{or } = ac + adi + bci + bdi^2$$

$$\text{or } = (ac - bd) + (ad + bc)i. \quad [\text{since } i^2 = -1]$$

If $(a + bi)$ and $(c + di)$ are two complex numbers, their product is defined as the complex number $(ac - bd) + (ad + bc)i$

Example 1.21 Evaluate:

$$(i) \quad (1 + 2i)(1 - 3i), \quad (ii) \quad (\sqrt{3} + i)(\sqrt{3} - i) \quad (iii) \quad (3 - 2i)^2$$

Solution:

$$(i) \quad (1 + 2i)(1 - 3i) = \{1 - (-6)\} + (-3 + 2)i \\ = 7 - i$$



MODULE - I
Algebra

Notes

$$\begin{aligned} \text{(ii)} \quad (\sqrt{3} + i)(\sqrt{3} - i) &= \{3 - (-1)\} + (-\sqrt{3} + \sqrt{3})i \\ &= 4 + 0i \end{aligned}$$

$$\begin{aligned} \text{(iii)} \quad (3 - 2i)^2 &= (3 - 2i)(3 - 2i) \\ &= (9 - 4) + (-6 - 6)i \\ &= 5 - 12i \end{aligned}$$

1.10.1 Properties of Multiplication

$$|z_1 \cdot z_2| = |z_1| \cdot |z_2|$$

Let $z_1 = r_1(\cos\theta_1 + i \sin\theta_1)$ and $z_2 = r_2(\cos\theta_2 + i \sin\theta_2)$

$$\therefore |z_1| = r_1 \sqrt{\cos^2 \theta_1 + \sin^2 \theta_1} = r_1$$

Similarly, $|z_2| = r_2$.

$$\begin{aligned} \text{Now,} \quad z_1 z_2 &= r_1(\cos\theta_1 + i \sin\theta_1) \cdot r_2(\cos\theta_2 + i \sin\theta_2) \\ &= r_1 r_2 [(\cos\theta_1 \cos\theta_2 - \sin\theta_1 \sin\theta_2) + (\cos\theta_1 \sin\theta_2 + \sin\theta_1 \cos\theta_2)i] \\ &= r_1 r_2 [\cos(\theta_1 + \theta_2) + i \sin(\theta_1 + \theta_2)] \\ &[\text{Since } \cos(\theta_1 + \theta_2) = \cos\theta_1 \cos\theta_2 - \sin\theta_1 \sin\theta_2 \text{ and} \\ &\quad \sin(\theta_1 + \theta_2) = \sin\theta_1 \cos\theta_2 + \cos\theta_1 \sin\theta_2] \end{aligned}$$

$$|z_1 \cdot z_2| = r_1 r_2 \sqrt{\cos^2(\theta_1 + \theta_2) + \sin^2(\theta_1 + \theta_2)} = r_1 r_2$$

$$\therefore |z_1 \cdot z_2| = r_1 r_2 = |z_1| \cdot |z_2|$$

and argument of $z_1 z_2 = \theta_1 + \theta_2 = \arg(z_1) + \arg(z_2)$

Example 1.22 Find the modulus of the complex number $(1 + i)(4 - 3i)$

Solution: Let $z = (1 + i)(4 - 3i)$

$$\begin{aligned} \text{then} \quad |z| &= |(1 + i)(4 - 3i)| \\ &= |(1 + i)| \cdot |(4 - 3i)| \quad (\text{since } |z_1 z_2| = |z_1| \cdot |z_2|) \end{aligned}$$

$$\text{But } |1 + i| = \sqrt{1^2 + 1^2} = \sqrt{2}$$

$$|4 - 3i| = \sqrt{4^2 + (-3)^2} = 5$$

$$\therefore |z| = \sqrt{2} \cdot 5 = 5\sqrt{2}$$



1.11 DIVISION OF TWO COMPLEX NUMBERS

Division of complex numbers involves multiplying both numerator and denominator with the conjugate of the denominator. We will explain it through an example.

Let $z_1 = a + bi$ and $z_2 = c + di$ then.

$$\frac{z_1}{z_2} = \frac{a + bi}{c + di} \quad (c + di \neq 0)$$

$$\frac{a + bi}{c + di} = \frac{(a + bi)(c - di)}{(c + di)(c - di)}$$

(multiplying numerator and denominator with the conjugate of the denominator)

$$= \frac{(ac + bd) + (bc - ad)i}{c^2 + d^2}$$

Thus,
$$\frac{a + bi}{c + di} = \frac{ac + bd}{c^2 + d^2} + \frac{bc - ad}{c^2 + d^2}i$$

Example 1.23 Divide $3 + i$ by $4 - 2i$

Solution:
$$\frac{3 + i}{4 - 2i} = \frac{(3 + i)(4 + 2i)}{(4 - 2i)(4 + 2i)}$$

Multiplying numerator and denominator by the conjugate of $(4 - 2i)$ we get

$$\begin{aligned} &= \frac{10 + 10i}{20} \\ &= \frac{1}{2} + \frac{1}{2}i \end{aligned}$$

Thus,
$$\frac{3 + i}{4 - 2i} = \frac{1}{2} + \frac{1}{2}i$$

1.11.1 Properties of Division

$$\left| \frac{z_1}{z_2} \right| = \frac{|z_1|}{|z_2|}$$

MODULE - I
Algebra

Notes

Proof: $z_1 = r_1(\cos\theta_1 + i \sin\theta_1)$

$$z_2 = r_2(\cos\theta_2 + i \sin\theta_2)$$

$$|z_1| = r_1 \sqrt{\cos^2 \theta_1 + \sin^2 \theta_1} = r_1$$

Similarly, $|z_2| = r_2$ and $\arg(z_1) = \theta_1$ and $\arg(z_2) = \theta_2$

$$\text{Then, } \frac{z_1}{z_2} = \frac{r_1(\cos\theta_1 + i \sin\theta_1)}{r_2(\cos\theta_2 + i \sin\theta_2)}$$

$$= \frac{r_1(\cos\theta_1 + i \sin\theta_1)(\cos\theta_2 - i \sin\theta_2)}{r_2(\cos\theta_2 + i \sin\theta_2)(\cos\theta_2 - i \sin\theta_2)}$$

$$= \frac{r_1}{r_2} \frac{(\cos\theta_1 \cos\theta_2 - i \cos\theta_1 \sin\theta_2 + i \sin\theta_1 \cos\theta_2 + \sin\theta_1 \sin\theta_2)}{(\cos^2 \theta_2 + \sin^2 \theta_2)}$$

$$= \frac{r_1}{r_2} [(\cos\theta_1 \cos\theta_2 + \sin\theta_1 \sin\theta_2) + i(\sin\theta_1 \cos\theta_2 - \cos\theta_1 \sin\theta_2)]$$

$$= \frac{r_1}{r_2} [\cos(\theta_1 - \theta_2) + i \sin(\theta_1 - \theta_2)]$$

$$\text{Thus, } \left| \frac{z_1}{z_2} \right| = \frac{r_1}{r_2} \sqrt{\cos^2(\theta_1 - \theta_2) + \sin^2(\theta_1 - \theta_2)} = \frac{r_1}{r_2}$$

$$\therefore \text{Argument of } \left| \frac{z_1}{z_2} \right| = \frac{r_1}{r_2} = \theta_1 - \theta_2$$

Example 1.24 Find the modulus of the complex number

$$\frac{2 + i}{3 - i}$$

$$\text{Solution : Let } z = \frac{2 + i}{3 - i}$$



$$\begin{aligned}\therefore |z| &= \left| \frac{2+i}{3-i} \right| = \frac{|2+i|}{|3-i|} \left(\sin ce \left| \frac{z_1}{z_2} \right| = \frac{|z_1|}{|z_2|} \right) \\ &= \frac{\sqrt{2^2+1^2}}{\sqrt{3^2+(-1)^2}} = \frac{\sqrt{5}}{\sqrt{10}} = \frac{1}{\sqrt{2}}\end{aligned}$$

$$\therefore |z| = \frac{1}{\sqrt{2}}$$

1.12 PROPERTIES OF MULTIPLICATION OF TWO COMPLEX NUMBERS

1. Closure

If $z_1 = a + bi$ and $z_2 = c + di$ be two complex numbers then their product $z_1 z_2$ is also a complex number.

2. Cummutative

If $z_1 = a + bi$ and $z_2 = c + di$ be two complex numbers then $z_1 z_2 = z_2 z_1$.

For example, let $z_1 = 3 + 4i$ and $z_2 = 1 - i$

$$\begin{aligned}\text{then } z_1 z_2 &= (3 + 4i)(1 - i) \\ &= 3(1 - i) + 4i(1 - i) \\ &= 3 - 3i + 4i - 4i^2 \\ &= 3 - 3i + 4i - 4(-1) \\ &= 3 + i + 4 = 7 + i\end{aligned}$$

$$\begin{aligned}\text{Again, } z_2 z_1 &= (1 - i)(3 + 4i) \\ &= (3 + 4i) - i(3 + 4i) \\ &= 3 + 4i - 3i - 4i^2 \\ &= 3 + i + 4 = 7 + i\end{aligned}$$

$$\therefore z_1 z_2 = z_2 z_1$$

3. Associativity

If $z_1 = (a + bi)$, $z_2 = c + di$ and $z_3 = (e + fi)$ then

$$z_1(z_2 \cdot z_3) = (z_1 \cdot z_2) \cdot z_3$$

MODULE - I
Algebra

Notes

Let us verify it with an example :

If $z_1 = (1 + i)$, $z_2 = (2 + i)$ and $z_3 = (3 + i)$ then

$$\begin{aligned} z_1(z_2 \cdot z_3) &= (1 + i)\{(2 + i)(3 + i)\} \\ &= (1 + i)\{(6 - 1) + (3 + 2)i\} \\ &= (1 + i)(5 + 5i) \\ &= (5 - 5) + (5 + 5)i \\ &= 0 + 10i = 10i \end{aligned}$$

$$\begin{aligned} \text{and } (z_1 \cdot z_2)z_3 &= \{(1 + i)(2 + i)\}(3 + i) \\ &= \{(2 - 1) + (1 + 2)i\}(3 + i) \\ &= (1 + 3i)(3 + i) \\ &= (3 - 3) + (1 + 9)i \\ &= 0 + 10i = 10i \end{aligned}$$

$$\therefore z_1(z_2 \cdot z_3) = (z_1 \cdot z_2)z_3$$

4. Existence of Multiplicative Identity: For every non-zero complex number $z_1 = a + bi$ there exists a unique complex number $(1 + 0i)$ such that

$$(a + bi) \cdot (1 + 0i) = (1 + 0i)(a + bi) = a + bi$$

Let $z_1 = x + yi$ be the multiplicative identity of $z = a + bi$

$$\text{Then } z \cdot z_1 = z.$$

$$\text{i.e. } (a + bi)(x + yi) = a + bi$$

$$\text{or } (ax - by) + (ay + bx)i = a + bi$$

$$\text{or } ax - by = a \text{ and } ay + bx = b$$

$$\text{pr } x = 1 \text{ and } y = 0$$

$$\text{i.e. } z_1 = x + yi = 1 + 0i \text{ is the multiplicative identity.}$$

The complex number $1 + 0i$ is the identity for multiplication.

Let us verify it with an example:

If $z = 2 + 3i$ then

$$\begin{aligned} z \cdot (1 + 0i) &= (2 + 3i)(1 + 0i) \\ &= (2 - 0) + (3 + 0)i \\ &= 2 + 3i \end{aligned}$$



5. Existence of Multiplicative inverse: Multiplicative inverse is a complex number that when multiplied to a given non-zero complex number yields one. In other words, for every non-zero complex number $z = a + bi$, there exists a unique complex number $(x + yi)$ such that their product is $(1 + 0i)$.

$$\text{i.e. } (a + bi)(x + yi) = 1 + 0i$$

$$\text{or } (ax - by) + (bx + ay)i = 1 + 0i$$

Equating real and imaginary parts, we have

$$ax - by = 1 \text{ and } bx + ay = 0$$

By cross multiplication

$$\frac{x}{a} = \frac{y}{-b} = \frac{1}{a^2 + b^2}$$

$$\Rightarrow x = \frac{a}{a^2 + b^2} = \frac{\operatorname{Re}(z)}{|z|^2} \text{ and } y = \frac{-b}{a^2 + b^2} = -\frac{\operatorname{Im}(z)}{|z|^2}$$

Thus, the multiplicative inverse of a non-zero complex number $z = (a + bi)$ is

$$x + yi = \left(\frac{\operatorname{Re}(z)}{|z|^2} - \frac{\operatorname{Im}(z)}{|z|^2}i \right) = \frac{\bar{z}}{|z|^2}$$

Example 1.25 Find the multiplication inverse of $2 - 4i$.

Solution: Let $z = 2 - 4i$

We have, $\bar{z} = 2 + 4i$ and $|z|^2 = 2^2 + (-4)^2 = 20$

\therefore Required multiplicative inverse is

$$\frac{\bar{z}}{|z|^2} = \frac{2 + 4i}{20} = \frac{1}{10} + \frac{1}{5}i$$

Verification:

If $\frac{1}{10} + \frac{1}{5}i$ be the multiplicative inverse of $2 - 4i$, their product must be equal to $1 + 0i$

$$\begin{aligned} \text{We have, } (2 - 4i)\left(\frac{1}{10} + \frac{1}{5}i\right) &= \left(\frac{2}{10} + \frac{4}{5}\right) + \left(\frac{2}{5} + \frac{4}{10}\right)i \\ &= 1 + 0i \text{ which is true.} \end{aligned}$$

MODULE - I
Algebra

Notes

6. Distributive Property of Multiplication over AdditionLet $z_1 = a_1 + b_1i$, $z_2 = a_2 + b_2i$ and $z_3 = a_3 + b_3i$ Then $z_1(z_2 + z_3) = z_1z_2 + z_1z_3$

Let us verify it with an example:

If $z_1 = 3 - 2i$, $z_2 = -1 + 4i$ and $z_3 = -3 - i$ then

$$\begin{aligned}
 z_1(z_2 + z_3) &= (3 - 2i)\{(-1 + 4i) + (-3 - i)\} \\
 &= (3 - 2i)(-1 + 4i - 3 - i) \\
 &= (3 - 2i)(-4 + 3i) \\
 &= (-12 + 6) + (9 + 8)i \\
 &= -6 + 17i
 \end{aligned}$$

$$\begin{aligned}
 \text{and } z_1z_2 &= (3 - 2i)(-1 + 4i) \\
 &= (-3 + 8) + (12 + 2)i \\
 &= 5 + 14i
 \end{aligned}$$

$$\begin{aligned}
 \text{Again } z_1z_3 &= (3 - 2i)(-3 - i) \\
 &= (-9 - 2) + (-3 + 6)i \\
 &= -11 + 3i
 \end{aligned}$$

$$\begin{aligned}
 \text{Now } z_1z_2 + z_1z_3 &= (5 + 14i) + (-11 + 3i) \\
 &= -6 + 17i
 \end{aligned}$$

$$\therefore z_1(z_2 + z_3) = z_1z_2 + z_1z_3$$

**CHECK YOUR PROGRESS 1.4**

1. Simplify each of the following:

(a) $(1 + 2i)(\sqrt{2} - i)$

(b) $(\sqrt{2} + i)^2$

(c) $(3 + i)(1 - i)(-1 + i)$

(d) $(2 + 3i) \div (1 - 2i)$

(e) $(1 + 2i) \div (1 + i)$

(f) $(1 + 0i) \div (3 + 7i)$

2. Compute multiplicative inverse of each of the following complex numbers:

(a) $3 - 4i$

(b) $\sqrt{3} + 7i$

(c) $\frac{3 + 5i}{2 - 3i}$

3. If $z_1 = 4 + 3i$, $z_2 = 3 - 2i$ and $z_3 = i + 5$, verify that $z_1(z_2 + z_3) = z_1z_2 + z_2z_3$.
4. If $z_1 = 2 + i$, $z_2 = -2 + i$ and $z_3 = 2 - i$ then verify that $(z_1 \cdot z_2)z_3 = z_1(z_2 \cdot z_3)$



LET US SUM UP

- $z = a + bi$ is a complex number in the standard form where $a, b \in \mathbb{R}$ and $i = \sqrt{-1}$.
- Any higher powers of 'i' can be expressed in terms of one of the four values $i, -1, -i, 1$.
- Conjugate of a complex number $z = a + bi$ is $a - bi$ and is denoted by \bar{z} .
- Modulus of a complex number $z = a + bi$ is $\sqrt{a^2 + b^2}$ i.e. $|z| = |a + bi| = \sqrt{a^2 + b^2}$

$$(a) |z| = 0 \Leftrightarrow z = 0 \qquad (b) |z| = |\bar{z}| \qquad (c) |z_1 + z_2| \leq |z_1| + |z_2|$$
- $z = r(\cos\theta + i\sin\theta)$ represents the polar form of a complex number $z = a + bi$ where $r = \sqrt{a^2 + b^2}$ is modulus and $\theta = \tan^{-1}\left(\frac{b}{a}\right)$ is its argument.
- Multiplicative inverse of a complex number $z = a + bi$ is $\frac{\bar{z}}{|z|^2}$



SUPPORTIVE WEB SITES

<http://www.wikipedia.org>

<http://mathworld.wolfram.com>



TERMINAL EXERCISE

1. Find real and imaginary parts of each of the following:

(a) $2 + 7i$

(b) $3 + 0i$

(c) $-\frac{1}{2}$

(d) $5i$

(e) $\frac{1}{2 + 3i}$



MODULE - I
Algebra

Notes

2. Simplify each of the following:

(a) $\sqrt{-3} \cdot \sqrt{-27}$

(b) $\sqrt{-3} \sqrt{-4} \sqrt{-72}$

(c) $3i^{15} - 5i^8 + 1$

3. Form complex number whose real and imaginary parts are given in the form of ordered pairs.

(a) $z(3, -5)$

(b) $z(0, -4)$

(c) $z(8, \pi)$

4. Find the conjugate of each of the following:

(a) $1 - 2i$

(b) $-1 - 2i$

(c) $6 - \sqrt{2}i$

(d) $4i$

(e) $-4i$

5. Find the modulus of each of the following:

(a) $1 - i$

(b) $3 + \pi i$

(c) $-\frac{3}{2}i$

(d) $-2 + \sqrt{3}i$

6. Express $7i^{17} - 6i^6 + 3i^3 - 2i^2 + 1$ in the form of $a + bi$.7. Find the values of x and y if:

(a) $(x - yi) + 7 - 2i = 9 - i$

(b) $2x + 3yi = 4 - 9i$

(c) $x - 3yi = 7 + 9i$

8. Simplify each of the following:

(a) $(3 + i) - (1 - i) + (-1 + i)$

(b) $\left(\frac{1}{7} + i\right) - \left(\frac{2}{7} - i\right) + \left(\frac{3}{7} - 2i\right)$

9. Write additive inverse and multiplicative inverse of each of the following:

(a) $3 - 7i$

(b) $11 - 2i$

(c) $\sqrt{3} + 2i$

(d) $1 - \sqrt{2}i$

(e) $\frac{1+5i}{1-i}$

10. Find the modulus of each of the following complex numbers:

(a) $\frac{1+i}{3-i}$

(b) $\frac{5+2i}{\sqrt{2} + \sqrt{3}i}$

(c) $(3 + 2i)(1 - i)$

(d) $(1 - 3i)(-2i^3 + i^2 + 3)$

11. For the following pairs of complex numbers verify that $|z_1 z_2| = |z_2| |z_1|$

(a) $z_1 = 3 - 2i$, $z_2 = 1 - 5i$

(b) $z_1 = 3 - \sqrt{7}i$, $z_2 = \sqrt{3} - i$

12. For the following pairs of complex numbers verify that $\left| \frac{z_1}{z_2} \right| = \frac{|z_1|}{|z_2|}$

(a) $z_1 = 1 + 3i$, $z_2 = 2 + 5i$

(b) $z_1 = -2 + 5i$, $z_2 = 3 - 4i$



MODULE - I
Algebra

Notes


**ANSWERS****CHECK YOUR PROGRESS 1.1**

- 1 (a) $3\sqrt{3}i$ (b) $-3i$ (c) $\sqrt{13}i$
 2. (a) $5 + 0i$ (b) $0 - 3i$ (c) $0 + 0i$
 3. $12 - 4i$


CHECK YOUR PROGRESS 1.2

- 1 (a) $2i$ (b) $-5 + 3i$ (c) $-\sqrt{2}$ (d) $3 + 4i$

2. (a)




(b)






Notes

(c)




(d)



MODULE - I
Algebra

Notes

(e)



3. (a) (i) 3 (ii) $\sqrt{10}$ (iii) $\sqrt{13}$ (iv) $\sqrt{21}$

CHECK YOUR PROGRESS 1.3

1. (a) $(\sqrt{2} + \sqrt{5}) + (\sqrt{5} - \sqrt{2})i$ (b) $\frac{1}{6}(6 + i)$
 (c) $7i$ (d) $\sqrt{2}(\sqrt{2} + 1) + (7 - \sqrt{3})$
2. (a) $11 + 3i$ (b) $11 + 3i$ (c) Yes
 (d) $-1 - i$ (e) $1 + i$ (f) No
3. (a) $4 + 3i$ (b) $4 + 3i$ (c) Yes
 (d) $2 + 5i$ (e) $-2 - i$ (f) No.
4. (a) $-12 + 7i$ (b) $-4 + 3i$
5. $18 - 6i$

**CHECK YOUR PROGRESS 1.4**

1. (a) $(\sqrt{2} + 2) + (2\sqrt{2} - 1)i$ (b) $1 + 2\sqrt{2}i$ (c) $-2 + 6i$
(d) $\frac{1}{\sqrt{5}}(-4 + 7i)$ (e) $\frac{1}{2}(3 + i)$ (f) $\frac{1}{58}(3 - 7i)$
2. (a) $\frac{1}{25}(3 + 4i)$ (b) $\frac{1}{52}(\sqrt{3} - 7i)$ (c) $\frac{1}{34}(-9 - 19i)$

TERMINAL EXERCISE

1. (a) 2, 7
(b) 3, 0
(c) $-\frac{1}{2}, 0$
(d) 0, 5
(e) $\frac{2}{\sqrt{13}}, -\frac{3}{\sqrt{13}}$
2. (a) -9
(b) $-12\sqrt{6}i$
(c) $-4 - 3i$
3. (a) $3 - 5i$
(b) $0 - 4i$
(c) $8 + \pi i$
4. (a) $1 + 2i$
(b) $-1 + 2i$
(c) $6 + \sqrt{2}i$
(d) $-4i$
(e) $4i$

MODULE - I
Algebra

Notes

5. (a) $\sqrt{2}$
 (b) $\sqrt{9 + \pi^2}$
 (c) $\frac{3}{2}$ (d) $\sqrt{7}$
6. $9 + 4i$
7. (a) $x = 2, y = -1$
 (b) $x = 2, y = -3$
 (c) $x = 7, y = -3$
8. (a) $1 + 3i$
 (b) $\frac{2}{7} + 0i$
9. (a) $-3 + 7i, \frac{1}{58}(3 + 7i)$
 (b) $-11 + 2i, \frac{1}{125}(-11 + 2i)$
 (c) $-\sqrt{3} - 2i, \frac{1}{7}(\sqrt{3} - 2i)$
 (d) $-1 + \sqrt{2}i, \frac{1}{3}(1 + \sqrt{2}i)$
 (e) $2 - 3i, \frac{1}{13}(2 + 3i)$
10. (a) $\frac{1}{\sqrt{5}}$ (b) $\frac{1}{5}\sqrt{145}$ (c) $\sqrt{26}$ (d) $4\sqrt{5}$