

# Study Material

Downloaded from Vedantu

## About Vedantu

Vedantu is India's largest **LIVE online teaching platform** with best teachers from across the country.

Vedantu offers Live Interactive Classes for **JEE, NEET, KVPY, NTSE, Olympiads, CBSE, ICSE, IGCSE, IB & State Boards** for Students Studying in **6-12th Grades** and Droppers.

FREE

LIVE ONLINE

### MASTER CLASSES

FREE Webinars by Expert Teachers



Register for **FREE**

## Awesome Master Teachers



**Anand Prakash**  
B.Tech, IIT Roorkee  
Co-Founder, Vedantu



**Pulkit Jain**  
B.Tech, IIT Roorkee  
Co-Founder, Vedantu



**Vamsi Krishna**  
B.Tech, IIT Bombay  
Co-Founder, Vedantu



“My mentor is approachable and **guides me in my future aspirations as well.**”

Student - **Ayushi**



“My son loves the sessions and **I can already see the change.**”

Parent - **Sreelatha**



**10,04,600+**  
Hours of LIVE Learning



**9,49,900+**  
Happy Students



**95%**  
Top Results

**95%** Students of Regular Tuitions on Vedantu scored above **90%** in exams!

Vedantu

## FREE MASTER CLASS SERIES

- ✓ For **Grades 6-12th** targeting **JEE, CBSE, ICSE** & much more
- ✓ **Free 60 Minutes Live Interactive** classes everyday
- ✓ Learn from the **Master Teachers** - India's best

Register for **FREE**

Limited Seats!

# Download Vedantu's App & Get



All Study Material  
with Solution





LIVE  
Doubt Solving



Daily  
LIVE Classes



FREE Tests and  
Reports



**DOWNLOAD THE APP**

**Question 1:**

In a cricket match, a batswoman hits a boundary 6 times out of 30 balls she plays. Find the probability that she did not hit a boundary.

**Solution 1:**

Number of times the batswoman hits a boundary = 6

Total number of balls played = 30

∴ Number of times that the batswoman does not hit a boundary =  $30 - 6 = 24$

$$\begin{aligned} P(\text{she does not hit a boundary}) &= \frac{\text{Number of times when she does not hit boundary}}{\text{Total number of balls played}} \\ &= \frac{24}{30} = \frac{4}{5} \end{aligned}$$

**Question 2:**

1500 families with 2 children were selected randomly, and the following data were recorded:

Number of girls in a family	2	1	0
Number of families	475	814	211

Compute the probability of a family, chosen at random, having

- (i) 2 girls
- (ii) 1 girl
- (iii) No girl

Also check whether the sum of these probabilities is 1.

**Solution 2:**

Total number of families =  $475 + 814 + 211 = 1500$

- (i) Number of families having 2 girls = 475

$$\begin{aligned} P_1(\text{a randomly chosen family has 2 girls}) &= \frac{\text{Number of families having 2 girls}}{\text{Total number of families}} \\ &= \frac{475}{1500} = \frac{19}{60} \end{aligned}$$

- (ii) Number of families having 1 girl = 814

# Vedantu

## Study Materials

[NCERT Solutions for Class 6 to 12 \(Math & Science\)](#)

[Revision Notes for Class 6 to 12 \(Math & Science\)](#)

[RD Sharma Solutions for Class 6 to 12 Mathematics](#)

[RS Aggarwal Solutions for Class 6, 7 & 10 Mathematics](#)

[Important Questions for Class 6 to 12 \(Math & Science\)](#)

[CBSE Sample Papers for Class 9, 10 & 12 \(Math & Science\)](#)

[Important Formula for Class 6 to 12 Math](#)

[CBSE Syllabus for Class 6 to 12](#)

[Lakhmir Singh Solutions for Class 9 & 10](#)

[Previous Year Question Paper](#)

[CBSE Class 12 Previous Year Question Paper](#)

[CBSE Class 10 Previous Year Question Paper](#)

[JEE Main & Advanced Question Paper](#)

[NEET Previous Year Question Paper](#)

$$P_2 \text{ (a randomly chosen family has 1 girl)} = \frac{\text{Number of families having 1 girl}}{\text{Total number of families}}$$

$$= \frac{814}{1500} = \frac{407}{750}$$

(iii) Number of families having no girl = 211

$$P_3 \text{ (a randomly chosen family has no girl)} = \frac{\text{Number of families having no girl}}{\text{Total number of families}}$$

$$= \frac{211}{1500}$$

$$\text{Sum of all these probabilities} = \frac{19}{60} + \frac{407}{750} + \frac{211}{1500}$$


$$= \frac{475 + 814 + 211}{1500}$$

$$= \frac{1500}{1500} = 1$$

Therefore, the sum of all these probabilities is 1.

### Question 3:

In a particular section of Class IX, 40 students were asked about the months of their birth and the following graph was prepared for the data so obtained:



Find the probability that a student of the class was born in August.

### Solution 3:

Number of students born in the month of August = 6

Total number of students = 40

$$P \text{ (Students born in the month of August)} = \frac{\text{Number of students born in August}}{\text{Total number of students}}$$

$$= \frac{6}{40} = \frac{3}{20}$$

#### Question 4:

Three coins are tossed simultaneously 200 times with the following frequencies of different outcomes:

Outcome	3 heads	2 heads	1 head	No head
Frequency	23	72	77	28

If the three coins are simultaneously tossed again, compute the probability of 2 heads coming up.

#### Solution 4:

Number of times 2 heads come up = 72

Total number of times the coins were tossed = 200

$$\begin{aligned}
 P(2 \text{ heads will come up}) &= \frac{\text{Number of times 2 heads come up}}{\text{Total number of times the coins were tossed}} \\
 &= \frac{72}{200} = \frac{9}{25}
 \end{aligned}$$

#### Question 5:

An organization selected 2400 families at random and surveyed them to determine a relationship between income level and the number of vehicles in a family. The information gathered is listed in the table below:

Monthly income (in Rs.)	Vehicles per family			
	0	1	2	Above 2
Less than 7000	10	160	25	0
7000 – 10000	0	305	27	2
10000 – 13000	1	535	29	1
13000 – 16000	2	469	59	25
16000 or more	1	579	82	88

Suppose a family is chosen, find the probability that the family chosen is

- earning Rs. 10000–13000 per month and owning exactly 2 vehicles.
- earning Rs. 16000 or more per month and owning exactly 1 vehicle.
- earning less than Rs. 7000 per month and does not own any vehicle.
- earning Rs. 13000–16000 per month and owning more than 2 vehicles.
- owning not more than 1 vehicle.

#### Solution 5:

Number of total families surveyed =  $10 + 160 + 25 + 0 + 0 + 305 + 27 + 2 + 1 + 535 + 29 + 1 + 2 + 469 + 59 + 25 + 1 + 579 + 82 + 88 = 2400$

(i) Number of families earning Rs. 10000–13000 per month and owning exactly 2 vehicles = 29

Hence, required probability,  $P = \frac{29}{2400}$

(ii) Number of families earning Rs. 16000 or more per month and owning exactly 1 vehicle = 579

Hence, required probability,  $P = \frac{579}{2400}$

(iii) Number of families earning less than Rs. 7000 per month and does not own any vehicle = 10

Hence, required probability,  $P = \frac{10}{2400} = \frac{1}{240}$

(iv) Number of families earning Rs. 13000–16000 per month and owning more than 2 vehicles = 25

Hence, required probability,  $P = \frac{25}{2400} = \frac{1}{96}$

(v) Number of families owning not more than 1 vehicle =  $10 + 160 + 0 + 305 + 1 + 535 + 2 + 469 + 1 + 579 = 2062$

Hence, required probability,  $P = \frac{2062}{2400} = \frac{1031}{1200}$

### Question 6:

A teacher wanted to analyse the performance of two sections of students in a mathematics test of 100 marks. Looking at their performances, she found that a few students got under 20 marks and a few got 70 marks or above. So she decided to group them into intervals of varying sizes as follows: 0–20, 20–30, ..., 60–70, 70–100. Then she formed the following table:

Marks	Number of student
0 – 20	7
20 – 30	10
30 – 40	10
40 – 50	20
50 – 60	20
60 – 70	15
70 – above	8
Total	90

(i) Find the probability that a student obtained less than 20% in the mathematics test.

(ii) Find the probability that a student obtained marks 60 or above.

### Solution 6:

Total number of students = 90

(i) Number of students getting less than 20 % marks in the test = 7



Hence, required probability,  $P = \frac{7}{90}$

(ii) Number of students obtaining marks 60 or above =  $15 + 8 = 23$

Hence, required probability,  $P = \frac{23}{90}$

### Question 7:

To know the opinion of the students about the subject statistics, a survey of 200 students was conducted. The data is recorded in the following table.

Opinion	Number of students
like	135
dislike	65

Find the probability that a student chosen at random

- (i) likes statistics
- (ii) does not like it

### Solution 7:

Total number of students =  $135 + 65 = 200$

(i) Number of students liking statistics = 135

$$P(\text{students liking statistics}) = \frac{135}{200} = \frac{27}{40}$$

(ii) Number of students who do not like statistics = 65

$$P(\text{students not liking statistics}) = \frac{65}{200} = \frac{13}{40}$$

### Question 8:

The distance (in km) of 40 engineers from their residence to their place of work were found as follows:

5	3	10	20	25	11	13	7	12	31
19	10	12	17	18	11	32	17	16	2
7	9	7	8	3	5	12	15	18	3
12	14	2	9	6	15	15	7	6	12

What is the empirical probability that an engineer lives:

- (i) less than 7 km from her place of work?
- (ii) more than or equal to 7 km from her place of work?
- (iii) within  $\frac{1}{2}$  km from her place of work?

### Solution 8:

(i) Total number of engineers = 40

Number of engineers living less than 7 km from their place of work = 9



Hence, required probability that an engineer lives less than 7 km from her place of work,  $P = \frac{9}{40}$

(ii) Number of engineers living more than or equal to 7 km from their place of work =  $40 - 9 = 31$

Hence, required probability that an engineer lives more than or equal to 7 km from her place of work,  $P = \frac{31}{40}$

(iii) Number of engineers living within  $\frac{1}{2}$  km from her place of work = 0

Hence, required probability that an engineer lives within  $\frac{1}{2}$  km from her place of work,  $P = 0$

### Question 11:

Eleven bags of wheat flour, each marked 5 kg, actually contained the following weights of flour (in kg): 4.97, 5.05, 5.08, 5.03, 5.00, 5.06, 5.08, 4.98, 5.04, 5.07, 5.00

Find the probability that any of these bags chosen at random contains more than 5 kg of flour.

### Solution 11 :

Number of total bags = 11

Number of bags containing more than 5 kg of flour = 7

Hence, required probability,  $P = \frac{7}{11}$

### Question 12:

The below frequency distribution table represents the concentration of sulphur dioxide in the air in parts per million of a certain city for 30 days. Using this table,

find the probability of the concentration of sulphur dioxide in the interval 0.12–0.16 on any of these days.

Concentration of $\text{SO}_2$ (in ppm)	Number of days (frequency)
0.00 – 0.04	4
0.04 – 0.08	9
0.08 – 0.12	9
0.12 – 0.16	2
0.16 – 0.20	4
0.20 – 0.24	2
Total	30

### Solution 12 :

Number days for which the concentration of sulphur dioxide was in the interval of 0.12–0.16 = 2

Total number of days = 30

Hence, required probability,  $P = \frac{2}{30} = \frac{1}{15}$

**Question 13:**

The below frequency distribution table represents the blood groups of 30 students of a class. Use this table to determine the probability that a student of this class, selected at random, has blood group AB.

Blood group	Number of students
A	9
B	6
AB	3
O	12
Total	30

**Solution 13:**

Number of students having blood group AB = 3

Total number of students = 30

Hence, required probability,  $P = \frac{3}{30} = \frac{1}{10}$

**Thank You**  
for downloading the PDF

**FREE** LIVE ONLINE

# MASTER CLASSES

FREE Webinars by Expert Teachers



Vedantu

## FREE MASTER CLASS SERIES

- ✓ For **Grades 6-12th** targeting **JEE, CBSE, ICSE** & much more
- ✓ **Free 60 Minutes Live Interactive** classes everyday
- ✓ Learn from the **Master Teachers** - India's best

Register for **FREE**

Limited Seats!