

Study Material

Downloaded from Vedantu

About Vedantu

Vedantu is India's largest **LIVE online teaching platform** with best teachers from across the country.

Vedantu offers Live Interactive Classes for **JEE, NEET, KVPY, NTSE, Olympiads, CBSE, ICSE, IGCSE, IB & State Boards** for Students Studying in **6-12th Grades** and Droppers.

FREE

LIVE ONLINE

MASTER CLASSES

FREE Webinars by Expert Teachers



Register for **FREE**

Awesome Master Teachers



Anand Prakash
B.Tech, IIT Roorkee
Co-Founder, Vedantu



Pulkit Jain
B.Tech, IIT Roorkee
Co-Founder, Vedantu



Vamsi Krishna
B.Tech, IIT Bombay
Co-Founder, Vedantu



“My mentor is approachable and **guides me in my future aspirations as well.**”

Student - **Ayushi**



“My son loves the sessions and **I can already see the change.**”

Parent - **Sreelatha**



10,04,600+
Hours of LIVE Learning



9,49,900+
Happy Students



95%
Top Results

95% Students of Regular Tuitions on Vedantu scored above **90%** in exams!

Vedantu

FREE MASTER CLASS SERIES

- ✓ For **Grades 6-12th** targeting **JEE, CBSE, ICSE** & much more
- ✓ **Free 60 Minutes Live Interactive** classes everyday
- ✓ Learn from the **Master Teachers** - India's best

Register for **FREE**

Limited Seats!

Download Vedantu's App & Get



All Study Material
with Solution




LIVE
Doubt Solving



Daily
LIVE Classes



FREE Tests and
Reports



GET IT ON
Google Play

DOWNLOAD THE APP

Question 1:

Use Euclid's division algorithm to find the HCF of:

- (i) 135 and 225 (ii) 196 and 38220 (iii) 867 and 255

Solution 1:**(i) 135 and 225**

Since $225 > 135$, we apply the division lemma to 225 and 135 to obtain

$$225 = 135 \times 1 + 90$$

Since remainder $90 \neq 0$, we apply the division lemma to 135 and 90 to obtain

$$135 = 90 \times 1 + 45$$

We consider the new divisor 90 and new remainder 45, and apply the division lemma to obtain

$$90 = 2 \times 45 + 0$$

Since the remainder is zero, the process stops.

Since the divisor at this stage is 45,

Therefore, the HCF of 135 and 225 is 45.

(ii) 196 and 38220

Since $38220 > 196$, we apply the division lemma to 38220 and 196 to obtain

$$38220 = 196 \times 195 + 0$$

Since the remainder is zero, the process stops.

Since the divisor at this stage is 196,

Therefore, HCF of 196 and 38220 is 196.

(iii) 867 and 255

Since $867 > 255$, we apply the division lemma to 867 and 255 to obtain

$$867 = 255 \times 3 + 102$$

Since remainder $102 \neq 0$, we apply the division lemma to 255 and 102 to obtain

$$255 = 102 \times 2 + 51$$

We consider the new divisor 102 and new remainder 51, and apply the division lemma to obtain

$$102 = 51 \times 2 + 0$$

Since the remainder is zero, the process stops.

Since the divisor at this stage is 51,

Therefore, HCF of 867 and 255 is 51.

Vedantu

Study Materials

[NCERT Solutions for Class 6 to 12 \(Math & Science\)](#)

[Revision Notes for Class 6 to 12 \(Math & Science\)](#)

[RD Sharma Solutions for Class 6 to 12 Mathematics](#)

[RS Aggarwal Solutions for Class 6, 7 & 10 Mathematics](#)

[Important Questions for Class 6 to 12 \(Math & Science\)](#)

[CBSE Sample Papers for Class 9, 10 & 12 \(Math & Science\)](#)

[Important Formula for Class 6 to 12 Math](#)

[CBSE Syllabus for Class 6 to 12](#)

[Lakhmir Singh Solutions for Class 9 & 10](#)

[Previous Year Question Paper](#)

[CBSE Class 12 Previous Year Question Paper](#)

[CBSE Class 10 Previous Year Question Paper](#)

[JEE Main & Advanced Question Paper](#)

[NEET Previous Year Question Paper](#)

Question 2:

Show that any positive odd integer is of the form, $6q + 1$ or $6q + 3$, or $6q + 5$, where q is some integer.

Solution 2:

Let a be any positive integer and $b = 6$. Then, by Euclid's algorithm, $a = 6q + r$ for some integer $q \geq 0$, and $r = 0, 1, 2, 3, 4, 5$ because $0 \leq r < 6$.

Therefore, $a = 6q$ or $6q + 1$ or $6q + 2$ or $6q + 3$ or $6q + 4$ or $6q + 5$
Also, $6q + 1 = 2 \times 3q + 1 = 2k_1 + 1$, where k_1 is a positive integer

$6q + 3 = (6q + 2) + 1 = 2(3q + 1) + 1 = 2k_2 + 1$, where k_2 is an Integer

$6q + 5 = (6q + 4) + 1 = 2(3q + 2) + 1 = 2k_3 + 1$, where k_3 is an Integer

Clearly, $6q + 1$, $6q + 3$, $6q + 5$ are of the form $2k + 1$, where k is an integer.

Therefore, $6q + 1$, $6q + 3$, $6q + 5$ are not exactly divisible by 2.

Hence, these expressions of numbers are odd numbers.

And therefore, any odd integer can be expressed in the form $6q + 1$,
or $6q + 3$,
or $6q + 5$

Question 3:

An army contingent of 616 members is to march behind an army band of 32 members in a parade. The two groups are to march in the same number of columns. What is the maximum number of columns in which they can march?

Solution 3:

HCF (616, 32) will give the maximum number of columns in which they can march.

We can use Euclid's algorithm to find the HCF.

$$616 = 32 \times 19 + 8$$

$$32 = 8 \times 4 + 0$$

The HCF (616, 32) is 8.

Therefore, they can march in 8 columns each.

Question 4:

Use Euclid's division lemma to show that the square of any positive integer is either of form $3m$ or $3m + 1$ for some integer m .

[**Hint:** Let x be any positive integer then it is of the form $3q$, $3q + 1$ or $3q + 2$. Now square each of these and show that they can be rewritten in the form $3m$ or $3m + 1$.]

Solution 4:

Let a be any positive integer and $b = 3$.

Then $a = 3q + r$ for some integer $q \geq 0$

And $r = 0, 1, 2$ because $0 \leq r < 3$

Therefore, $a = 3q$ or $3q + 1$ or $3q + 2$

Or,

$$a^2 = (3q)^2 \text{ or } (3q + 1)^2 \text{ or } (3q + 2)^2$$

$$a^2 = (9q^2) \text{ or } 9q^2 + 6q + 1 \text{ or } 9q^2 + 12q + 4$$

$$= 3 \times (3q^2) \text{ or } (3q^2 + 2q) + 1 \text{ or } 3(3q^2 + 4q + 1) + 1$$

$$= 3k_1 \text{ or } 3k_2 + 1 \text{ or } 3k_3 + 1$$

$$a^2 = (3q)^2 \text{ or } (3q+1)^2 \text{ or } (3q+2)^2$$

$$a^2 = (9q^2) \text{ or } (9q^2 + 6q + 1) \text{ or } (9q^2 + 12q + 4)$$

$$= 3 \times (3q^2) \text{ or } 3 \times (3q^2 + 2q) + 1 \text{ or } 3 \times (3q^2 + 4q + 1) + 1$$

$$= 3k_1 \text{ or } 3k_2 + 1 \text{ or } 3k_3 + 1$$

Where k_1 , k_2 , and k_3 are some positive integers

Hence, it can be said that the square of any positive integer is either of the form $3m$ or $3m + 1$.

Question 5:

Use Euclid's division lemma to show that the cube of any positive integer is of the form $9m$, $9m + 1$ or $9m + 8$.

Solution 5:

Let a be any positive integer and $b = 3$

$a = 3q + r$, where $q \geq 0$ and $0 \leq r < 3$

$\therefore a = 3q$ or $3q + 1$ or $3q + 2$

Therefore, every number can be represented as these three forms.

There are three cases.

Case 1: When $a = 3q$,

$$a^3 = (3q)^3 = 27q^3 = 9(3q^3) = 9m,$$

Where m is an integer such that $m = 3q^3$

Case 2: When $a = 3q + 1$,

$$a^3 = (3q + 1)^3$$

$$a^3 = 27q^3 + 27q^2 + 9q + 1$$

$$a^3 = 9(3q^3 + 3q^2 + q) + 1$$

$$a^3 = 9m + 1$$

Where m is an integer such that $m = (3q^3 + 3q^2 + q)$

Case 3: When $a = 3q + 2$,

$$a^3 = (3q + 2)^3$$

$$a^3 = 27q^3 + 54q^2 + 36q + 8$$

$$a^3 = 9(3q^3 + 6q^2 + 4q) + 8$$

$$a^3 = 9m + 8$$

Where m is an integer such that $m = (3q^3 + 6q^2 + 4q)$

Therefore, the cube of any positive integer is of the form $9m$, $9m + 1$, or $9m + 8$.

EXERCISE NO: 1.2

Question 1:

Express each number as product of its prime factors:

- (i) 140 (ii) 156 (iii) 3825 (iv) 5005 (v) 7429

Solution 1:

- (i) $140 = 2 \times 2 \times 5 \times 7 = 2^2 \times 5 \times 7$
(ii) $156 = 2 \times 2 \times 3 \times 13 = 2^2 \times 3 \times 13$
(iii) $3825 = 3 \times 3 \times 5 \times 5 \times 17 = 3^2 \times 5^2 \times 17$
(iv) $5005 = 5 \times 7 \times 11 \times 13$
(v) $7429 = 17 \times 19 \times 23$

Question 2:

Find the LCM and HCF of the following pairs of integers and verify that $\text{LCM} \times \text{HCF} = \text{product of the two numbers}$.

- (i) 26 and 91 (ii) 510 and 92 (iii) 336 and 54

Solution 2:

- (i) 26 and 91

$$26 = 2 \times 13$$

$$91 = 7 \times 13$$

$$\text{HCF} = 13$$

$$\text{LCM} = 2 \times 7 \times 13 = 182$$

$$\text{Product of the two numbers} = 26 \times 91 = 2366$$

$$\text{HCF} \times \text{LCM} = 13 \times 182 = 2366$$

$$\text{Hence, product of two numbers} = \text{HCF} \times \text{LCM}$$

- (ii) 510 and 92

$$510 = 2 \times 3 \times 5 \times 17$$

$$92 = 2 \times 2 \times 23$$

$$\text{HCF} = 2$$

$$\text{LCM} = 2 \times 2 \times 3 \times 5 \times 17 \times 23 = 23460$$

$$\text{Product of the two numbers} = 510 \times 92 = 46920$$

$$\text{HCF} \times \text{LCM} = 2 \times 23460$$

$$= 46920$$

$$\text{Hence, product of two numbers} = \text{HCF} \times \text{LCM}$$

- (iii) 336 and 54

$$336 = 2 \times 2 \times 2 \times 2 \times 3 \times 7$$

$$54 = 2 \times 3 \times 3 \times 3$$

$$54 = 2 \times 3 \times 3 \times 3$$

$$54 = 2 \times 3^3$$

$$\text{HCF} = 2 \times 3 = 6$$

$$\text{LCM} = 2^4 \times 3^3 \times 7 = 3024$$

$$\text{Product of the two numbers} = 336 \times 54 = 18144$$

$$\text{HCF} \times \text{LCM} = 6 \times 3024 = 18144$$

$$\text{Hence, product of two numbers} = \text{HCF} \times \text{LCM}$$

Question 3:

Find the LCM and HCF of the following integers by applying the prime factorisation method.

- (i) 12, 15 and 21 (ii) 17, 23 and 29 (iii) 8, 9 and 25

Solution 3:

- (i) 12, 15 and 21

$$12 = 2^2 \times 3$$

$$15 = 3 \times 5$$

$$21 = 3 \times 7$$

$$\text{HCF} = 3$$

$$\text{LCM} = 2^2 \times 3 \times 5 \times 7 = 420$$

- (ii) 17, 23 and 29

$$17 = 1 \times 17$$

$$23 = 1 \times 23$$

$$29 = 1 \times 29$$

$$\text{HCF} = 1$$

$$\text{LCM} = 17 \times 23 \times 29 = 11339$$

- (iii) 8, 9 and 25

$$8 = 2 \times 2 \times 2$$

$$9 = 3 \times 3$$

$$25 = 5 \times 5$$

$$\text{HCF} = 1$$

$$\text{LCM} = 2 \times 2 \times 2 \times 3 \times 3 \times 5 \times 5 = 1800$$

Question 4:

Given that $\text{HCF}(306, 657) = 9$, find $\text{LCM}(306, 657)$.

Solution 4:

$$\text{HCF}(306, 657) = 9$$

We know that, $\text{LCM} \times \text{HCF} = \text{Product of two numbers}$

$$\therefore \text{LCM} \times \text{HCF} = 306 \times 657$$

$$\text{LCM} = \frac{306 \times 657}{\text{HCF}} = \frac{306 \times 657}{9}$$

$$\text{LCM} = 22338$$

Question 5:

Check whether 6^n can end with the digit 0 for any natural number n .

Solution 5:

If any number ends with the digit 0, it should be divisible by 10 or in other words, it will also be divisible by 2 and 5 as $10 = 2 \times 5$

Prime factorisation of $6^n = (2 \times 3)^n$

It can be observed that 5 is not in the prime factorisation of $6n$.

Hence, for any value of n , 6^n will not be divisible by 5.

Therefore, 6^n cannot end with the digit 0 for any natural number n .

Question 6:

Explain why $7 \times 11 \times 13 + 13$ and $7 \times 6 \times 5 \times 4 \times 3 \times 2 \times 1 + 5$ are composite numbers.

Solution 6:

Numbers are of two types - prime and composite. A prime number can be divided by 1 and only itself, whereas a composite number have factors other than 1 and itself.

It can be observed that

$$\begin{aligned} 7 \times 11 \times 13 + 13 &= 13 \times (7 \times 11 + 1) = 13 \times (77 + 1) \\ &= 13 \times 78 \\ &= 13 \times 13 \times 6 \end{aligned}$$

The given expression has 6 and 13 as its factors. Therefore, it is a composite number.

$$\begin{aligned} 7 \times 6 \times 5 \times 4 \times 3 \times 2 \times 1 + 5 &= 5 \times (7 \times 6 \times 4 \times 3 \times 2 \times 1 + 1) \\ &= 5 \times (1008 + 1) \\ &= 5 \times 1009 \end{aligned}$$

1009 cannot be factorised further. Therefore, the given expression has 5 and 1009 as its factors. Hence, it is a composite number.

Question 7:

There is a circular path around a sports field. Sonia takes 18 minutes to drive one round of the field, while Ravi takes 12 minutes for the same. Suppose they both start at the same point and at the same time, and go in the same direction. After how many minutes will they meet again at the starting point?

Solution 7:

It can be observed that Ravi takes lesser time than Sonia for completing 1 round of the circular path. As they are going in the same direction, they will meet again at the same time when Ravi will have completed 1 round of that circular path with respect to Sonia. And the total time taken for completing this 1 round of circular path will be the LCM of time taken by Sonia and Ravi for completing 1 round of circular path respectively i.e., LCM of 18 minutes and 12 minutes.

$$18 = 2 \times 3 \times 3$$

$$\text{And, } 12 = 2 \times 2 \times 3$$

$$\text{LCM of 12 and 18} = 2 \times 2 \times 3 \times 3 = 36$$

Therefore, Ravi and Sonia will meet together at the starting point after 36 minutes.

EXERCISE NO: 1.3

Question 1:

Prove that $\sqrt{5}$ is irrational.

Solution 1:

Let $\sqrt{5}$ is a rational number. Therefore, we can find two integers a, b ($b \neq 0$)

such that $\sqrt{5} = \frac{a}{b}$

Let a and b have a common factor other than 1. Then we can divide them by the common factor, and assume that a and b are co-prime.

$$a = \sqrt{5}b$$

$$a^2 = 5b^2$$

Therefore, a^2 is divisible by 5 and it can be said that a is divisible by 5. Let $a = 5k$, where k is an integer

$$(5k)^2 = 5b^2$$

$b^2 = 5k^2$ This means that b^2 is divisible by 5 and hence, b is divisible by 5.

This implies that a and b have 5 as a common factor. And this is a contradiction to the fact that a and b are co-prime.

Hence, cannot be expressed as $\frac{p}{q}$ or it can be said that $\sqrt{5}$ is irrational.

Question 2:

Prove that $3 + 2\sqrt{5}$ is irrational.

Solution 2:

Let $3 + 2\sqrt{5}$ rational.

Therefore, we can find two integers a, b ($b \neq 0$) such that

$$3 + 2\sqrt{5} = \frac{a}{b}$$

$$2\sqrt{5} = \frac{a}{b} - 3$$

$$\sqrt{5} = \frac{1}{2} \left(\frac{a}{b} - 3 \right)$$

Since a and b are integers, $\frac{1}{2}\left(\frac{a}{b} - 3\right)$ will also be rational and

therefore, $\sqrt{5}$ is rational.

This contradicts the fact that $\sqrt{5}$ is irrational. Hence, our assumption that $3 + 2\sqrt{5}$ is rational is false. Therefore, $3 + 2\sqrt{5}$ is irrational.

Question 3:

Prove that the following are irrationals:

- (i) $\frac{1}{\sqrt{2}}$ (ii) $7\sqrt{5}$ (iii) $6 + \sqrt{2}$

Solution 3:

- (i) $\frac{1}{\sqrt{2}}$

Let $\frac{1}{\sqrt{2}}$ is rational

Therefore, we can find two integers a, b ($b \neq 0$) such that

$$\frac{1}{\sqrt{2}} = \frac{a}{b}$$

$$\sqrt{2} = \frac{b}{a}$$

$\frac{b}{a}$ is rational as a and b are integers. Therefore, $\sqrt{2}$ is rational which

contradicts to the fact that $\sqrt{2}$ is irrational.

Hence, our assumption is false and $\frac{1}{\sqrt{2}}$ is irrational.

- (ii) $7\sqrt{5}$

Let $7\sqrt{5}$ is rational.

Therefore, we can find two integers a, b ($b \neq 0$) such that $7\sqrt{5} = \frac{a}{b}$ for

some integers a and b

$$\therefore \sqrt{5} = \frac{a}{7b}$$

$\frac{a}{7b}$ is rational as a and b are integers.

Therefore, $\sqrt{5}$ should be rational.

This contradicts the fact that $\sqrt{5}$ is irrational. Therefore, our assumption that $7\sqrt{5}$ is rational is false. Hence, $7\sqrt{5}$ is irrational.

(iii) $6 + \sqrt{2}$

Let $6 + \sqrt{2}$ be rational.

Therefore, we can find two integers a, b ($b \neq 0$) such that

$$6 + \sqrt{2} = \frac{a}{b}$$

$$\sqrt{2} = \frac{a}{b} - 6$$

Since a and b are integers, $\frac{a}{b} - 6$ is also rational and hence, $\sqrt{2}$ should

be rational. This contradicts the fact that $\sqrt{2}$ is irrational. Therefore, our assumption is false and hence, $6 + \sqrt{2}$ is irrational.

EXERCISE NO: 1.4

Question 1:

Without actually performing the long division, state whether the following rational numbers will have a terminating decimal expansion or a non-terminating repeating decimal expansion:

(i) $\frac{13}{3125}$

(ii) $\frac{17}{8}$

(iii) $\frac{64}{455}$

(iv) $\frac{15}{1600}$

(v) $\frac{29}{343}$

(vi) $\frac{23}{2^3 5^2}$

(vii) $\frac{129}{2^2 5^7 7^5}$

(viii) $\frac{6}{15}$

(ix) $\frac{35}{50}$

(x) $\frac{77}{210}$

Solution 1:

(i) $\frac{13}{3125}$

$$3125 = 5^5$$

The denominator is of the form 5^m .

Hence, the decimal expansion of $\frac{13}{3125}$ is terminating.

(ii) $\frac{17}{8}$

$$8 = 2^3$$

The denominator is of the form 2^m .

Hence, the decimal expansion of $\frac{17}{8}$ is terminating.

(iii) $\frac{64}{455}$

$$455 = 5 \times 7 \times 13$$

Since the denominator is not in the form $2^m \times 5^n$, and it also contains 7 and 13 as its factors, its decimal expansion will be non-terminating repeating.

(iv) $\frac{15}{1600}$

$$1600 = 2^6 \times 5^2$$

The denominator is of the form $2^m \times 5^n$.

Hence, the decimal expansion of $\frac{15}{1600}$ is terminating.

$$(v) \frac{29}{343}$$

$$343 = 7^3$$

Since the denominator is not in the form $2^m \times 5^n$, and it has 7 as its factor, the decimal expansion of $\frac{29}{343}$ is non-terminating repeating.

$$(vi) \frac{23}{2^3 \times 5^2}$$

The denominator is of the form $2^m \times 5^n$.

Hence, the decimal expansion of $\frac{23}{2^3 \times 5^2}$ is terminating.

$$(vii) \frac{129}{2^2 \times 5^7 \times 7^5}$$

Since the denominator is not of the form $2^m \times 5^n$, and it also has 7 as its factor, the decimal expansion of $\frac{129}{2^2 \times 5^7 \times 7^5}$ is non-terminating repeating.

$$(viii) \frac{6}{15} = \frac{2 \times 3}{3 \times 5} = \frac{2}{5}$$

The denominator is of the form 5^n .

Hence, the decimal expansion of $\frac{6}{15}$ is terminating.

$$(ix) \frac{35}{50} = \frac{7 \times 5}{10 \times 5} = \frac{7}{10}$$

$$10 = 2 \times 5$$

The denominator is of the form $2^m \times 5^n$.

Hence, the decimal expansion of $\frac{35}{50}$ is terminating.

$$(x) \frac{77}{210} = \frac{11 \times 7}{30 \times 7} = \frac{11}{30}$$

$$30 = 2 \times 3 \times 5$$

Since the denominator is not of the form $2^m \times 5^n$, and it also has 3 as its factors, the decimal expansion of $\frac{77}{210}$ is non-terminating repeating.

Question 2:

Write down the decimal expansions of those rational numbers in Question 1 above which have terminating decimal expansions.

Solution 2:

$$(i) \frac{13}{3125} = 0.00416$$

$$\begin{array}{r} 0.00416 \\ 3125 \overline{) 13.00000} \end{array}$$

0

130

0

13000

12500

5000

3125

18750

18750

x

(ii) $\frac{17}{8} = 2.125$

$$\begin{array}{r} 2.125 \\ 8 \overline{) 17} \\ \underline{16} \\ 10 \\ \underline{8} \\ 20 \\ \underline{16} \\ 40 \\ \underline{40} \\ x \\ \underline{} \end{array}$$

(iv) $\frac{15}{1600} = 0.009375$

$$\begin{array}{r} 0.009375 \\ 1600 \overline{) 15.000000} \end{array}$$

0

150

0

1500

0

15000

14400

6000

4800

12000

11200

8000

8000

x

$$(vi) \frac{23}{2^3 \times 5^2} = \frac{23}{200} = 0.115$$

$$\begin{array}{r}
 00.115 \\
 200 \overline{) 23.000} \\
 \underline{0} \\
 23 \\
 \underline{- 0} \\
 230 \\
 \underline{- 200} \\
 300 \\
 \underline{- 200} \\
 1000 \\
 \underline{- 1000} \\
 0
 \end{array}$$

(viii) $\frac{6}{15} = \frac{2 \times 3}{3 \times 5} = \frac{2}{5} = 0.4$

$$\begin{array}{r}
 0.4 \\
 5 \overline{) 2.0} \\
 \underline{0} \\
 20 \\
 \underline{20} \\
 x
 \end{array}$$

(ix) $\frac{35}{50} = 0.7$

$$\begin{array}{r}
 0.7 \\
 50 \overline{) 35.0} \\
 \underline{0} \\
 350 \\
 \underline{350} \\
 x \\
 \underline{}
 \end{array}$$

Question 3:

The following real numbers have decimal expansions as given below. In each case, decide whether they are rational or not. If they are rational, and of the form $\frac{p}{q}$, what can you say about the prime factor

of q ?

- (i) 43.123456789
- (ii) 0.120120012000120000...
- (iii) $\overline{43.123456789}$

Solution 3:

(i) 43.123456789

Since this number has a terminating decimal expansion, it is a rational number of the form $\frac{p}{q}$ and q is of the form $2^m \times 5^n$

i.e., the prime factors of q will be either 2 or 5 or both.

(ii) 0.120120012000120000 ...

The decimal expansion is neither terminating nor recurring. Therefore, the given number is an irrational number.

(iii) $\overline{43.123456789}$

Since the decimal expansion is non-terminating recurring, the given

number is a rational number of the form $\frac{p}{q}$ and q is not of the form $2^m \times 5^n$

i.e., the prime factors of q will also have a factor other than 2 or 5.

Thank You
for downloading the PDF

FREE LIVE ONLINE

MASTER CLASSES

FREE Webinars by Expert Teachers



Vedantu

FREE MASTER CLASS SERIES

- ✓ For **Grades 6-12th** targeting **JEE, CBSE, ICSE** & much more
- ✓ **Free 60 Minutes Live Interactive** classes everyday
- ✓ Learn from the **Master Teachers** - India's best

Register for **FREE**

Limited Seats!