

UNIT 8

THE *d*- AND *f*-BLOCK ELEMENTS

1. Introduction

d-block elements are present from fourth period onwards. There are mainly three series of the transition metals – 3*d* series (Sc to Zn), 4*d* series (Y to Cd) and 5*d* series (La to Hg, omitting Ce to Lu).

d-block elements are known as transition elements because their position in the periodic table is between the s-block and p-block elements. Electronic configuration of the *d*-block elements is $(n-1)d^{1-10}ns^{0-2}$ but Cu⁺, Zn, Cd, Hg etc. $[(n-1)d^{10}]$ are *d*-block elements, but not transition metals because these have completely filled *d*-orbitals.

Transition Metals of *d*-block Elements

3rd group ns^2	4th group $d^1 ns^2 d^2$	5th group $ns^2 d^3$	6th group $ns^2 d^5$	7th group $ns^2 d^5$	8th group $ns^2 d^6$	9th group $ns^2 d^7$	10th group $ns^2 d^8$	11th group $ns^2 d^{10}$	12th group $ns^2 d^{10}$
$(n-1)$	$(n-1)$	$(n-1)$	$(n-1)$	$(n-1)$	$(n-1)$	$(n-1)$	$(n-1)$	$(n-1)$	$(n-1)$
Sc	Ti	V	Cr	Mn	Fe	Co	Ni	Cu	Zn
Y	Zr	Nb	Mo	Tc	Ru	Rh	Pd	Ag	Cd
La	Hf	Ta	W	Re	Os	Ir	Pt	Au	Hg

2. General Properties of the Transition Elements

(i) Atomic and Ionic Radii

In transition metals, left to right net nuclear charge increases due to poor shielding effect. Due to this, the atomic and ionic radii for transition elements for a given series show a decreasing trend for first five elements and then becomes almost constant for next five elements of the series.

(ii) Enthalpies of Atomisation

Transition elements exhibit higher enthalpies of atomization because of large number of unpaired electrons in their atoms. They have stronger interatomic interaction and hence, stronger bond.

(iii) Ionisation Enthalpies

- In a series from left to right, ionization enthalpy increases due to increase in nuclear charge.

- The irregular trend in the first ionization enthalpy of the 3*d* metals, though of little chemical significance, can be accounted for by considering that the removal of one electron alters the relative energies of 4*s* and 3*d* orbitals.

(iv) Oxidation States

- Transition metals show variable oxidation state due to two incomplete outermost shells. Only stable oxidation states of the first row transition metals are

Sc(+ 3), Ti(+ 4), V(+ 5), Cr(+ 3, + 6), Mn(+ 2, + 7), Fe(+ 2, + 3), Co(+ 2, + 3), Ni(+ 2), Cu(+ 2), Zn(+ 2)

- The transition elements in their lower oxidation states (+ 2 and + 3) usually form ionic compounds. In higher oxidation state compounds are normally covalent.
- Only Os and Ru show + 8 oxidation states in their compounds.
- Ni and Fe in Ni(CO)₄ and Fe(CO)₅ show zero oxidation state.

(v) Trends in the Standard Electrode Potentials

- Transformation of the solid metal atoms to M²⁺ ions in solution and their standard electrode potentials.
- If sum of the first and second ionization enthalpies is greater than hydration enthalpy standard potential ($E^\circ_{M^{2+}/M}$) will be positive and reactivity will be lower and vice-versa.

(vi) Trends in Stability of Higher Oxidation States

The higher oxidation numbers are achieved in TiX₄, VF₅ and CrF₆. The + 7 state for Mn is not represented in simple halides but MnO₃F is known and beyond Mn no metal has a trihalide except FeX₃ and CoF₃ and increasing order of oxidizing power in the series VO₂⁺ < Cr₂O₇²⁻ < MnO₄⁻.

(vii) Magnetic Properties

- When a magnetic field is applied to substances, mainly two types of magnetic behavior are observed : diamagnetism and paramagnetism. Paramagnetism due to presence of unpaired electrons, each such electron having a magnetic moment associated with its spin angular momentum.
- The magnetic moment is determined by the number of unpaired electrons.

$$\text{Magnetic moment} = \sqrt{n(n+2)}$$

where, *n* = number of unpaired electrons.

If all electrons are paired, substance will be diamagnetic and magnetic moment will be zero.

(viii) Formation of Coloured Ions

- The d -orbitals are non-degenerated in presence of ligands. When an electron from a lower energy d -orbital is excited to a higher energy d -orbital, the energy of required wavelength is absorbed and rest light is transmitted out. Therefore, the colour observed corresponds to the complementary colour of the light absorbed.

- In V_2O_5 , V is in + 5 oxidation state. It is coloured due to defects in crystal lattice.

(ix) Formation of Complex Compounds

- Transition metals have small size high nuclear charge which facilitates the acceptance of lone pair of electron from ligands.

- They have vacant d -orbitals of appropriate energy in order to accommodate the lone pair of electrons.

(x) Catalytic Properties

- Transition metals have two outermost shells incomplete and ability to adopt multiple oxidation states and to form complexes, therefore used as a catalyst.

- Transition metals also provide larger surface area for the reactant to be adsorbed.

(xi) Formation of Interstitial Compounds

- Small size of non-metals (H, C, N) fit into the voids of crystalline solid of transition metals and form interstitial compounds.

- The principal physical and chemical characteristics of these compounds are as follows :

- They have high melting points, higher than those of pure metals.
- They are very hard, some borides approach diamond in hardness.
- They retain metallic conductivity.
- They are chemically inert.

(xii) Alloy Formation

Alloy is the homogeneous mixture of two or more metals. Transition metals have approximate same size therefore, in molten form they can fit to each other crystalline structure and form homogeneous mixture and form the alloy.

E.g., Brass (copper-zinc) and bronze (copper-tin) etc.




3. Some Important Compounds of Transition Elements

Potassium Dichromate ($K_2Cr_2O_7$)

(i) Ore


Ferrochrome or chromate ($FeO.Cr_2O_3$) or ($FeCr_2O_4$)

(ii) Preparation






Sodium dichromate is more soluble than potassium dichromate.

- Chromates and dichromates are interconvertible in aqueous solution depending upon pH of the solutions.



(iii) Properties

Sodium and potassium dichromates are strong oxidizing agents, thus, acidified $\text{K}_2\text{Cr}_2\text{O}_7$ will oxidise iodides to iodine, sulphides to sulphur, tin (II) to tin (IV) and iron (II) salts to iron (III).



(iv) Uses


- $\text{K}_2\text{Cr}_2\text{O}_7$ is used as oxidizing agent in volumetric analysis.
- It is used in mordant dyes, leather industry, photography (for hardening of film).
- It is used in chromyl chloride test.
- It is used in cleaning glassware.

Potassium Permanganate (KMnO_4)



(i) Ore

Pyrolusite (MnO_2)

(ii) Preparation




(iii) Commercial preparation





(iv) Properties

KMnO₄ acts as strong oxidizing agent.

(a) In presence of dilute H₂SO₄, KMnO₄ is reduced to manganous salt.




Acidic KMnO₄ solution oxidizes oxalates to CO₂, iron (II) and iron (III), nitrites to nitrates and iodides to iodine. The half reactions of reductants are



To acidify KMnO₄, only H₂SO₄ is used and not HCl or HNO₃ because HCl reacts with KMnO₄ and produce Cl₂ while HNO₃, itself acts as oxidizing agent.

(b) In alkaline medium, KMnO₄ is reduced to insoluble MnO₂.



Alkaline or neutral KMnO₄ solution oxidizes I⁻ to IO₃⁻, S₂O₃²⁻ to SO₄²⁻, Mn²⁺ to MnO₂ etc.

(v) Uses

(a) In laboratory preparation of Cl₂.

(b) KMnO₄ is used as an oxidizing agent, disinfectant.

(c) In making Baeyer's reagent.

4. The Inner Transition Elements (f-Block)


The f-block consists of the two series, lanthanoids and actinoids. Lanthanoids are known as rare earth metals and actinoids are known as radioactive elements (Th to Lr).

Lanthanoids

General characteristics

- General configuration [Xe] 4f¹⁻¹⁴, 5d⁰⁻¹, 6s².
- Atomic and ionic size from left to right, decreases due to increase in nuclear charge. This is known as lanthanoid contraction.
- All the lanthanoids are silvery white soft metals and tarnish rapidly in air.

- Many trivalent lanthanoid ions are coloured both in the solid state and in aqueous solutions. Neither La^{3+} nor Lu^{3+} ion shows any colour but the rest do so.
- The lanthanoid ions other than the f^0 type (La^{3+} and Ce^{4+}) and the f^{14} type (Yb^{2+} and Lu^{3+}) are all paramagnetic. The paramagnetism arises to maximum in neodymium.
- Oxidation states $\rightarrow \text{Ce}^{4+}$; (Some elements) is favoured by its noble gas configuration, but it is a strong oxidant reverting to the common + 3 state. The E°_{value} for $\text{Ce}^{4+}/\text{Ce}^{3+}$ is + 1.74 V, the reaction rate is very slow and hence, Ce(IV) is a good analytical reagent. Pr, Nd, Tb and Dy also exhibit + 4 state but only in oxides. Eu^{2+} is formed by losing the two s-electrons and its f^7 configuration accounts for the formation of this ion. However, Eu^{2+} is a strong reducing agent changing to the common + 3 state. Similarly, Yb^{2+} which has f^{14} configuration is a reductant, Tb^{4+} has half-filled *f*-orbitals and is an oxidant.
- Lanthanoid are very reactive metals like alkaline earth metals.



- Misch metals, contain lanthanoids about 90-95% (Ce 40-5%, Lanthanum and neodymium 44%) iron 4.5%, calcium, carbon and silicon about 10.5% are used in cigarette and gas lighters, toys, tank and tracer bullets.

Actinoids

- Genral configuration $[\text{Rn}] 5f^{1-14}, 6d^{0-2}, 7s^2$.
- Actinoids exhibit a range of oxidation states due to comparable energies of $5f$, $6d$ and $7s$ orbitals. The genral oxidation state of actinoids is + 3.
- All the actinoids are strong reducing agents and very reactive.
- Actinoids also react with oxygen, halogen, hydrogen and sulphur, etc. like lanthanoids.
- Actinoids are radioactive in nature and therefore, it is difficult to study their chemical nature.

VERY SHORT ANSWER TYPE QUESTIONS (1 Mark)

Q. 1. Explain $\text{CuSO}_4 \cdot 5\text{H}_2\text{O}$ is blue while CuSO_4 is colourless ?

Ans. Because water molecules act as ligands and results in crystal field splitting of *d*-orbitals of Cu^{2+} ion.

Q. 2. Which element among $3d$ series exhibit highest oxidation state ?

Ans. Mn

Q. 3. In 3d series (Sc to Zn), the enthalpy of atomization of Zn is low. Why ?

Ans. Due to absence of unpaired electrons.

Q. 4. Which element among 3d series exhibit only one oxidation state ?

Ans. Sc

Q. 5. Why is the 3rd ionization energy of Mn ($Z = 25$) is unexpectedly high ?

Ans. Due to half-filled electronic configuration.

Q. 6. Define alloy.

Ans. Alloys are homogeneous solid solutions of two or more metals.

Q. 7. Transition metals show zero oxidation state with ligands like CO. Explain.

Ans. Co form synergic bonding with metal ion.

Q. 8. Why can't HCl acid be used to acidify KMnO_4 solution ?

Ans. Because KMnO_4 oxidize HCl into Cl_2 .

Q. 9. Name one ore of Mn and Cr.

Ans. Mn : MnO_2

Cr : FeCr_2O_4

Q. 10. Why Mn^{2+} compounds are more stable than Fe^{2+} compounds towards oxidation to their + 3 state ?

Ans. Mn^{+2} has half-filled electronic configuration.

Q. 11. Why do transition metal (elements) show variable oxidation states ?

Ans. Due to presence of vacant d-orbitals.

Q. 12. Write any uses of pyrophoric alloy.

Ans. Making bullets, shells and lighter flints.

Q. 13. Which is more basic – $\text{La}(\text{OH})_3$ or $\text{Lu}(\text{OH})_3$? Why ?

Ans. $\text{La}(\text{OH})_3$, due to lanthanide contraction, lower size, more covalent character, least basic.

Q. 14. Find out number of Cr – O – Cr bond/bonds in $\text{Cr}_2\text{O}_7^{2-}$ ion.

Ans. 1

Q. 15. What is effect of increasing pH on $\text{K}_2\text{Cr}_2\text{O}_7$ solution ?

Ans. It changes into K_2CrO_4 solution/orange colour change into yellow colour.

Q. 16. Why is Ce^{4+} in aqueous solution a good oxidizing agent ?

Ans. Because Ce^{4+} is most stable in Ce^{+3} state in aqueous solution.

Q. 17. Why do Zr and Hf exhibit similar properties ?

Ans. Due to lanthanide contraction.

Q. 18. What is lanthanoid contraction ?

Ans. The regular slow decrease in the atomic or ionic radii of lanthanoids with increasing atomic number.

Q. 19. Why is Cu ($Z = 29$) considered a transition metal ?

Ans. Due to its partially filled *d*-orbital in Cu^{2+} state.

Q. 20. Arrange the given in increasing order of acidic character :

$\text{CrO}_3, \text{CrO}, \text{Cr}_2\text{O}_3$.

Ans. $\text{CrO} < \text{Cr}_2\text{O}_3 < \text{CrO}_3$

Q. 21. Why KMnO_4 or MnO_4^- ion is coloured ?

Ans. Due to charge transfer complex formation.

SHORT ANSWER-I TYPE QUESTIONS (2 Marks)

Q. 1. Chromium is typical hard metal while mercury is a liquid. Explain why ?

Ans. Cr has five unpaired *d*-electrons. Hence metallic bonds are strong. In Hg, there is absence of unpaired electrons and size is larger.

Q. 2. Why KMnO_4 is used in cleaning surgical instruments in hospitals ?

Ans. This is because KMnO_4 has a germicidal action.

Q. 3. Most of the transition metals do not displace hydrogen from dilute acids, why ?

Ans. Due to their –ve reduction potential.

Q. 4. Explain why Cu^+ is not stable in aqueous solution ?

Ans. Due to less –ve $\Delta_{\text{hyd}}H^\circ$ of Cu^+ /it cannot compensate 2nd ionization potential of Cu.

Q. 5. Why is the highest oxidation state of a metal exhibited in its oxide or fluoride only ?




Ans. Oxygen and fluoride have small size and high electronegativity. They can oxidise the metal.

Q. 6. Write electronic configuration of Cu^{+2} and Co^{+2} .

Ans. $\text{Cu}^{+2} = [\text{Ar}] 3d^9 4s^0$

$\text{Co}^{+2} = [\text{Ar}] 3d^7$

Q. 7. Balance the following equations :



Q. 8. Briefly explain why electronic configuration of lanthanoids are not known with certainty ?

Ans. $4f/5d$ subshells are very close in energy. +ve electrons can jump from $4f$ to $5d$ or vice-versa.

Q. 9. Why Zn, Cd, Hg are soft and have low melting point ?

Ans. Due to weak interatomic attraction/absence of unpaired electrons.

Q. 10. What is the effect of pH on the solution of $\text{K}_2\text{Cr}_2\text{O}_4$ solution ?

Ans. $\text{K}_2\text{Cr}_2\text{O}_4$ solution changes into $\text{K}_2\text{Cr}_2\text{O}_7$ /yellow colour changes into orange colour.

Q. 11. Which of the following is/are transition element and why ?

Zn, Cd, Ag, Fe, Ni

Ans. Fe, Ni, Ag

Q. 12. What are interstitial compounds ? Give example.

Ans. When small atoms like C, H, B and N occupy interstitial site in their lattice. Example, TiC, Fe_3H ,

Q. 13. Why are Zn, Cd, Hg volatile metals ? Explain.



Ans. Due to weak interatomic attraction/low boiling point.

Q. 14. Why is first ionization energy of $5d$ elements higher than those of $3d$ and $4d$ elements ?

Q. 15. Explain 'Misch metal' and write its use.

Ans. It is an alloy of 95% lanthanoid and 5% iron and traces of S, C, Ca and Al. Used in lighter flint, bullet tips etc.

Q. 16. The following two reactions of HNO_3 with Zn are given :





Identify X and Y.

Ans. $\text{X} = \text{NO}_2$

$\text{Y} = \text{N}_2\text{O}$

Q. 17. Complete the equations :



Q. 18. Out of Fe and Cu, which one would exhibit higher melting point ?


Ans. Fe, due to large number of unpaired *d*-electrons/more interatomic attraction.

Q. 19. Sc, the first member of first transition series does not exhibit variable oxidation state. Why ?

Ans. Due to noble gas electronic configuration in + 3 oxidation state no other oxidation state is stable.

SHORT ANSWER-II TYPE QUESTIONS (3 Marks)

Q. 1. (a) Deduce the number of 3*d* electrons in the following ions :



(b) **Why do transition metals form alloy ?**

(c) **Why Zn^{+2} salts are white ?**





Ans. (a) Cu^{+2} : 9 electrons

Sc^{+3} : 0 electron

(b) Transition metals have similar atomic radii.

(c) Absence of unpaired electron.


Q. 2. Complete and balance the following equations :




Q. 3. Describe the oxidizing action of $\text{K}_2\text{Cr}_2\text{O}_7$ with the following. Write ionic equations for its reaction (acidic medium) with :

(a) I^- (b) Iron(II) (c) H_2S


Ans. (a) It liberates I_2 from I^- .



- (b) It oxidizes Fe^{+2} to Fe^{+3} .



- (b) It oxidizes H_2S to sulphur.



Q. 4. Write any four differences between lanthanoids and actinoids.

Ans.

Lanthanoids

Actinoids

- | | |
|--|--|
| (a) They are generally non-radioactive. | They are radioactive. |
| (b) Most of their ions are colourless. | Coloured ions. |
| (c) Show + 3, + 4 and + 2 oxidation state. | Beside + 3, show higher oxidation state. |
| (d) Less tendency of complex formation. | Higher tendency. |

Q. 5. (a) Why is separation of lanthanoid elements difficult ?

- (b) Transition metal exhibit higher enthalpies of atomization. Explain why ?**

- (c) Why have the transition metal high enthalpy of hydration ?**

Ans.

- (a) Due to lanthanide contraction, the size of these elements is nearly same.
- (b) Transition metal contain large number of unpaired electrons, and they have strong interatomic attractions.
- (c) Due to their small size and large nuclear charge.

Q. 6. (a) Use Hund's rule to derive the electronic configuration of Ce^{+3} ions and calculate its magnetic moment.

- (b) Is lanthanum a *f*-block element ?**

Ans.

- (a) ${}_{58}\text{Ce} = [\text{Xe}] 4f^1 5d^1 6s^2$
 $\text{Ce}^{+3} = 4f^1$ one unpaired electron

$$\mu = \sqrt{n(n+2)} = 1.73 \text{ BH}$$


- (b) No, it is a *d*-block element.

Q. 7. Account for the following :

- (a) Silver chloride dissolves in excess of NH_3 .
- (b) Cuprous chloride is diamagnetic while cupric chloride is paramagnetic.
- (c) In CrO_4^{2-} ion, all the Cr – O bond length are equal.

Ans.

- (a) AgCl forms a soluble complex with NH_3 .



- (b) $\text{Cu}^+ : 3d^{10} 4s^0$ – All electrons are paired.
 $\text{Cu}^{+2} : 3d^9$ – Here, one unpaired electron is present.
- (c) Due to resonance.

Q. 8. The E° values in respect of electrodes of Cr, Mn and iron are :

$$\text{Cr}^{+3}/\text{Cr}^{+2} = -0.4 \text{ V}$$

$$\text{Mn}^{+3}/\text{Mn}^{+2} = +1.5 \text{ V}$$

$$\text{Fe}^{+3}/\text{Fe}^{+2} = +0.8 \text{ V}$$

Compare the feasibilities of further oxidation of these ions.

Ans. Cr^{+3} is more stable than Cr^{+2} .

Mn^{+2} is more stable than Mn^{+3} .

Fe^{+3} is more stable than Fe^{+2} .

Order of feasibility of + 2 oxidation state is :

$$\text{Mn}^{+2} > \text{Fe}^{+2} > \text{Cr}^{+2}$$

Q. 9. Write any three properties of interstitial compounds.

- Ans.** (a) They are chemically inert.
 (b) They retain metallic conductivity.
 (c) They have high melting point than their pure metals.
 (d) These are harder and more corrosion resistant.

Q. 10. Account for the following :

- (a) All Scandium salts are white.
 (b) The 1st ionization energy of the 5*d* series are higher than 3*d* and 4*d* transition elements in respective groups.
 (c) Ce^{+3} can be easily oxidized to Ce^{+4} .

- Ans.** (a) Sc has only + 3 oxidation state, there is no unpaired electron.
 (b) Due to lanthanide contraction, effective nuclear charge increase.
 (c) Due to gain noble gas electron configuration.


Q. 11. A green chromium compound (A) on fusion with alkali gives a yellow compound (B) which on acidification gives a orange coloured compound (C). Identify A, B, C. Write equations for reactions.

- Ans.** A : FeCr_2O_4 B : Na_2CrO_4 C. $\text{Na}_2\text{Cr}_2\text{O}_7$

$$4\text{FeCr}_2\text{O}_4 + 8\text{Na}_2\text{CO}_3 + 7\text{O}_2 \rightarrow 8\text{Na}_2\text{CrO}_4 + 2\text{Fe}_2\text{O}_3 + 8\text{CO}_2$$

(A)

(B)



(C)

Q. 12. When an oxide of Mn (A) is fused with KOH in the presence of an oxidizing agent and dissolved in water, it gives a dark solution of compound (B). Compound (B) disproportionate in neutral or acidic solution to give purple compound (C). Identify A, B, C.

Ans. A : MnO_2 B : K_2MnO_4 C : KMnO_4

LONG ANSWER TYPE QUESTIONS (5 Marks)

Q. 1. A violet compound of manganese (A) decomposes on heating to liberate oxygen and compounds (B) and (C) of manganese are formed. Compound (C) reacts with KOH in the presence of KNO_3 to give compound (B). On heating compound (C) with conc. H_2SO_4 and NaCl, Cl_2 gas is liberated and compound (D) of manganese is formed. Identify A, B, C, D alongwith reactions involved.


Ans. A : KMnO_4 B : K_2MnO_4 C : MnO_2 D : MnCl_2



(A) (B)



(C) (B)




(C) (D)

Q. 2. (a) What is meant by disproportionation of an oxidation state ? Give one example.

(b) Draw the structures of $\text{Cr}_2\text{O}_7^{2-}$, CrO_4^{2-} , MnO_4^- .

(c) What is the effect of lanthanoids contraction beyond lanthanoid ?

Ans. (a) When any atom or ion undergo oxidation and reduction simultaneously it is called disproportionation.



(c) Size of respective 4d and 5d series elements becomes comparable from fourth group onwards (e.g., Zr and Hf).

VALUE BASED QUESTIONS (4 Marks)

- Q. 1.** A group of students was smoking cigarettes in college premises. A social activist noticed and advised them not to smoke.
- By forbidding them not to smoke, which value of social activity wants to be inculcated among the youth ?
 - As a chemistry student, write the name of 'alloy' used in lighter's flint.
 - Mention diseases caused by smoking.
 - Name the chemical present in the smoke which is habit forming.
- Q. 2.** It is a general myth that we should not come out of the house to see solar eclipse because it can have evil impact on life but nowadays educated people allow their children to see solar eclipse, treating them as a natural science phenomenon, but children are advised to see them by uv protected sun glasses (crooke's lenses) to avoid harmful impact of uv light on eyes.
- Write the name of *d*-block metal oxide used in making uv protected lens.
 - By allowing the children to see solar eclipse using uv protected lens, which value the educated people are trying to inculcate in the children ?
 - Which rays are present in the light which can damage the eye while viewing solar eclipse with naked eye ?
 - Name the oxide of a lanthanoid also present in uv protected lenses.
- Q. 3.** In Chemistry lab, Lalit used $\text{Na}_2\text{Cr}_2\text{O}_7$ for the estimation of Fe^{2+} and I^- in redox reactions while Sahil uses $\text{K}_2\text{Cr}_2\text{O}_7$ for the same estimation of ions.
- Out of these two, who is performing correct estimation of ions ?
 - Draw the structure of $\text{Cr}_2\text{O}_7^{2-}$ ion.
 - Explain why CrO_4^{2-} and $\text{Cr}_2\text{O}_7^{2-}$ are interconvertible by the change in pH of the medium.
 - Name the value learnt by the use of $\text{K}_2\text{Cr}_2\text{O}_7$ in the estimation of ions.
- Q. 4.** For automotive catalytic converters, transition metals like Pt, Pd, Rh, Fe, Cr etc. used in industry are expensive while ceramic cores are inexpensive.
- Which property of transition metal makes them useful as catalyst ?
 - What is the function of catalytic convertor ?
 - As a student of chemistry, would you suggest the use of explosive metals like Pt, Pd and Rh ? Give reasons.
 - Which value is promoted through the use of catalytic convertors in industry ?

- Q. 5.** Due to scarcity to water, the resident of a colony decided to refuse their well which was not used for the past six months. The secretary of RWA decided to clean the well by adding small amount of potassium permanganate in the well and then loaded the well water with alum crystals.
- (a) Why did the secretary of RWA add potassium permanganate in well ?
 - (b) Draw the structure of permanganate ion, MnO_4^- .
 - (c) Why did secretary of RWA add alum to the well ?
 - (d) What values are associated with the use of KMnO_4 and alum ?

