

Chemistry

(www.tiwariacademy.com)

(Chapter 1)(The Solid State)

XII

Intext Questions

Question 1.1:

Why are solids rigid?

Answer

The intermolecular forces of attraction that are present in solids are very strong. The constituent particles of solids cannot move from their positions i.e., they have fixed positions. However, they can oscillate about their mean positions. This is the reason solids are rigid.

Question 1.2:

Why do solids have a definite volume?

Answer

The intermolecular forces of attraction that are present in solids are very strong. The constituent particles of solids have fixed positions i.e., they are rigid. Hence, solids have a definite volume.

Question 1.3:

Classify the following as amorphous or crystalline solids:

Polyurethane, naphthalene, benzoic acid, teflon, potassium nitrate, cellophane, polyvinyl chloride, fibre glass, copper.

Answer

Amorphous solids

Polyurethane, teflon, cellophane, polyvinyl chloride, fibre glass

Crystalline solids

Naphthalene, benzoic acid, potassium nitrate, copper

Question 1.4:

Why is glass considered a super cooled liquid?



Chemistry

(www.tiwariacademy.com)

(Chapter 1)(The Solid State)

XII

Answer

Similar to liquids, glass has a tendency to flow, though very slowly. Therefore, glass is considered as a super cooled liquid. This is the reason that glass windows and doors are slightly thicker at the bottom than at the top.

Question 1.5:

Refractive index of a solid is observed to have the same value along all directions. Comment on the nature of this solid. Would it show cleavage property?

Answer

An isotropic solid has the same value of physical properties when measured along different directions. Therefore, the given solid, having the same value of refractive index along all directions, is isotropic in nature. Hence, the solid is an amorphous solid. When an amorphous solid is cut with a sharp edged tool, it cuts into two pieces with irregular surfaces.

Question 1.6:

Classify the following solids in different categories based on the nature of intermolecular forces operating in them:

Potassium sulphate, tin, benzene, urea, ammonia, water, zinc sulphide, graphite, rubidium, argon, silicon carbide.

Answer

Potassium sulphate → Ionic solid

Tin → Metallic solid

Benzene → Molecular (non-polar) solid

Urea → Polar molecular solid

Ammonia → Polar molecular solid

Water → Hydrogen bonded molecular solid

Zinc sulphide → Ionic solid

Graphite → Covalent or network solid

Rubidium → Metallic solid

Argon → Non-polar molecular solid



Chemistry

(www.tiwariacademy.com)

(Chapter 1)(The Solid State)

XII

Silicon carbide → Covalent or network solid

Question 1.7:

Solid A is a very hard electrical insulator in solid as well as in molten state and melts at extremely high temperature. What type of solid is it?

Answer

The given properties are the properties of a covalent or network solid. Therefore, the given solid is a covalent or network solid. Examples of such solids include diamond (C) and quartz (SiO_2).

Question 1.8:

Ionic solids conduct electricity in molten state but not in solid state. Explain.

Answer

In ionic compounds, electricity is conducted by ions. In solid state, ions are held together by strong electrostatic forces and are not free to move about within the solid. Hence, ionic solids do not conduct electricity in solid state. However, in molten state or in solution form, the ions are free to move and can conduct electricity.

Question 1.9:

What type of solids are electrical conductors, malleable and ductile? Answer

Metallic solids are electrical conductors, malleable, and ductile.

Question 1.10:

Give the significance of a 'lattice point'.

Answer

The significance of a lattice point is that each lattice point represents one constituent particle of a solid which may be an atom, a molecule (group of atom), or an ion.

Question 1.11:

Name the parameters that characterize a unit cell.

Answer



Chemistry

(www.tiwariacademy.com)

(Chapter 1)(The Solid State)

XII

The six parameters that characterise a unit cell are as follows.

(i) Its dimensions along the three edges, a , b , and c . These edges may or may not be equal.

(ii) Angles between the edges

These are the angle α (between edges b and c), β (between edges a and c), and γ (between edges a and b).

Question 1.12:

Distinguish between

(i) Hexagonal and monoclinic unit cells

(ii) Face-centred and end-centred unit cells.

Answer

(i) Hexagonal unit cell

For a hexagonal unit cell,

$$a = b \neq c$$

$$\text{and } \alpha = \beta = 90^\circ$$

$$\gamma = 120^\circ$$

Monoclinic unit cell

For a monoclinic cell,

$$a \neq b \neq c$$

$$\text{and } \alpha = \gamma = 90^\circ$$


$$\beta \neq 90^\circ$$

(ii) Face-centred unit cell

In a face-centred unit cell, the constituent particles are present at the corners and one at the centre of each face.

End-centred unit cell

An end-centred unit cell contains particles at the corners and one at the centre of any two opposite faces.



Chemistry

(www.tiwariacademy.com)

(Chapter 1)(The Solid State)

XII

Question 1.13:

Explain how much portion of an atom located at **(i)** corner and **(ii)** body-centre of a cubic unit cell is part of its neighbouring unit cell.

Answer

(i) An atom located at the corner of a cubic unit cell is shared by eight adjacent unit cells.

Therefore, $\frac{1}{8}$ th portion of the atom is shared by one unit cell.

(ii) An atom located at the body centre of a cubic unit cell is not shared by its neighbouring unit cell. Therefore, the atom belongs only to the unit cell in which it is present i.e., its contribution to the unit cell is 1.

Question 1.14:

What is the two dimensional coordination number of a molecule in square close packed layer?

Answer

In square close-packed layer, a molecule is in contact with four of its neighbours. Therefore, the two-dimensional coordination number of a molecule in square closepacked layer is 4.

Question 1.15:

A compound forms *hexagonal close-packed* structure. What is the total number of voids in 0.5 mol of it? How many of these are tetrahedral voids?


Answer

Number of close-packed particles = $0.5 \times 6.022 \times 10^{23} = 3.011 \times 10^{23}$

Therefore, number of octahedral voids = 3.011×10^{23}

And, number of tetrahedral voids = $2 \times 3.011 \times 10^{23} = 6.022 \times 10^{23}$

Therefore, total number of voids = $3.011 \times 10^{23} + 6.022 \times 10^{23} = 9.033 \times 10^{23}$



Chemistry

(www.tiwariacademy.com)

(Chapter 1)(The Solid State)

XII

Question 1.16:

A compound is formed by two elements M and N. The element N forms *ccp* and atoms of M occupy $\frac{1}{3}$ rd of tetrahedral voids. What is the formula of the compound?

Answer

The *ccp* lattice is formed by the atoms of the element N.

Here, the number of tetrahedral voids generated is equal to twice the number of atoms of the element N.

According to the question, the atoms of element M occupy $\frac{1}{3}$ rd of the tetrahedral voids.

$$2 \times \frac{1}{3} = \frac{2}{3}$$

Therefore, the number of atoms of M is equal to $\frac{2}{3}$ of the number of atoms of N.

Therefore, ratio of the number of atoms of M to that of N is M: N $= \frac{2}{3} : 1$
 $= 2 : 3$

Thus, the formula of the compound is $M_2 N_3$.

Question 1.17:

Which of the following lattices has the highest packing efficiency (i) simple cubic (ii) body-centred cubic and (iii) hexagonal close-packed lattice?

Answer

Hexagonal close-packed lattice has the highest packing efficiency of 74%. The packing efficiencies of simple cubic and body-centred cubic lattices are 52.4% and 68% respectively.

Question 1.18:

An element with molar mass $2.7 \times 10^{-2} \text{ kg mol}^{-1}$ forms a cubic unit cell with edge length 405 pm. If its density is $2.7 \times 10^3 \text{ kg m}^{-3}$, what is the nature of the cubic unit cell?

Answer

It is given that density of the element, $d = 2.7 \times 10^3 \text{ kg m}^{-3}$



Chemistry

(www.tiwariacademy.com)

(Chapter 1)(The Solid State)

XII

Molar mass, $M = 2.7 \times 10^{-2} \text{ kg mol}^{-1}$

Edge length, $a = 405 \text{ pm} = 405 \times 10^{-12} \text{ m}$

$= 4.05 \times 10^{-10} \text{ m}$

It is known that, Avogadro's number, $N_A = 6.022 \times 10^{23} \text{ mol}^{-1}$ Applying the relation,

$$\begin{aligned}d &= \frac{z \cdot M}{a^3 \cdot N_A} \\z &= \frac{d \cdot a^3 N_A}{M} \\&= \frac{2.7 \times 10^{-2} \text{ kg mol}^{-1} \times (4.05 \times 10^{-10} \text{ m})^3 \times 6.022 \times 10^{23} \text{ mol}^{-1}}{2.7 \times 10^{-2} \text{ kg mol}^{-1}} \\&= 4.004 \\&= 4\end{aligned}$$

This implies that four atoms of the element are present per unit cell. Hence, the unit cell is face-centred cubic (fcc) or cubic close-packed (ccp).

Question 1.19:

What type of defect can arise when a solid is heated? Which physical property is affected by it and in what way?

Answer

When a solid is heated, vacancy defect can arise. A solid crystal is said to have vacancy defect when some of the lattice sites are vacant.

Vacancy defect leads to a decrease in the density of the solid.

Question 1.20:

What type of stoichiometric defect is shown by:

(i) ZnS (ii) AgBr

Answer

(i) ZnS shows Frenkel defect.

(ii) AgBr shows Frenkel defect as well as Schottky defect.



Chemistry

(www.tiwariacademy.com)

(Chapter 1)(The Solid State)

XII

Question 1.21:

Explain how vacancies are introduced in an ionic solid when a cation of higher valence is added as an impurity in it.

Answer

When a cation of higher valence is added to an ionic solid as an impurity to it, the cation of higher valence replaces more than one cation of lower valence so as to keep the crystal electrically neutral. As a result, some sites become vacant. For example, when Sr^{2+} is added to NaCl, each Sr^{2+} ion replaces two Na^+ ions. However, one Sr^{2+} ion occupies the site of one Na^+ ion and the other site remains vacant. Hence, vacancies are introduced.

Question 1.22:

Ionic solids, which have anionic vacancies due to metal excess defect, develop colour.

Explain with the help of a suitable example.

Answer

The colour develops because of the presence of electrons in the anionic sites. These electrons absorb energy from the visible part of radiation and get excited.

For example, when crystals of NaCl are heated in an atmosphere of sodium vapours, the sodium atoms get deposited on the surface of the crystal and the chloride ions from the crystal diffuse to the surface to form NaCl with the deposited Na atoms. During this process, the Na atoms on the surface lose electrons to form Na^+ ions and the released electrons diffuse into the crystal to occupy the vacant anionic sites. These electrons get excited by absorbing energy from the visible light and impart yellow colour to the crystals.

Question 1.23:

A group 14 element is to be converted into n -type semiconductor by doping it with a suitable impurity. To which group should this impurity belong?

Answer

An n -type semiconductor conducts because of the presence of extra electrons. Therefore, a group 14 element can be converted to n -type semiconductor by doping it with a group 15 element.



Chemistry

(www.tiwariacademy.com)

(Chapter 1)(The Solid State)

XII

Question 1.24:

What type of substances would make better permanent magnets, ferromagnetic or ferrimagnetic. Justify your answer.

Answer

Ferromagnetic substances would make better permanent magnets.

In solid state, the metal ions of ferromagnetic substances are grouped together into small regions. These regions are called domains and each domain acts as a tiny magnet. In an unmagnetised piece of a ferromagnetic substance, the domains are randomly oriented. As a result, the magnetic moments of the domains get cancelled. However, when the substance is placed in a magnetic field, all the domains get oriented in the direction of the magnetic field and a strong magnetic effect is produced.

The ordering of the domains persists even after the removal of the magnetic field. Thus, the ferromagnetic substance becomes a permanent magnet.

