

QUESTION PAPER CODE 65/2/2/F
EXPECTED ANSWER/VALUE POINTS
SECTION A

1. $P = \frac{1}{2}(A + A')$ $\therefore P = \begin{bmatrix} 3 & 6 \\ 6 & 9 \end{bmatrix}$ $\frac{1}{2} + \frac{1}{2}$

2. $(\vec{a} + \vec{b} + \vec{c}) \cdot (\vec{a} + \vec{b} + \vec{c}) = 0$ $\frac{1}{2}$

$$\Rightarrow |\vec{a}|^2 + |\vec{b}|^2 + |\vec{c}|^2 + 2(\vec{a} \cdot \vec{b} + \vec{b} \cdot \vec{c} + \vec{c} \cdot \vec{a}) = 0$$

$$\Rightarrow \vec{a} \cdot \vec{b} + \vec{b} \cdot \vec{c} + \vec{c} \cdot \vec{a} = -\frac{3}{2} \quad \frac{1}{2}$$

3. $a^2 b^2 \sin^2 \theta + a^2 b^2 \cos^2 \theta = 400$ $\frac{1}{2}$

$$\Rightarrow |\vec{b}| = 4 \quad \frac{1}{2}$$

4. $\frac{x}{\sqrt{3}} + \frac{y}{\sqrt{3}} + \frac{z}{\sqrt{3}} = 5\sqrt{3}$ or $x + y + z = 15$ $\left[\frac{1}{2} \text{ mark for dc's of normal} \right]$ 1

5. 1×1 1

6. Expanding we get

$$x^3 = -8 \Rightarrow x = -2 \quad \frac{1}{2} + \frac{1}{2}$$

SECTION B

7. $f'_{1-} = 2x + 3 = 5$
 $f'_{1+} = b$

$$f'_{1-} = f'_{1+} \Rightarrow [b = 5] \quad 1+1$$

$$\lim_{x \rightarrow 1^-} f(x) = f(1) = \lim_{x \rightarrow 1^+} f(x)$$

$$\Rightarrow 4 + a = b + 2 \quad 1$$

$$\Rightarrow [a = 3] \quad 1$$

8. Let $u = \tan^{-1} \frac{\sqrt{1+x^2} - 1}{x}$

$$\text{Put } x = \tan \theta \Rightarrow \theta = \tan^{-1} x \quad \frac{1}{2}$$

$$\therefore u = \tan^{-1} \left[\frac{\sec \theta - 1}{\tan \theta} \right]$$

$$= \tan^{-1} \left[\frac{1 - \cos \theta}{\sin \theta} \right]$$

$$= \tan^{-1} \left(\tan \frac{\theta}{2} \right)$$

$$= \frac{\theta}{2} = \frac{1}{2} \tan^{-1} x$$

$$\Rightarrow \frac{du}{dx} = \frac{1}{2(1+x^2)}$$

$$v = \sin^{-1} \left(\frac{2x}{1+x^2} \right)$$

$$= 2 \tan^{-1} x$$

$$\Rightarrow \frac{dv}{dx} = \frac{2}{1+x^2}$$

$$\therefore \frac{du}{dv} = \frac{du/dx}{dv/dx} = \frac{1}{4}$$

OR

$$x = \sin t \Rightarrow \frac{dx}{dt} = \cos t$$

$$y = \sin pt \Rightarrow \frac{dy}{dt} = p \cos pt$$

$$\frac{dy}{dx} = \frac{p \cos pt}{\cos t}$$

$$\frac{d^2y}{dx^2} = \frac{\cos t (-p^2 \sin pt) - p \cos pt (-\sin t)}{\cos^2 t} \cdot \frac{dt}{dx}$$

$$= \frac{-p^2 \sin pt \cos t + p \cos pt \sin t}{\cos^3 t}$$

$$\text{Now } (1-x^2) \frac{d^2y}{dx^2} - x \frac{dy}{dx} + p^2 y = 0 \quad \left[\text{Substituting values of } y, \frac{dy}{dx} \text{ & } \frac{d^2y}{dx^2} \right]$$

9. Eqn of given curves

$$y^2 = 4ax \text{ and } x^2 = 4by$$

Their point of intersections are $(0, 0)$ and $(4a^{1/3} b^{2/3}, 4a^{2/3} b^{1/3})$

$$y^2 = 4ax \Rightarrow \frac{dy}{dx} = \frac{2a}{y}, \text{ slope} = \frac{a^{1/3}}{2b^{1/3}} \quad \dots(i)$$

$$x^2 = 4by \Rightarrow \frac{dy}{dx} = \frac{x}{2b}, \text{ slope} = \frac{2a^{1/3}}{b^{1/3}} \quad \dots(ii)$$

At $(0, 0)$, angle between two curves is 90°

or

Acute angle θ between (i) and (ii) is

$$\theta = \tan^{-1} \left\{ \frac{3}{2} \left(\frac{a^{1/3} b^{1/3}}{a^{2/3} + b^{2/3}} \right) \right\}$$

(13)

10. $I = \int_0^\pi \frac{(\pi - x)}{1 + \sin \alpha \sin(\pi - x)} dx$ 1

$$2I = \pi \int_0^\pi \frac{dx}{1 + \sin \alpha \sin x}$$

$$= 2\pi \int_0^{\pi/2} \frac{dx}{1 + \sin \alpha \sin x}$$

$$\Rightarrow I = \pi \int_0^{\pi/2} \frac{dx}{1 + \sin \alpha \frac{2 \tan \frac{x}{2}}{1 + \tan^2 \frac{x}{2}}} \quad 1$$

$$I = \pi \int_0^1 \frac{2dt}{1 + t^2 + 2t \sin \alpha} \quad \text{Put } \tan \frac{x}{2} = t \quad \frac{1}{2}$$

$$\Rightarrow I = 2\pi \int_0^1 \frac{dt}{(t + \sin \alpha)^2 + \cos^2 \alpha} \quad 1$$

$$= \frac{2\pi}{\cos \alpha} \left[\tan^{-1} \left(\frac{t + \sin \alpha}{\cos \alpha} \right) \right]_0^1$$

$$\Rightarrow I = \frac{\pi}{\cos \alpha} \left(\frac{\pi}{2} - \alpha \right) \quad \frac{1}{2}$$

11. $I = \int (2x + 5) \sqrt{10 - 4x - 3x^2} dx$

$$= -\frac{1}{3} \int (-4 - 6x) \sqrt{10 - 4x - 3x^2} dx + \frac{11}{3} \int \sqrt{10 - 4x - 3x^2} dx \quad 1$$

$$= -\frac{2}{9} (10 - 4x - 3x^2)^{3/2} + \frac{11\sqrt{3}}{3} \int \sqrt{\left(\frac{\sqrt{34}}{3}\right)^2 - \left(x - \frac{2}{3}\right)^2} dx \quad 1+1$$

$$= -\frac{2}{9} (10 - 4x - 3x^2)^{3/2} + \frac{11\sqrt{3}}{3} \left[\frac{\left(x - \frac{2}{3}\right) - \sqrt{\left(\frac{\sqrt{34}}{3}\right)^2 - \left(x - \frac{2}{3}\right)^2}}{2} + \frac{17}{9} \sin^{-1} \frac{3x - 2}{\sqrt{34}} \right] + C \quad 1$$

OR

$$x^2 = y \text{ (say)} \quad \frac{1}{2}$$

$$\frac{(y+1)(y+4)}{(y+3)(y-5)} = 1 + \frac{A}{y+3} + \frac{B}{y-5} \quad \frac{1}{2}$$

$$\text{using partial fraction we get } A = \frac{1}{4}, B = \frac{27}{4} \quad 1$$

$$\int \frac{(x^2 + 1)(x^2 + 4)}{(x^2 + 3)(x^2 - 5)} dx = \int 1 dx + \frac{1}{4} \int \frac{dx}{x^2 + 3} + \frac{27}{4} \int \frac{dx}{x^2 - 5} \quad 1$$

$$= x + \frac{1}{4\sqrt{3}} \tan^{-1} \frac{x}{\sqrt{3}} + \frac{27}{8\sqrt{5}} \log \left| \frac{x - \sqrt{5}}{x + \sqrt{5}} \right| + C \quad 1$$

12. $I = \int \frac{x \sin^{-1} x}{\sqrt{1-x^2}} dx$

put $\sin^{-1} x = t \Rightarrow \frac{dx}{\sqrt{1-x^2}} = dt$

$$\frac{1}{2} + \frac{1}{2}$$

$$= \int t \cdot \sin t dt$$

$$1$$

$$= -t \cos t + \sin t + c$$

$$1\frac{1}{2}$$

$$= -\sqrt{1-x^2} \sin^{-1} x + x + c$$

$$\frac{1}{2}$$

13. $y^2 dx + (x^2 - xy + y^2) dy = 0$

$$\Rightarrow \frac{dx}{dy} = -\frac{(x^2 - xy + y^2)}{y^2}$$

$$\frac{1}{2}$$

put $x = vy \Rightarrow \frac{dx}{dy} = v + y \frac{dv}{dy}$

$$\frac{1}{2}$$

$$v + y \frac{dv}{dy} = \frac{(v^2 y^2 - y^2 v + y^2)}{y^2}$$

$$\Rightarrow \frac{dv}{v^2 + 1} = -\frac{dy}{y}$$

$$1\frac{1}{2}$$

Integrating both sides

$$\tan^{-1} v = -\log y + c$$

$$\frac{1}{2} + \frac{1}{2}$$

$$\Rightarrow \tan^{-1} \frac{x}{y} = -\log y + c$$

$$\frac{1}{2}$$

14. $\frac{dx}{dy} - \frac{x}{1+y^2} = \frac{\cot^{-1} y}{1+y^2}$

$$\frac{1}{2}$$

$$I.F = e^{-\int \frac{1}{1+y^2} dy} = e^{\cot^{-1} y}$$

$$1$$

$$\Rightarrow \frac{d}{dy} (x \cdot e^{\cot^{-1} y}) = \frac{\cot^{-1} t e^{\cot^{-1} y}}{1+y^2}$$

Integrating, we get

$$x \cdot e^{\cot^{-1} y} = \int \frac{\cot^{-1} y e^{\cot^{-1} y}}{1+y^2} dy$$

$$1\frac{1}{2}$$

put $\cot^{-1} y = t$

$$= - \int t e^t dt$$

$$= (1-t) e^t + c$$

$$\Rightarrow x = (1 - \cot^{-1} y) + c e^{-\cot^{-1} y}$$

$$1$$

15. $\vec{a} \times \vec{b} = \vec{c} \times \vec{d}$... (i) 2
 $\vec{a} \times \vec{c} = \vec{b} \times \vec{d}$... (ii) 1
- (1) - (2) $\Rightarrow \vec{a} \times (\vec{b} - \vec{c}) = \vec{d} \times (\vec{b} - \vec{c})$ 1
- $\Rightarrow (\vec{a} - \vec{d}) \times (\vec{b} - \vec{c}) = \vec{0}$ 1
- $\Rightarrow (\vec{a} - \vec{d}) \parallel (\vec{b} - \vec{c})$ 1

16. Equation of line \overline{AB}
 $\vec{r} = (-\hat{j} - \hat{k}) + \lambda (4\hat{i} + 6\hat{j} + 2\hat{k})$ 1
- Equation of line \overline{CD}
 $\vec{r} = (3\hat{i} + 9\hat{j} + 4\hat{k}) + \mu (-7\hat{i} - 5\hat{j})$ 1
- $\vec{a}_2 - \vec{a}_1 = 3\hat{i} + 10\hat{j} + 5\hat{k}$ 1
- $\vec{b}_1 \times \vec{b}_2 = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 4 & 6 & 2 \\ -7 & -5 & 0 \end{vmatrix} = 10\hat{i} - 14\hat{j} + 22\hat{k}$ 1
- $(\vec{a}_2 - \vec{a}_1) \cdot (\vec{b}_1 \times \vec{b}_2) = 30 - 140 + 110 = 0$ 1
- \Rightarrow Lines intersect

17. Let selection of defective pen be considered success
 $p = \frac{2}{20} = \frac{1}{10}, q = \frac{9}{10}$ 1
- Reqd probability = $P(x=0) + P(x=1) + P(x=2)$ 1
- $= {}^5C_0 \left(\frac{1}{10}\right)^0 \left(\frac{9}{10}\right)^5 + {}^5C_1 \left(\frac{1}{10}\right)^1 \left(\frac{9}{10}\right)^4 + {}^5C_2 \left(\frac{1}{10}\right)^2 \left(\frac{9}{10}\right)^3$
- $= \left(\frac{9}{10}\right)^5 + \frac{1}{2} \left(\frac{9}{10}\right)^4 + \frac{1}{10} \times \left(\frac{9}{10}\right)^3$
- $= \left(\frac{9}{10}\right)^3 \times \frac{34}{25}$ 1

OR

- $\sum_{i=0}^4 P(x_i) = 1$ 1
- $\Rightarrow 8k = 1 \Rightarrow k = \frac{1}{8}$ 1
- (i) $P(x=1) = \frac{1}{8}$ 1

(ii) $P(\text{at most 2 colleges}) = P(0) + P(1) + P(2)$

$$= \frac{5}{8}$$

1

(iii) $P(\text{at least 2 colleges}) = 1 - [P(x=0) + P(x=1)]$

$$= 1 - \frac{1}{8} = \frac{7}{8}$$

1

18. $\text{LHS} = \cot^{-1} \left[\frac{\left(\cos \frac{x}{2} + \sin \frac{x}{2} \right) + \left(\cos \frac{x}{2} - \sin \frac{x}{2} \right)}{\left(\cos \frac{x}{2} + \sin \frac{x}{2} \right) - \left(\cos \frac{x}{2} - \sin \frac{x}{2} \right)} \right]$

1+1

$$= \cot^{-1} \left(\cot \frac{x}{2} \right)$$

1

$$= \frac{x}{2} = \text{RHS}$$

1

OR

$$\tan^{-1} \left[\frac{\frac{x-2}{x-1} + \frac{x+2}{x+1}}{1 - \frac{x-2}{x-1} \cdot \frac{x+2}{x+1}} \right] = \frac{\pi}{4}$$

1 $\frac{1}{2}$

$$\Rightarrow \frac{2x^2 - 4}{3} = \tan \frac{\pi}{4}$$

1 $\frac{1}{2}$

$$\Rightarrow x = \pm \sqrt{\frac{7}{2}}$$

1

19. Let each poor child pay ₹ x per month and each rich child pay ₹ y per month.

$$\therefore 20x + 5y = 9000$$

1 $\frac{1}{2}$

$$5x + 25y = 26000$$

1 $\frac{1}{2}$

In matrix form,

$$\begin{bmatrix} 20 & 5 \\ 5 & 25 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 9000 \\ 26000 \end{bmatrix}$$

1

$$AX = B \Rightarrow X = A^{-1} B$$

$$A^{-1} = \frac{1}{475} \begin{bmatrix} 25 & -5 \\ -5 & 20 \end{bmatrix}$$

$$\begin{bmatrix} x \\ y \end{bmatrix} = \frac{1}{475} \begin{bmatrix} 25 & -5 \\ -5 & 20 \end{bmatrix} \begin{bmatrix} 9000 \\ 26000 \end{bmatrix} = \begin{bmatrix} 200 \\ 1000 \end{bmatrix}$$

$$\Rightarrow x = 200, y = 1000$$


1

Value: Compassion or any relevant value

1

SECTION C

- 20.** Their point of intersection $(1, \sqrt{3})$



Correct Figure

1

$$\text{Required Area} = \int_0^1 \sqrt{(2)^2 - (x-2)^2} dx + \int_1^2 \sqrt{2^2 - x^2} dx$$

2

$$= \left[\frac{(x-2)\sqrt{4x-x^2}}{2} + 2\sin^{-1}\frac{x-2}{2} \right]_0^1 + \left[\frac{x\sqrt{4-x^2}}{2} + 2\sin^{-1}\frac{x}{2} \right]_1^2$$

1

$$= \left(\frac{5\pi}{3} - \sqrt{3} \right) \text{ Sq. units}$$

1

- 21.** Equation of family of planes passing through two given planes

$$(x + 2y + 3z - 4) + k(2x + y - z + 5) = 0$$

1

$$\Rightarrow (1 + 2k)x + (2 + k)y + (3 - k)z = 4 - 5k \quad \dots(i)$$

$$\Rightarrow \frac{x}{4-5k} + \frac{y}{4-5k} + \frac{z}{4-5k} = 1$$

$$\frac{1+2k}{4-5k} + \frac{2+k}{4-5k} + \frac{3-k}{4-5k} = 1$$

1

As per condition

$$\frac{4-5k}{1+2k} = \frac{2(4-5k)}{(3-k)}$$

1

$$\Rightarrow k = \frac{4}{5} \text{ or } \frac{1}{5}$$

1

$$\text{For } k = \frac{1}{5}, \text{ Eqn. of plane is } 7x + 11y + 14z = 15$$

1

$$\text{For } k = \frac{4}{5}, \text{ Eqn. of plane is } 13x + 14y + 11z = 0$$

1
2

Equation of plane passing through $(2, 3, -1)$

1

and parallel to the plane is:

$$7(x-2) + 11(y-3) + 14(z+1) = 0$$

1

$$\Rightarrow 7x + 11y + 14z = 33$$

$$\text{Vector form: } \vec{r} \cdot (7\hat{i} + 11\hat{j} + 14\hat{k}) = 33$$

1
2

- 22.** Let H_1 be the event 2 red balls are transferred

H_2 be the event 1 red and 1 black ball, transferred

H_3 be the event 2 black and 1 black ball transferred

E be the event that ball drawn from B is red.

1

$$P(H_1) = \frac{^3C_2}{^8C_2} = \frac{3}{28} \quad P(E/H_1) = \frac{6}{10}$$

$$P(H_2) = \frac{^3C_1 \times ^5C_1}{^8C_2} = \frac{15}{28} \quad P(E/H_2) = \frac{5}{10}$$

$$P(H_3) = \frac{^5C_2}{^8C_2} = \frac{10}{28}$$

$$P(E/H_3) = \frac{4}{10}$$

$$1\frac{1}{2} + 1\frac{1}{2}$$

$$P(H_1/E) = \frac{\frac{3}{28} \times \frac{6}{10}}{\frac{3}{28} \times \frac{6}{10} + \frac{15}{28} \times \frac{5}{10} + \frac{10}{28} \times \frac{4}{10}}$$

$$= \frac{18}{133}$$

$$1\frac{1}{2}$$

23.


Let x tablets of type X and y tablets of type Y are taken

$$\text{Minimise } C = 2x + y$$

1

subjected to

$$\left. \begin{array}{l} 6x + 2y \geq 18 \\ 3x + 3y \geq 21 \\ 2x + 4y \geq 16 \\ x, y \geq 0 \end{array} \right\}$$



Correct Graph

$$1\frac{1}{2}$$

$$C|_{A(0, 9)} = 9$$

$$C|_{B(1, 6)} = 8 \leftarrow \text{Minimum value}$$

$$C|_{C(6, 1)} = 13$$

$$C|_{D(8, 0)} = 16$$

$$2x + y < 8 \text{ does not pass through unbounded region}$$

$$\frac{1}{2}$$

Thus, minimum value of $C = 8$ at $x = 1, y = 6$.

1

$$24. \quad f(x) = |x| + x, \quad g(x) = |x| - x \quad \forall x \in \mathbb{R}$$

$$(fog)(x) = f(g(x))$$

1

$$= ||x| - 1| + |x| - x$$

1

$$(gof)(x) = g(f(x))$$

$$\frac{1}{2}$$

$$= ||x| + x| - |x| - x$$

$$\frac{1}{2}$$

$$(fog)(-3) = 6$$

1

$$(fog)(5) = 0$$

1

$$(gof)(-2) = 2$$

1

$$25. \quad abc \begin{vmatrix} \frac{1}{a} + 1 & \frac{1}{b} & \frac{1}{c} \\ \frac{1}{a} & \frac{1}{b} + 1 & \frac{1}{c} \\ \frac{1}{a} & \frac{1}{b} & \frac{1}{c} + 1 \end{vmatrix} = 0$$

$$1\frac{1}{2}$$

$$C_1 \rightarrow C_1 + C_2 + C_3$$

$$\Rightarrow abc \begin{vmatrix} 1 + \frac{1}{a} + \frac{1}{b} + \frac{1}{c} & \frac{1}{b} & \frac{1}{c} \\ 1 + \frac{1}{a} + \frac{1}{b} + \frac{1}{c} & \frac{1}{b} + 1 & \frac{1}{c} \\ 1 + \frac{1}{a} + \frac{1}{b} + \frac{1}{c} & \frac{1}{b} & \frac{1}{c} + 1 \end{vmatrix} = 0 \quad 1\frac{1}{2}$$

$$\Rightarrow abc \left(1 + \frac{1}{a} + \frac{1}{b} + \frac{1}{c} \right) \begin{vmatrix} 1 & \frac{1}{b} & \frac{1}{c} \\ 1 & \frac{1}{b} + 1 & \frac{1}{c} \\ 1 & \frac{1}{b} & \frac{1}{c} + 1 \end{vmatrix} = 0$$

$$R_2 \rightarrow R_2 - R_1, R_3 \rightarrow R_3 - R_1 \quad 1\frac{1}{2}$$

$$\Rightarrow abc \left(1 + \frac{1}{a} + \frac{1}{b} + \frac{1}{c} \right) = 0$$

$$\therefore a, b, c, \neq 0$$

$$\therefore 1 + \frac{1}{a} + \frac{1}{b} + \frac{1}{c} = 0 \quad 1\frac{1}{2}$$

OR


$$|A| = 1 \quad 1$$

$$\text{adj } A = \begin{bmatrix} \cos \alpha & \sin \alpha & 0 \\ -\sin \alpha & \cos \alpha & 0 \\ 0 & 0 & 1 \end{bmatrix} \quad 3$$

$$A(\text{adj } A) = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} = I \quad 1$$

$$|A| I_3 = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} = I \quad 1$$

$$26. \quad S = 6x^2 + 4\pi r^2$$



$$\Rightarrow r = \sqrt{\frac{S - 6x^2}{4\pi}} \quad \dots(i) \quad \frac{1}{2}$$

$$V = \frac{2x^3}{3} + \frac{4}{3}\pi r^3 \quad 1$$

$$= \frac{2}{3}x^3 + \frac{4}{3}\pi \left(\frac{S - 6x^2}{4\pi} \right)^{\frac{3}{2}}$$

$$= \frac{2x^3}{3} + \frac{(S - 6x^2)^{\frac{3}{2}}}{6\sqrt{\pi}} \quad 1$$

$$\frac{dV}{dx} = 2x^2 - \frac{3x}{\sqrt{\pi}} \sqrt{S - 6x^2}$$

1

$$\frac{dV}{dx} = 0 \Rightarrow 2x^2 = \frac{3x \sqrt{S - 6x^2}}{\sqrt{\pi}}$$

$$\Rightarrow r = \frac{x}{3} \text{ [using (i)]}$$

1

$$\left. \begin{aligned} \frac{d^2V}{dx^2} &= 4x \left[\frac{3x}{\sqrt{\pi}} \cdot \frac{(-12x)}{2\sqrt{S - 6x^2}} + \frac{3}{\sqrt{\pi}} \sqrt{S - 6x^2} \right] \\ \frac{d^2V}{dx^2} \Big|_{x=\frac{r}{3}} &> 0 \end{aligned} \right\}$$

1

$$\Rightarrow V \text{ is minimum at } x = \frac{r}{3} \text{ i.e. } r = 3x$$

$$\text{Minimum value of sum of volume} = \left(\frac{2x^3}{3} + 36\pi x^3 \right) \text{ cubic units}$$

1
2**OR**

Equation of given curve

$$y = \cos(x + y) \quad \dots(i)$$

$$\Rightarrow \frac{dy}{dx} = -\sin(x + y) \left(1 + \frac{dy}{dx} \right)$$

$$\Rightarrow \frac{dy}{dx} = \frac{-\sin(x + y)}{1 + \sin(x + y)} \quad 2$$

$$\text{given line } x + 2y = 0, \text{ its slope} = -\frac{1}{2} \quad \frac{1}{2}$$

condition of || lines

$$\frac{-\sin(x + y)}{1 + \sin(x + y)} = -\frac{1}{2} \quad 1$$

$$\Rightarrow \sin(x + y) = 1$$

$$\Rightarrow \cos(x + y) = 0 \quad y = 0 \quad \text{using (i)}$$

$$\Rightarrow \cos x = 0 \Rightarrow x = (2n+1)\frac{\pi}{2}, n \in I$$

$$\therefore x = \frac{-3\pi}{2}, \frac{\pi}{2} \in [-2\pi, 2\pi] \quad 1$$

Thus tangents are || to the line $x + 2y = 0$

$$\text{only at pts } \left(-\frac{3\pi}{2}, 0 \right) \text{ and } \left(\frac{\pi}{2}, 0 \right) \quad \frac{1}{2}$$

 \therefore Required equation of tangents are

$$y - 0 = -\frac{1}{2} \left(x + \frac{3\pi}{2} \right) \Rightarrow 2x + 4y + 3\pi = 0 \quad \frac{1}{2}$$

$$y - 0 = -\frac{1}{2} \left(x - \frac{\pi}{2} \right) \Rightarrow 2x + 4y - \pi = 0 \quad \frac{1}{2}$$