

ANON'S ACADEMY FOR MATHS

SECOND TERMINAL EXAMINATION - 2022 MATHAMATICS (SCIENCE) HSE II - ANSWERS

	QUESTIONS FROM 1 TO 8. 3 MARK	MARK
1 (a)	(i) y	1
(b)	$f(x) = x^2 f(x_1) = x_1^2 \quad f(x_2) = x_2^2$ $f(x_1) = f(x_2)$ $x_1^2 = x_2^2 \Rightarrow x_1 = x_2 \text{ or } x_1 = -x_2 \text{ it is not one-one}$ $\text{let } f(x) = y$ $x^2 = y$ $x = \pm\sqrt{y} \text{ y is real number so can be negative too hence } x^2 \text{ is not onto}$	2
2 (a)	(iii) $\frac{\pi}{4}$	1
(b)	$\cos^{-1}\left(\frac{1}{2}\right) + 2\sin^{-1}\left(\frac{1}{2}\right) = \frac{\pi}{3} + 2 \times \frac{\pi}{6}$ $= \frac{\pi}{3} + \frac{\pi}{3} = \frac{2\pi}{3}$	2
3 (a)	$a_{ij} = 2i - j$ $A = \begin{bmatrix} 1 & 0 \\ 3 & 2 \end{bmatrix}$	1
(b)	$AB = \begin{bmatrix} 1 & 0 \\ 3 & 2 \end{bmatrix} \begin{bmatrix} 1 & 2 \\ 2 & 0 \end{bmatrix} = \begin{bmatrix} 1 & 2 \\ 7 & 6 \end{bmatrix}$	2
4	<p>Write equation as $AX = B$</p> $\begin{bmatrix} 2 & 5 \\ 3 & 2 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 1 \\ 7 \end{bmatrix}$ <p>calculate $A = \begin{bmatrix} 2 & 5 \\ 3 & 2 \end{bmatrix} = -11$</p> $AX = B$ $X = A^{-1}B$ <p>Here $A^{-1} = \frac{1}{ A } adj(A)$ $adj(A) = \begin{bmatrix} 2 & -5 \\ -3 & 2 \end{bmatrix}$</p> <p>Now $A^{-1} = \frac{1}{-11} \begin{bmatrix} 2 & -5 \\ -3 & 2 \end{bmatrix}$</p> $X = A^{-1}B \begin{bmatrix} x \\ y \end{bmatrix} = \frac{-1}{11} \begin{bmatrix} 2 & -5 \\ -3 & 2 \end{bmatrix} \begin{bmatrix} 1 \\ 7 \end{bmatrix} = \begin{bmatrix} 3 \\ -1 \end{bmatrix}$ $x = 3, y = -1$	3

ANON'S ACADEMY FOR MATHS


<p>5 Let x be the length of an edge of cube and V be the volume of the cube. Then, $V = x^3$ \therefore Rate of change of volume w.r.t time $\frac{dV}{dt} = \frac{d(x^3)}{dt} = 3x^2 \frac{dx}{dt}$ <p>It is given that edge of the cube is increasing at the rate of 3 cm/s, so $\frac{dx}{dt} = 3\text{cm/s}$ $\therefore \frac{dV}{dt} = 3x^2(3) = 9x^2\text{cm}^3/\text{s}$</p> <p>thus , when $x = 10\text{cm}$, $\frac{dV}{dt} = 9(10)^2 = 900\text{cm}^3/\text{s}$</p> <p>the volume of the cube is increasing at the rate of 900 cm³/s when the edge is 10 cm long</p> </p>	3
<p>6 $\int 2x\sin(x^2 + 1). dx$ let $u = x^2 + 1$ $\frac{du}{dx} = 2x$ $du = 2x dx$ substituting $\int 2x\sin(x^2 + 1)dx = \int \sin u du$ $= (-\cos u) + c$ putting value of $u = x^2 + 1$ $= -\cos(x^2 + 1) + c$</p>	3
<p>7(a) order = 2 degree = 1</p>	2
<p>(b) 0</p>	1
<p>8(a) $x = 2, y = 3$</p>	1
<p>(b) $drs = (1, 2, 2)$ $dcs = (\frac{1}{3}, \frac{2}{3}, \frac{2}{3})$</p>	2
QUESTIONS FROM 9 TO 16. 4 MARK	
<p>9(a) $2x + 3y = \sin x$ Differentiating both sides w.r.t. x, we obtain $\frac{d}{dx}(2x + 3y) = \frac{d}{dx}(\sin x) \Rightarrow \frac{d}{dx}(2x) + \frac{d}{dx}(3y) = \cos x$ $\Rightarrow 2 + 3 \frac{dy}{dx} = \cos x$ $\Rightarrow 3 \frac{dy}{dx} = \cos x - 2$</p>	2

ANON'S ACADEMY FOR MATHS

	$\therefore \frac{dy}{dx} = \frac{\cos x - 2}{3}$							
(b)	<p>let $y = e^{\sin^{-1}x}$ Differentiating both sides w.r.t. x,</p> $\frac{d(y)}{dx} = \frac{d(e^{\sin^{-1}x})}{dx}$ $\Rightarrow \frac{dy}{dx} = \frac{e^{\sin^{-1}x}}{\sqrt{1-x^2}}$	2						
10(a)	(i) f is increasing in $[a, b]$ if $f'(x) > 0$	1						
(b)	<p>$f'(x) = -\sin x$</p> <p>(i) for each $x \in (0, \pi)$, $\sin x > 0$, $f'(x) < 0$ and so f is decreasing in $(0, \pi)$.</p> <p>(ii) for each $x \in (\pi, 2\pi)$, $\sin x < 0$, $f'(x) > 0$ and so f is increasing in $(\pi, 2\pi)$.</p>	3						
11	$f(x) = 3x^4 + 4x^3 - 12x^2 + 12$ $f'(x) = 12x^3 + 12x^2 - 24x = 12x(x-1)(x+2)$ $f'(x) = 0 \text{ at } x = 0, x = 1 \text{ and } x = -2.$ <p>Now $f''(x) = 36x^2 + 24x - 24 = 12(3x^2 + 2x - 2)$</p> <table border="0" style="width: 100%;"> <tr> <td style="width: 50%;">$f''(0) = -24 < 0$</td> <td>0 is the local maxima</td> </tr> <tr> <td>$f''(1) = 36 > 0$</td> <td></td> </tr> <tr> <td>$f''(-2) = 72 > 0$</td> <td>-2 is the local minima</td> </tr> </table> <p>Therefore, by second derivative test, $x = 0$ is a point of local maxima and local maximum value of f at $x = 0$ is $f(0) = 12$ while $x = 1$ and $x = -2$ are the points of local minima and local minimum values of f at $x = -1$ and -2 are $f(1) = 7$ and $f(-2) = -20$, respectively</p>	$f''(0) = -24 < 0$	0 is the local maxima	$f''(1) = 36 > 0$		$f''(-2) = 72 > 0$	-2 is the local minima	4
$f''(0) = -24 < 0$	0 is the local maxima							
$f''(1) = 36 > 0$								
$f''(-2) = 72 > 0$	-2 is the local minima							
12	$I = \int e^x \sin x dx$ <p>By using integration by parts</p> $I = e^x(-\cos x) - \int e^x(-\cos x)dx$ $= -e^x \cos x + \int e^x(\cos x)dx$ $I = -e^x \cos x + e^x \sin x - \int e^x \sin x dx$ $I = -e^x \cos x + e^x \sin x - I$	4						

ANON'S ACADEMY FOR MATHS

	$2I = e^x(\sin x - \cos x) + c$ $I = \frac{e^x}{2}(\sin x - \cos x) + c$	
13(a)	(iv) $\int_a^b f(x) dx$	1
(b)	$\begin{aligned} \text{Area} &= \int_1^4 y dx = \int_1^4 \sqrt{x} dx \\ &= \frac{2}{3} \left[(4)^{\frac{3}{2}} - (1)^{\frac{3}{2}} \right] \\ &= \frac{2}{3} [8 - 1] = \frac{14}{3} \text{ units} \end{aligned}$	3
14(a)	integrating factor = $e^{\int pdx}$	1
(b)	$\begin{aligned} x \cdot \frac{dy}{dx} + 2y &= x^2 \\ p &= \frac{2}{x}, Q = x \\ \int pdx &= \int \frac{2}{x} dx = 2\log x \\ IF &= e^{\int pdx} = e^{2\log x} = e^{\log x^2} = x^2 \end{aligned}$	1
(c)	$\begin{aligned} y \cdot If &= \int Q \cdot If dx + c \\ y \cdot x^2 &= \int x \cdot x^2 dx + c \\ x^2 y &= \int x^3 dx + c \\ x^2 y &= \frac{x^4}{4} + c. \end{aligned}$	2
15(a)	$\vec{a} \cdot \vec{b} = 10$	1
(b)	$\begin{aligned} \cos \theta &= \frac{\vec{a} \cdot \vec{b}}{ \vec{a} \vec{b} } = \frac{10}{\sqrt{14} \sqrt{14}} = \frac{5}{7} \\ \theta &= \cos^{-1} \left(\frac{5}{7} \right) \end{aligned}$	2



ANON'S ACADEMY FOR MATHS

(c)	$projection = \frac{10}{\sqrt{14}}$	1
16(a)	(1,0,0)	1
(b)	$dc's = \left(\frac{3}{\sqrt{77}}, \frac{-2}{\sqrt{77}}, \frac{8}{\sqrt{77}} \right)$	3
QUESTIONS FROM 17 TO 20 . 6 MARK		
17(a)	$\int \frac{1}{\sqrt{7-6x-x^2}} dx = \int \frac{1}{\sqrt{-x^2-6x+7}} dx$ $= \int \frac{1}{\sqrt{-(x^2+6x-7)}} dx$ $= \int \frac{1}{\sqrt{-((x+3)^2-4^2)}} dx$ $= \int \frac{1}{\sqrt{4^2-(x+3)^2}} dx = \sin^{-1} \left(\frac{x+3}{4} \right) + C$	3
(b)	$\int_1^2 \frac{x \cdot dx}{(x+1)(x+2)}$ $\frac{x \cdot dx}{(x+1)(x+2)} = \frac{A}{x+1} + \frac{B}{x+2} \quad (A = -1, B = 2)$ $\frac{x \cdot dx}{(x+1)(x+2)} = \frac{-1}{x+1} + \frac{2}{x+2}$ $\int \frac{x \cdot dx}{(x+1)(x+2)} = -\log x+1 + 2\log x+2 $ $\int_1^2 \frac{x \cdot dx}{(x+1)(x+2)} = -3\log 3 + \log 2 + 2\log 4$	3
18(a)	$\frac{dy}{dx} = \frac{x+y}{x}$ <p>Let $x = \lambda x, y = \lambda y$</p> $\therefore \frac{dy}{dx} = \frac{\lambda x + \lambda y}{\lambda x} = \frac{\lambda(x+y)}{\lambda x} = \frac{x+y}{x}$ <p>\therefore Homogeneous differential eq.</p>	2

ANON'S ACADEMY FOR MATHS

(b)

$$\text{Now let } y = vx \Rightarrow \frac{dy}{dx} = v + x \frac{dv}{dx}$$

$$\therefore v + x \frac{dv}{dx} = \frac{x + vx}{x} 1 + v$$

$$\Rightarrow v + x \frac{dv}{dx} = 1 + v$$

$$\Rightarrow dv = \frac{dx}{x}$$

$$\Rightarrow \int dv = \int \frac{dx}{x}$$

$$\Rightarrow v = \log x + c$$

$$\Rightarrow \frac{y}{x} = \log x + c$$

$$\Rightarrow y = x \log x + c.$$

4

19(a)

$$\hat{i} \times \hat{j} = \hat{k}$$

1

(b)(i)

$$\vec{P} = \vec{a} + \vec{b} = 2\hat{i} + 3\hat{j} + 4\hat{k}$$

2

$$\vec{Q} = \vec{a} - \vec{b} = 0\hat{i} - \hat{j} - 2\hat{k}$$

(ii)

$$\text{unit vector} = \frac{\vec{P} \times \vec{Q}}{|\vec{P} \times \vec{Q}|} = \frac{-2\hat{i} + 4\hat{j} - 2\hat{k}}{\sqrt{24}}$$

3

**20(a) Deleted portion questions
(passing through two points) any related attempt full score**

20(a)

$$\vec{r} = -1\hat{i} + 0\hat{j} + 2\hat{k} + \lambda(4\hat{j} + 4\hat{j} + 4\hat{k})$$

2

(b)

$$S.D = \left| \frac{(\vec{b}_1 \times \vec{b}_2) \cdot (\vec{a}_2 - \vec{a}_1)}{|\vec{b}_1 \times \vec{b}_2|} \right|$$

4

$$\vec{b}_1 \times \vec{b}_2 = 8\hat{i} + 4\hat{j} - 12\hat{k}$$

$$(\vec{b}_1 \times \vec{b}_2) \cdot (\vec{a}_2 - \vec{a}_1) = 44$$

$$|\vec{b}_1 \times \vec{b}_2| = \sqrt{224}$$

$$S.D = \left| \frac{44}{\sqrt{224}} \right|$$